

NUCLEIC ACID MOLECULES AND OTHER MOLECULES ASSOCIATED WITH
THE TOCOPHEROL PATHWAY

in D15
CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority under 35 U.S.C. § 119(e) to application serial no.

5 60/078,031, which is herein incorporated by reference in its entirety.

FIELD OF THE INVENTION

The present invention is in the field of plant biochemistry. More specifically the invention relates to nucleic acid sequences from plant cells, in particular, nucleic acid sequences from maize and soybean plants associated with the tocopherol synthesis pathway in plants. The invention encompasses nucleic acid molecules that encode proteins and fragments of proteins. In addition, the invention also encompasses proteins and fragments of proteins so encoded and antibodies capable of binding these proteins or fragments. The invention also relates to methods of using the nucleic acid molecules, proteins and fragments of proteins and antibodies, for example for genome mapping, gene identification and analysis, plant breeding, preparation of constructs for use in plant gene expression and transgenic plants.

BACKGROUND OF THE INVENTION

I. TOCOPHEROL SYNTHESIS PATHWAY

The chloroplast of higher plants exhibit interconnected biochemical pathways that lead to secondary metabolites, including tocopherols, that not only perform functions in plants but can also be important for mammalian nutrition. In plastids, tocopherols account up to 40% of the total quinone pool. The biosynthetic pathway of α -tocopherol in higher plants involves condensation of homogentisic acid and phytylpyrophosphate to form 2-methyl-6

phytylbenzoquinol (Fiedler *et al.*, *Planta* 155: 511-515 (1982); Soll *et al.*, *Arch. Biochem.*

Biophys. 204: 544-550 (1980); Marshall *et al.*, *Phytochem.* 24: 1705-1711 (1985), all of which

are herein incorporated by reference in their entirety). The plant tocopherol biosynthetic

pathway can be divided into four parts: synthesis of homogentisic acid, which contributes to the

5 aromatic ring of tocopherol; synthesis of phytylpyrophosphate, which contributes to the side chain of tocopherol; cyclization which plays a role in chirality and chromanol substructure of the vitamin E family; and S-adenosyl methionine dependent methylation of an aromatic ring, which effects the compositional quality of the vitamin E family.

Homogentisate is an aromatic precursor in the biosynthesis of tocopherols in chloroplasts and is formed from the aromatic shikimate metabolite, p-hydroxyphenylpyruvate. The aromatic amino acids phenylalanine, tyrosine, and tryptophan are formed by a reaction sequence that initiates from the two carbohydrate precursors, D-erythrose 4-phosphate and phosphoenolpyruvate, via shikimate, and forms prearomatic and aromatic compounds (Bentley, *Critical Rev. Biochem. Mol. Biol.* 25: 307-384 (1990), the entirety of which is herein incorporated by reference). Approximately 20% of the total carbon fixed by green plants is routed through the shikimate pathway with end products being aromatic amino acids and other aromatic secondary metabolites such as flavonoids, vitamins, lignins, alkaloids, and phenolics (Herrmann, *Plant Physiol.* 107: 7-12 (1995), Kishore and Shah, *Ann. Rev. Biochem.* 57:67-663 (1988), both of which are herein incorporated by reference in their entirety). Various aspects of
20 the shikimate pathway have been reviewed (Bentley, *Critical Rev. Biochem. Mol. Biol.* 25: 307-384 (1990); Herrmann, *Plant Physiol.* 107: 7-12 (1995); Kishore and Shah, *Ann. Rev. Biochem.* 57:67-663 (1988)).

The first reported committed reaction in the shikimate pathway is catalyzed by the enzyme 3-deoxyarabino-heptulosonate 7-phosphate synthase (also known as 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase, deoxyarabino-heptulosonate-P-synthase, and DAHP synthase (EC. 4.1.2.15)), which has been reported to control carbon flow into the shikimate pathway. The plastid localized DAHP synthase catalyzes the formation of 3-deoxy-D-arabino-heptulosonate 7-phosphate by condensing D-erythrose 4-phosphate with phosphoenolpyruvate. DAHP synthase has been isolated from plant sources including carrot and potato. DAHP synthase has substrate specificity for D-erythrose 4-phosphate and phosphoenolpyruvate, is a dimer of subunits having a molecular weight of 53KD and is activated by Mn^{2+} (Herrmann, *Plant Physiol.* 107: 7-12 (1995)). Aromatic amino acids are not reported to act as feedback regulators. Purified DAHP synthase is activated by tryptophan and, to a lesser extent, by tyrosine in a hysteric fashion (Suzich *et al.*, *Plant Physiol.* 79: 765-770 (1985), the entirety of which is herein incorporated by reference).

The next reported enzyme in the shikimate pathway is 3-dehydroquinate synthase (EC 4.6.1.3), which catalyzes the formation of dehydroquinate, the first carbocyclic metabolite in the biosynthesis of aromatic amino acids, from the substrates D-erythrose 4-phosphate and phosphoenolpyruvate. The enzyme reaction involves a NAD (nicotinamide adenine dinucleotide) cofactor dependent oxidation-reduction, β -elimination, and an intramolecular aldol condensation. 3-dehydroquinate synthase has been purified from *Phaseolus mungo* seedlings and pea seedlings, has a native molecular weight of 66 KD and is a dimer (Yamamoto, *Phytochem.* 19: 779-802 (1980); Pompliano *et al.*, *J. Am. Chem. Soc.* 111: 1866-1871-1871 (1989), both of which are herein incorporated by reference in their entirety).

3-dehydroquinate dehydratase (EC 4.2.1.10) catalyzes the stereospecific syn-dehydration of dehydroquinate to dehydroshikimate and has been reported to be responsible for initiating the process of aromatization by introducing the first of three double bonds of the aromatic ring system. 3-dehydroquinate dehydratase has been cloned from *E. coli* (Duncan *et al.*, *Biochem. J.* 238:475-483 (1986), the entirety of which is herein incorporated by reference).

Shikimate dehydrogenase (EC 1.1.1.25) catalyzes the NADPH (reduced nicotinamide adenine dinucleotide phosphate)-dependent conversion of dehydroshikimate to shikimate. Bifunctional 3-dehydroquinate dehydratase-shikimate dehydrogenase has been reported in spinach, pea seedling, and Maize (Bentley, *Critical Rev. Biochem. Mol. Biol.* 25: 307-384 (1990), Kishore and Shah, *Ann. Rev. Biochem.* 57:67-663 (1988)). *E. coli* shikimate dehydrogenase has been reported to be a monomeric, monofunctional protein with a molecular weight of 32,000 daltons (Chaudhuri and Coggins, *Biochem. J.* 226:217-223 (1985), the entirety of which is herein incorporated by reference).

Shikimate kinase (EC 2.7.1.71) catalyzes the phosphorylation of shikimate to shikimate-3-phosphate. Shikimate kinase exists as isoforms in *E. coli* and *S. typhimurium*. Plant shikimate kinase has been partially purified from mung bean and sorghum (Bentley, *Critical Rev. Biochem. Mol. Biol.* 25: 307-384 (1990); Kishore and Shah, *Ann. Rev. Biochem.* 57:67-663 (1988)).

Certain plant species accumulate shikimate and shikimate kinase may play a role in regulating flux in the tocopherol synthesis pathway.

5-Enolpyruvyl-shikimate-3-phosphate synthase (also known as enolpyruvyl-shikimate-P-synthase, and EPSPS (EC 2.5.1.19)) catalyzes the reversible transfer of the carboxyvinyl moiety of phosphoenolpyruvate to shikimate-3-phosphate, yielding 5-enolpyruvyl-shikimate-3-phosphate. 5-Enolpyruvyl-shikimate-3-phosphate synthase is a target of the broad spectrum,

nonselective, postemergence herbicide, glyphosate. Chemical modification studies indicate that lysine, arginine, and histidine residues are essential for activity of the enzyme (Kishore and Shah, *Ann. Rev. Biochem.* 57:67-663 (1988)). 5-Enolpyruvyl-shikimate-3-phosphate synthase has been isolated and characterized from microbial and plant sources including tomato, petunia,
5 *Arabidopsis*, and *Brassica* (Kishore and Shah, *Ann. Rev. Biochem.* 57:67-663 (1988)).

Chorismate synthase (EC 4.6.1.4) catalyzes the conversion of 5-enolpyruvyl-shikimate-3-phosphate to chorismic acid and introduces a second double bond in an aromatic ring and a trans-1,4-elimination of inorganic phosphorous. Chorismate is the last reported common intermediate in the biosynthesis of aromatic compounds via the shikimate pathway. The enzyme reaction involves no change in the oxidation state of the substrate. Chorismate synthase from various sources requires a reduced flavin cofactor, FMNH₂ (reduced flavin mononucleotide) or FADH₂ (reduced flavin adenine dinucleotide), for catalytic activity (Bentley, *Critical Rev. Biochem. Mol. Biol.* 25: 307-384 (1990); Kishore and Shah, *Ann. Rev. Biochem.* 57:67-663 (1988)).

The next reported enzyme in the tocopherol biosynthetic pathway is chorismate mutase (EC 5.4.99.5), which catalyzes the conversion of chorismic acid to prephenic acid. Chorismic acid is a substrate for a number of enzymes involved in the biosynthesis of aromatic compounds. Plant chorismate mutase exists in two isoforms, chorismate mutase-1 and chorismate mutase-2, that differ in feedback regulation by aromatic amino acids (Singh *et al.*, *Arch. Biochem. Biophys.* 243: 374-384 (1985); Goers *et al.*, *Planta* 162: 109-124 (1984), both of which are herein
20 incorporated by reference in their entirety). It has been reported that chloroplastic chorismate mutase-1 may play a role in biosynthesis of aromatic amino acids as this enzyme is activated by tyrosine and phenylalanine. Cytosolic isozyme chorismate mutase-2 is not regulated by aromatic amino acids and may play a role in providing the aromatic nucleus for synthesis of aromatic

secondary metabolites including tocopherol (d'Amato *et al.*, *Planta*, 162: 104-108 (1984), the entirety of which is herein incorporated by reference).

The metabolic pathways branch after prephenic acid and lead not only to phenylalanine and tyrosine, but also to a number of secondary metabolites. Tyrosine is synthesized from prephenate via either 4-hydroxyphenylpyruvate or arogenate. Both routes have been reported in plants (Bentley, *Critical Rev. Biochem. Mol. Biol.* 25: 307-384 (1990)).

The formation of 4-hydroxyphenylpyruvate from prephenate is catalyzed by prephenate dehydrogenase (EC 1.3.1.12 for NAD specific prephenate dehydrogenase and EC 1.3.1.13 for NADP specific prephenate dehydrogenase). 4-Hydroxyphenylpyruvate associated with tocopherol biosynthesis may also come from tyrosine pool by the action of tyrosine transaminase (EC 2.6.1.5) or L-amino acid oxidase (EC 1.4.3.2). Tyrosine transaminase catalyzes the pyridoxal-phosphate dependent conversion of L-tyrosine to 4-hydroxyphenylpyruvate. This reversible enzyme reaction transfers the amino group of tyrosine to 2-oxoglutarate to form 4-hydroxyphenylpyruvate and glutamate. L-amino acid oxidase (EC 1.4.3.2) catalyzes the conversion of tyrosine to 4-hydroxyphenylpyruvate by acting on the amino group of tyrosine with oxygen acting as an acceptor. L-amino acid oxidase is not specific to tyrosine. In *E. coli*, aromatic amino acid amino transferase (also referred to as aromatic-amino-acid transaminase (EC 2.6.1.57)) converts 4-hydroxyphenylpyruvate to tyrosine and plays a role in phenylalanine and tyrosine biosynthesis (Oue *et al.*, *J. Biochem. (Tokyo)* 121: 161-171 (1997); Soto-Urzua *et al.*, *Can. J. Microbiol.* 42: 294-298 (1996); Hayashi *et al.*, *Biochemistry* 32: 12229-12239 (1993), all of which are herein incorporated by reference in their entirety).

Aspartic acid amino transferase or transaminase A (EC 2.6.1.1) exhibits a broad substrate specificity and may utilize phenylpyruvate or p-hydroxyphenylpyruvate to form phenylalanine

and tyrosine, respectively. Transaminase A has been characterized in *Arabidopsis* (Wilkie *et al.*, *Biochem J.* 319: 969-976 (1996); Wilkie *et al.*, *Plant Mol. Biol.* 27: 1227-1233 (1995), both of which are herein incorporated by reference in their entirety), rice (Song *et al.*, *DNA Res.* 3: 303-310 (1996), herein incorporated by reference in its entirety), *Panicum miliaceum* L (Taniguchi *et al.*, *Arch. Biochem. Biophys.* 318: 295-306 (1995), herein incorporated by reference in its entirety), *Lupinus angustifolius* (Winefield *et al.*, *Plant Physiol.* 104: 417-423 (1994), herein incorporated by reference in its entirety), and soybean (Wadsworth *et al.*, *Plant Mol. Biol.* 21: 993-1009 (1993), herein incorporated by reference in its entirety).

A precursor molecule, homogentisic acid, is produced in the chloroplast from the shikimate pathway intermediate p-hydroxyphenylpyruvate. p-Hydroxyphenylpyruvate dioxygenase (also known as 4-hydroxyphenylpyruvate dioxygenase (EC 1.13.11.27)) catalyzes the formation of homogentisate from hydroxyphenylpyruvate through an oxidative decarboxylation of the 2-oxoacid side chain accompanied by hydroxylation of the aromatic ring and a 1,2 migration of the carboxymethyl group. Norris *et al.* reported functional identification of a *pdsI* gene that encodes p-Hydroxyphenylpyruvate dioxygenase (Norris *et al.*, *Plant Cell* 7: 2139-2149 (1995), the entirety of which is herein incorporated by reference). p-Hydroxyphenylpyruvate dioxygenase has been cloned from *Arabidopsis* and carrot (GenBank accession numbers U89267, AF000228, and U87257; Garcia *et al.*, *Biochem. J.* 325: 761-769 (1997), herein incorporated by reference in its entirety). Fiedler *et al.* reported the localization and presence of this enzyme in both isolated spinach chloroplast and the peroxisome (Fiedler *et al.*, *Planta*, 155: 511-515 (1982)). Garcia *et al.* reported the purification of the cytosolic form of hydroxyphenylpyruvate dioxygenase from cultured carrot protoplast (Garcia *et al.*, *Biochem. J.* 325: 761-769 (1997), the entirety of which is herein incorporated by reference). It has been

reported that the chloroplastic isoform may be involved in the biosynthesis of prenylquinones, and that the peroxisomal and cytosolic isoform may be involved in the degradation of tyrosine.

The carbon flow to the pool of phytol, *i.e.*, the isoprene-derived side chain of tocopherol, occurs via the mevalonate pathway or non-mevalonate pathway. Geranylgeranyl-pyrophosphate synthase (GGPP synthase (EC 2.5.1.29)) catalyzes the formation of geranylgeranylpyrophosphate by prenyltransferring an isoprene moiety from isopentenylpyrophosphate to farnesylpyrophosphate. A gene encoding geranylgeranylpyrophosphate synthase has been isolated from *Arabidopsis* and *Cantharanthus roseus* (Zhu *et al.*, *Plant Cell Physiol.* 38: 357-361 (1997), Bantignies *et al.*, *Plant Physiol.* 110: 336-336 (1995), both of which are herein incorporated by reference in their entirety).

Geranylgeranylpyrophosphate synthesized by GGPP synthase is used in the carotenoid and tocopherol biosynthesis pathways.

The NADPH-dependent hydrogenation of geranylgeranylpyrophosphate is catalyzed by geranylgeranylpyrophosphate hydrogenase (also called geranylgeranylpyrophosphate reductase) to form phytylpyrophosphate (Soll *et al.*, *Plant Physiol.* 71: 849-854 (1983), the entirety of which is herein incorporated by reference). Geranylgeranylpyrophosphate hydrogenase appears to be localized to two sites: the chloroplast envelope and the thylakoids. The chloroplast envelope form is reported to be responsible for the hydrogenation of geranylgeranylpyrophosphate to a phytyl moiety. The thylakoids form is reported to be responsible for the stepwise reduction of chlorophyll esterified with geranylgeraniol to chlorophyll esterified with phytol. The chloroplast envelope form of geranylgeranylpyrophosphate may play a role in tocopherol and phylloquinone synthesis. A *chlP* gene cloned from *Synechocystis* has been functionally assigned by complementation in

Rhodobactor sphaeroids to catalyze the stepwise hydrogenation of geranylgeraniol to phytol (Addlesse *et al.*, *FEBS Lett.* 389: 126-130 (1996), the entirety of which is herein incorporated by reference).

Homogentisate:phytyl transferase (also referred to as phytyl/prenyltransferase) catalyzes the decarboxylation followed by condensation of homogentisic acid with a phytol moiety from phytylpyrophosphate to form 2-methyl-6 phytylbenzoquinol. Prenyltransferase activity has been reported in spinach chloroplasts and such activity is located in chloroplast envelope membranes (Fiedler *et al.*, *Planta* 155: 511-515 (1982)). A reported prenyltransferase gene, termed *pdsII*, specific to tocopherol biosynthesis has been identified in *Arabidopsis* (Norris *et al.*, *Plant Cell* 7: 2139-2149 (1995)).

Tocopherol cyclase catalyzes the cyclization of 2,3-dimethyl-6-phytylbenzoquinol to form γ -tocopherol and plays a role in the biosynthesis of enantioselective chromanol substructure of the vitamin E subfamily (Stocker *et al.*, *Bioorg. Medic. Chem.* 4: 1129-1134 (1996), the entirety of which is herein incorporated by reference). The preferred substrate specificity of tocopherol cyclase may be either 2,3-dimethyl-6-phytylbenzoquinol or 2-methyl-5-phytylbenzoquinol or both. The substrate, 2-methyl-6 phytylbenzoquinol, is formed by prenyltransferase and requires methylation by an *S*-adenosylmethionine-dependent methyltransferase before cyclization. Tocopherol cyclase has been purified from green algae *Chlorella protothecoids*, *Dunaliella salina* and from wheat leafs (U.S. Patent No. 5,432,069, the entirety of which is herein incorporated by reference).

Synthesis of γ -tocopherol from 2-methyl-6 phytylbenzoquinol occurs by two pathways with either δ -tocopherol or 2,3 dimethyl-5-phytylbenzoquinol acting as an intermediate. α -

Tocopherol is then synthesized from γ -tocopherol in a final methylation step with S-adenosylmethionine. These steps of α -tocopherol biosynthesis are located in the chloroplast membrane in higher plants. Formation of α -tocopherol from other tocopherols is catalyzed by S-adenosyl methionine (SAM)-dependent γ -tocopherol methyltransferase (EC 2.1.1.95). This enzyme has been partially purified from *Capsicum* and *Euglena gracilis* (Shigeoka *et al.*, *Biochim. Biophys. Acta* 1128: 220-226 (1992), d'Harlingue and Camara, *J. Biol. Chem.* 260: 15200-15203 (1985), both of which are herein incorporated by reference in their entirety).

Tocotrienols are similar to tocopherols in molecular structure except that there are three double bonds in the isoprenoid side chain. Although tocotrienols have not been reported in soybean, they are found within in the plant kingdom. The tocotrienol biosynthetic pathway is similar to that of tocopherol up to the formation of homogentisic. It has been reported that homogentisate:phytyl transferase is able to transfer geranylgeranyl-pyrophosphate ("GGPP") to homogentisic acid. A side chain of GGPP may be desaturated by the addition of phytylpyrophosphate to homogentisate. Stocker *et al.* report that a reduction of the side chain's double bond occurs at an earlier stage of the biosynthesis. Phytylpyrophosphate or GGPP are condensed with homogentisic acid ("HGA") to yield different hydroquinone precursors which are cyclized by the same enzyme (Stocker *et al.*, *Bioorg. Medicinal Chem.* 4:1129-1134 (1996), the entirety of which is herein incorporated by reference).

The primary oxidation product of tocopherol is tocopheryl quinone, which can be conjugated to yield glucuronate after prior reduction to the hydroquinone. In animals, glucuronate can be excreted into bile or further catabolized to tocopheronic acid in the kidney

and processed for urinary excretion (Traber and Sies, *Ann. Rev. Nutr.* 16:321-347 (1996), the entirety of which is herein incorporated by reference).

In *Aspergillus nidulans*, the aromatic amino acid catabolic pathway involves formation of homogentisic acid followed by aromatic ring cleavage by an homogentisic acid dioxygenase (EC 1.13.11.5) to yield, after an isomerization step, fumarylacetoacetate (Fernandez-Canon *et al.*, *Anal. Biochem.* 245: 218-22 (1997); Hudecova *et al.*, *Int. J. Biochem. Cell Biol.* 27: 1357-1363 (1995); Fernandez-Canon *et al.*, *J. Biol. Chem.* 270: 21199-21205 (1995), all of which are herein incorporated by reference in their entirety). Fumarylacetoacetate, is then split by fumarylacetoacetate (Fernandez-Canon and Penalva, *J. Biol. Chem.* 270:21199-21205 (1995), the entirety of which is herein incorporated by reference). Homogentisic acid dioxygenase uses a tocopherol biosynthetic metabolite homogentisic acid for hydrolysis.

Tocopherol levels are reported to vary in different plants, tissues, and developmental stages. The production of homogentisic acid by p-hydroxyphenylpyruvate dioxygenase may be a regulatory point for bulk flow through the pathway due to the irreversible nature of the enzyme reaction and due to the fact that homogentisic acid production is the first committed step in tocopherol biosynthesis (Norris *et al.*, *Plant Cell* 7: 2139-2149 (1995)). Another regulatory step in tocopherol biosynthesis may be associated with the availability of phytylpyrophosphate pool. Feeding studies in Safflower callus culture showed 1.8-fold and 18-fold increase in tocopherol synthesis by feeding homogentisate and phytol, respectively (Fury *et al.*, *Phytochem.* 26: 2741-2747 (1987), the entirety of which is herein incorporated by reference). In meadow rescue leaf, vitamin E increases in the initial phase of senescence when phytol is cleaved off from the chlorophylls and when a free phytol pool is available (Peskier *et al.*, *J. Plant Physiol.* 135: 428-432 (1989), the entirety of which is herein incorporated by reference).

II. EXPRESSED SEQUENCE TAG NUCLEIC ACID MOLECULES

Expressed sequence tags, or ESTs are randomly sequenced members of a cDNA library (or complementary DNA)(McCombie *et al.*, *Nature Genetics* 1:124-130 (1992); Kurata *et al.*, *Nature Genetics* 8: 365-372 (1994); Okubo, *et al.* *Nature Genetics* 2: 173-179 (1992), all of which references are incorporated herein in their entirety). The randomly selected clones comprise insets that can represent a copy of up to the full length of a mRNA transcript.

Using conventional methodologies, cDNA libraries can be constructed from the mRNA (messenger RNA) of a given tissue or organism using poly dT primers and reverse transcriptase (Efstratiadis *et al.* *Cell* 7:279-288 (1976), the entirety of which is herein incorporated by reference; Higuchi *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 73:3146-3150 (1976), the entirety of which is herein incorporated by reference; Maniatis *et al.*, *Cell* 8:163-182 (1976) the entirety of which is herein incorporated by reference; Land *et al.*, *Nucleic Acids Res.* 9:2251-2266 (1981), the entirety of which is herein incorporated by reference; Okayama *et al.*, *Mol. Cell. Biol.* 2:161-170 (1982), the entirety of which is herein incorporated by reference; Gubler *et al.*, *Gene* 25:263-269 (1983), the entirety of which is herein incorporated by reference).

Several methods may be employed to obtain full-length cDNA constructs. For example, terminal transferase can be used to add homopolymeric tails of dC residues to the free 3' hydroxyl groups (Land *et al.*, *Nucleic Acids Res.* 9:2251-2266 (1981), the entirety of which is herein incorporated by reference). This tail can then be hybridized by a poly dG oligo which can act as a primer for the synthesis of full length second strand cDNA. Okayama and Berg, *Mol. Cell. Biol.* 2: 161-170 (1982), the entirety of which is herein incorporated by reference, report a method for obtaining full length cDNA constructs. This method has been simplified by using synthetic primer-adapters that have both homopolymeric tails for priming the synthesis of the

first and second strands and restriction sites for cloning into plasmids (Coleclough *et al.*, *Gene* 34:305-314 (1985), the entirety of which is herein incorporated by reference) and bacteriophage vectors (Krawinkel *et al.*, *Nucleic Acids Res.* 14:1913 (1986), the entirety of which is herein incorporated by reference; Han *et al.*, *Nucleic Acids Res.* 15:6304 (1987), the entirety of which is
5 herein incorporated by reference).

These strategies have been coupled with additional strategies for isolating rare mRNA populations. For example, a typical mammalian cell contains between 10,000 and 30,000 different mRNA sequences (Davidson, *Gene Activity in Early Development*, 2nd ed., Academic Press, New York (1976). The number of clones required to achieve a given probability that a
10 low-abundance mRNA will be present in a cDNA library is $N = (\ln(1-P))/(\ln(1-1/n))$ where N is the number of clones required, P is the probability desired, and 1/n is the fractional proportion of the total mRNA that is represented by a single rare mRNA (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press (1989), the entirety of which is herein incorporated by reference).

A method to enrich preparations of mRNA for sequences of interest is to fractionate by size. One such method is to fractionate by electrophoresis through an agarose gel (Pennica *et al.*, *Nature* 301:214-221 (1983), the entirety of which is herein incorporated by reference). Another such method employs sucrose gradient centrifugation in the presence of an agent, such as methylmercuric hydroxide, that denatures secondary structure in RNA (Schweinfest *et al.*, *Proc.*
15
20 *Natl. Acad. Sci. (U.S.A.)* 79:4997-5000 (1982), the entirety of which is herein incorporated by reference).

A frequently adopted method is to construct equalized or normalized cDNA libraries (Ko, *Nucleic Acids Res.* 18:5705-5711 (1990), the entirety of which is herein incorporated by

reference; Patanjali, S. R. *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 88:1943-1947 (1991), the entirety of which is herein incorporated by reference). Typically, the cDNA population is normalized by subtractive hybridization (Schmid *et al.*, *J. Neurochem.* 48:307-312 (1987) the entirety of which is herein incorporated by reference; Fargnoli *et al.*, *Anal. Biochem.* 187:364-373 (1990) the
5 entirety of which is herein incorporated by reference; Travis *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 85:1696-1700 (1988) the entirety of which is herein incorporated by reference; Kato, *Eur. J. Neurosci.* 2:704-711 (1990); and Schweinfest *et al.*, *Genet. Anal. Tech. Appl.* 7:64-70 (1990), the entirety of which is herein incorporated by reference). Subtraction represents another method for reducing the population of certain sequences in the cDNA library (Swaroop *et al.*, *Nucleic Acids Res.* 19:1954 (1991), the entirety of which is herein incorporated by reference).

ESTs can be sequenced by a number of methods. Two basic methods may be used for DNA sequencing, the chain termination method of Sanger *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 74: 5463-5467 (1977), the entirety of which is herein incorporated by reference, and the chemical degradation method of Maxam and Gilbert, *Proc. Nat. Acad. Sci. (U.S.A.)* 74: 560-564 (1977), the entirety of which is herein incorporated by reference. Automation and advances in technology such as the replacement of radioisotopes with fluorescence-based sequencing have reduced the effort required to sequence DNA (Craxton, *Methods* 2: 20-26 (1991), the entirety of which is herein incorporated by reference; Ju *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 92: 4347-4351 (1995), the entirety of which is herein incorporated by reference; Tabor and Richardson,
20 *Proc. Natl. Acad. Sci. (U.S.A.)* 92: 6339-6343 (1995), the entirety of which is herein incorporated by reference). Automated sequencers are available from, for example, Pharmacia Biotech, Inc., Piscataway, New Jersey (Pharmacia ALF), LI-COR, Inc., Lincoln, Nebraska (LI-COR 4,000) and Millipore, Bedford, Massachusetts (Millipore BaseStation).

In addition, advances in capillary gel electrophoresis have also reduced the effort required to sequence DNA and such advances provide a rapid high resolution approach for sequencing DNA samples (Swerdlow and Gesteland, *Nucleic Acids Res.* 18:1415-1419 (1990); Smith, *Nature* 349:812-813 (1991); Luckey *et al.*, *Methods Enzymol.* 218:154-172 (1993); Lu *et al.*, *J. Chromatog. A.* 680:497-501 (1994); Carson *et al.*, *Anal. Chem.* 65:3219-3226 (1993); Huang *et al.*, *Anal. Chem.* 64:2149-2154 (1992); Kheterpal *et al.*, *Electrophoresis* 17:1852-1859 (1996); Quesada and Zhang, *Electrophoresis* 17:1841-1851 (1996); Baba, *Yakugaku Zasshi* 117:265-281 (1997), all of which are herein incorporated by reference in their entirety).

ESTs longer than 150 nucleotides have been found to be useful for similarity searches and mapping (Adams *et al.*, *Science* 252:1651-1656 (1991), herein incorporated by reference). ESTs, which can represent copies of up to the full length transcript, may be partially or completely sequenced. Between 150-450 nucleotides of sequence information is usually generated as this is length of sequence information that is routinely and reliably produced using single run sequence data. Typically, only single run sequence data is obtained from the cDNA library (Adams *et al.*, *Science* 252:1651-1656 (1991). Automated single run sequencing typically results in an approximately 2-3% error or base ambiguity rate (Boguski *et al.*, *Nature Genetics* 4:332-333 (1993), the entirety of which is herein incorporated by reference).

EST databases have been constructed or partially constructed from, for example, *C. elegans* (McCombie *et al.*, *Nature Genetics* 1:124-131 (1992)), human liver cell line HepG2 (Okubo *et al.*, *Nature Genetics* 2:173-179 (1992)), human brain RNA (Adams *et al.*, *Science* 252:1651-1656 (1991)); Adams *et al.*, *Nature* 355:632-635 (1992)), *Arabidopsis*, (Newman *et al.*, *Plant Physiol.* 106:1241-1255 (1994)); and rice (Kurata *et al.*, *Nature Genetics* 8:365-372 (1994)).

III. SEQUENCE COMPARISONS

A characteristic feature of a protein or DNA sequence is that it can be compared with other known protein or DNA sequences. Sequence comparisons can be undertaken by determining the similarity of the test or query sequence with sequences in publicly available or proprietary databases ("similarity analysis") or by searching for certain motifs ("intrinsic sequence analysis") (e.g. *cis* elements) (Coulson, *Trends in Biotechnology* 12: 76-80 (1994), the entirety of which is herein incorporated by reference); Birren *et al.*, *Genome Analysis I*: Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York 543-559 (1997), the entirety of which is herein incorporated by reference).

Similarity analysis includes database search and alignment. Examples of public databases include the DNA Database of Japan (DDBJ) (<http://www.ddbj.nig.ac.jp/>); Genebank (<http://www.ncbi.nlm.nih.gov/Web/Search/Index.html>); and the European Molecular Biology Laboratory Nucleic Acid Sequence Database (EMBL) (http://www.ebi.ac.uk/ebi_docs/embl_db/embl-db.html). A number of different search algorithms have been developed, one example of which are the suite of programs referred to as BLAST programs. There are five implementations of BLAST, three designed for nucleotide sequences queries (BLASTN, BLASTX, and TBLASTX) and two designed for protein sequence queries (BLASTP and TBLASTN) (Coulson, *Trends in Biotechnology* 12: 76-80 (1994); Birren *et al.*, *Genome Analysis I*: Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York 543-559 (1997)).

BLASTN takes a nucleotide sequence (the query sequence) and its reverse complement and searches them against a nucleotide sequence database. BLASTN was designed for speed, not maximum sensitivity, and may not find distantly related coding sequences. BLASTX takes a

nucleotide sequence, translates it in three forward reading frames and three reverse complement reading frames, and then compares the six translations against a protein sequence database.

BLASTX is useful for sensitive analysis of preliminary (single-pass) sequence data and is tolerant of sequencing errors (Gish and States, *Nature Genetics* 3: 266-272 (1993), the entirety of which is herein incorporated by reference). BLASTN and BLASTX may be used in concert for analyzing EST data (Coulson, *Trends in Biotechnology* 12: 76-80 (1994); Birren *et al.*, *Genome Analysis* 1: 543-559 (1997)).

Given a coding nucleotide sequence and the protein it encodes, it is often preferable to use the protein as the query sequence to search a database because of the greatly increased sensitivity to detect more subtle relationships. This is due to the larger alphabet of proteins (20 amino acids) compared with the alphabet of nucleic acid sequences (4 bases), where it is far easier to obtain a match by chance. In addition, with nucleotide alignments, only a match (positive score) or a mismatch (negative score) is obtained, but with proteins, the presence of conservative amino acid substitutions can be taken into account. Here, a mismatch may yield a positive score if the non-identical residue has physical/chemical properties similar to the one it replaced. Various scoring matrices are used to supply the substitution scores of all possible amino acid pairs. A general purpose scoring system is the BLOSUM62 matrix (Henikoff and Henikoff, *Proteins* 17: 49-61 (1993), the entirety of which is herein incorporated by reference), which is currently the default choice for BLAST programs. BLOSUM62 is tailored for alignments of moderately diverged sequences and thus may not yield the best results under all conditions. Altschul, *J. Mol. Biol.* 36: 290-300 (1993), the entirety of which is herein incorporated by reference, describes a combination of three matrices to cover all contingencies. This may improve sensitivity, but at the expense of slower searches. In practice, a single

BLOSUM62 matrix is often used but others (PAM40 and PAM250) may be attempted when additional analysis is necessary. Low PAM matrices are directed at detecting very strong but localized sequence similarities, whereas high PAM matrices are directed at detecting long but weak alignments between very distantly related sequences.

5 Homologues in other organisms are available that can be used for comparative sequence analysis. Multiple alignments are performed to study similarities and differences in a group of related sequences. CLUSTAL W is a multiple sequence alignment package that performs progressive multiple sequence alignments based on the method of Feng and Doolittle, *J. Mol. Evol.* 25: 351-360 (1987), the entirety of which is herein incorporated by reference. Each pair of sequences is aligned and the distance between each pair is calculated; from this distance matrix, a guide tree is calculated, and all of the sequences are progressively aligned based on this tree. A feature of the program is its sensitivity to the effect of gaps on the alignment; gap penalties are varied to encourage the insertion of gaps in probable loop regions instead of in the middle of structured regions. Users can specify gap penalties, choose between a number of scoring matrices, or supply their own scoring matrix for both pairwise alignments and multiple alignments. CLUSTAL W for UNIX and VMS systems is available at: <ftp://ebi.ac.uk>. Another program is MACAW (Schuler *et al.*, *Proteins Struct. Func. Genet.* 9:180-190 (1991), the entirety of which is herein incorporated by reference, for which both Macintosh and Microsoft Windows versions are available. MACAW uses a graphical interface, provides a choice of several
20 alignment algorithms, and is available by anonymous ftp at: [ncbi.nlm.nih.gov](ftp://ncbi.nlm.nih.gov) (directory/pub/macaw).

Sequence motifs are derived from multiple alignments and can be used to examine individual sequences or an entire database for subtle patterns. With motifs, it is sometimes

possible to detect distant relationships that may not be demonstrable based on comparisons of primary sequences alone. Currently, the largest collection of sequence motifs in the world is PROSITE (Bairoch and Bucher, *Nucleic Acid Research* 22: 3583-3589 (1994), the entirety of which is herein incorporated by reference). PROSITE may be accessed via either the ExPASy
5 server on the World Wide Web or anonymous ftp site. Many commercial sequence analysis packages also provide search programs that use PROSITE data.

A resource for searching protein motifs is the BLOCKS E-mail server developed by S. Henikoff, *Trends Biochem Sci.* 18:267-268 (1993), the entirety of which is herein incorporated by reference; Henikoff and Henikoff, *Nucleic Acid Research* 19:6565-6572 (1991), the entirety of which is herein incorporated by reference; Henikoff and Henikoff, *Proteins*, 17: 49-61 (1993). BLOCKS searches a protein or nucleotide sequence against a database of protein motifs or "blocks." Blocks are defined as short, ungapped multiple alignments that represent highly conserved protein patterns. The blocks themselves are derived from entries in PROSITE as well as other sources. Either a protein query or a nucleotide query can be submitted to the BLOCKS
10 server; if a nucleotide sequence is submitted, the sequence is translated in all six reading frames and motifs are sought for these conceptual translations. Once the search is completed, the server will return a ranked list of significant matches, along with an alignment of the query sequence to the matched BLOCKS entries.

Conserved protein domains can be represented by two-dimensional matrices, which
20 measure either the frequency or probability of the occurrences of each amino acid residue and deletions or insertions in each position of the domain. This type of model, when used to search against protein databases, is sensitive and usually yields more accurate results than simple motif searches. Two popular implementations of this approach are profile searches (such as GCG

program ProfileSearch) and Hidden Markov Models (HMMs)(Krough. *et al.*, *J. Mol. Biol.* 235:1501-1531, (1994); Eddy, *Current Opinion in Structural Biology*, 6:361-365, (1996), both of which are herein incorporated by reference in their entirety). In both cases, a large number of common protein domains have been converted into profiles, as present in the PROSITE library, or HHM models, as in the Pfam protein domain library (Sonnhammer *et al.*, *Proteins* 28:405-420 (1997), the entirety of which is herein incorporated by reference). Pfam contains more than 500 HMM models for enzymes, transcription factors, signal transduction molecules, and structural proteins. Protein databases can be queried with these profiles or HMM models, which will identify proteins containing the domain of interest. For example, HMMSW or HMMFS, two programs in a public domain package called HMMER (Sonnhammer *et al.*, *Proteins* 28:405-420, (1997)) can be used.

PROSITE and BLOCKS represent collected families of protein motifs. Thus, searching these databases entails submitting a single sequence to determine whether or not that sequence is similar to the members of an established family. Programs working in the opposite direction compare a collection of sequences with individual entries in the protein databases. An example of such a program is the Motif Search Tool, or MoST (Tatusov *et al. Proc. Natl. Acad. Sci.* 91: 12091-12095 (1994), the entirety of which is herein incorporated by reference). On the basis of an aligned set of input sequences, a weight matrix is calculated by using one of four methods (selected by the user). A weight matrix is simply a representation, position by position of how likely a particular amino acid will appear. The calculated weight matrix is then used to search the databases. To increase sensitivity, newly found sequences are added to the original data set, the weight matrix is recalculated, and the search is performed again. This procedure continues until no new sequences are found.

SUMMARY OF THE INVENTION

The present invention provides a substantially purified nucleic acid molecule that encodes a maize or soybean tocopherol synthesis pathway enzyme or fragment thereof, wherein the maize
5 or soybean tocopherol synthesis pathway enzyme is selected from the group consisting of: (a) deoxyarabiono-heptulosonate-P-synthase; (b) putative deoxyarabiono-heptulosonate-P-synthase; (c) dehydroquinase synthase; (d) dehydroquinase dehydratase; (e) putative dehydroquinase dehydratase; (f) shikimate dehydrogenase; (g) shikimate kinase; (h) enolpyruvylshikimate-P-synthase; (i) chorismate synthase; (j) chorismate mutase; (k) tyrosine transaminase; (l) putative
10 tyrosine transaminase; (m) transaminase A; (n) putative Transaminase A; (o) 4-hydroxyphenylpyruvate dioxygenase; (p) homogentisic acid dioxygenase; and (q) geranylgeranylpyrophosphate synthase.

The present invention also provides a substantially purified nucleic acid molecule that encodes a plant tocopherol synthesis pathway enzyme or fragment thereof, wherein the nucleic
15 acid molecule is selected from the group consisting of a nucleic acid molecule that encodes a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean dehydroquinase synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a
20 maize or soybean dehydroquinase dehydratase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean putative dehydroquinase dehydratase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean shikimate dehydrogenase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or

soybean shikimate kinase enzyme or fragment thereof; a nucleic acid molecule that encodes a
maize or soybean enolpyruvylshikimate-P-synthase enzyme or fragment thereof; a nucleic acid
molecule that encodes a maize or soybean chorismate synthase enzyme or fragment thereof; a
nucleic acid molecule that encodes a maize or soybean chorismate mutase enzyme or fragment
5 thereof; a nucleic acid molecule that encodes a maize or soybean tyrosine transaminase enzyme
or fragment thereof; a nucleic acid molecule that encodes a maize or soybean putative Tyrosine
transaminase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or
soybean transaminase A enzyme or fragment thereof; a nucleic acid molecule that encodes a
maize or soybean putative transaminase A enzyme or fragment thereof; a nucleic acid molecule
that encodes a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment
thereof; a nucleic acid molecule that encodes a maize or soybean homogentisic acid dioxygenase
enzyme or fragment thereof; and a nucleic acid molecule that encodes a maize or soybean
geranylgeranylpyrophosphate synthase enzyme or fragment thereof.

The present invention also provides a substantially purified maize or soybean tocopherol
15 synthesis pathway enzyme or fragment thereof, wherein the maize or soybean tocopherol
synthesis pathway enzyme is selected from the group consisting of (a) deoxyarabiono-
heptulosonate-P-synthase or fragment thereof; (b) putative deoxyarabiono-heptulosonate-P-
synthase or fragment thereof; (c) dehydroquinase synthase or fragment thereof; (d)
dehydroquinase dehydratase or fragment thereof; (e) putative dehydroquinase dehydratase or
20 fragment thereof; (f) shikimate dehydrogenase or fragment thereof; (g) shikimate kinase or
fragment thereof; (h) enolpyruvylshikimate-P-synthase or fragment thereof; (i) chorismate
synthase or fragment thereof; (j) chorismate mutase or fragment thereof; (k) tyrosine
transaminase or fragment thereof; (l) putative tyrosine transaminase or fragment thereof; (m)

transaminase A or fragment thereof; (n) putative Transaminase A or fragment thereof; (o) 4-hydroxyphenylpyruvate dioxygenase or fragment thereof; (p) homogentisic acid dioxygenase or fragment thereof; and (q) geranylgeranylpyrophosphate synthase or fragment thereof.

The present invention also provides a substantially purified maize or soybean tocopherol
5 synthesis pathway enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 1 through SEQ ID NO: 627.

The present invention also provides a substantially purified maize or soybean
10 deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 1 through SEQ ID NO: 97 and SEQ ID NO: 100 through SEQ ID NO: 146.

The present invention also provides a substantially purified maize or soybean
15 deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 97 and SEQ ID NO: 100 through SEQ ID NO: 146.

The present invention also provides a substantially purified maize or soybean putative
20 deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a

complement of SEQ ID NO: 98 through SEQ ID NO: 99 and SEQ ID NO: 147 through SEQ ID NO: 152.

The present invention also provides a substantially purified maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 98 through SEQ ID NO: 99 and
5 SEQ ID NO: 147 through SEQ ID NO: 152.

The present invention also provides a substantially purified maize dehydroquate synthase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 153 through
10 SEQ ID NO: 157.

The present invention also provides a substantially purified maize dehydroquate synthase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 153 through SEQ ID NO: 157.

The present invention also provides a substantially purified soybean dehydroquate dehydratase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence of a complement of SEQ ID NO: 160.
15

The present invention also provides a substantially purified soybean dehydroquate dehydratase enzyme or fragment thereof encoded by a nucleic acid sequence of SEQ ID NO: 160.
20

The present invention also provides a substantially purified maize putative dehydroquate dehydratase enzyme or fragment thereof encoded by a first nucleic acid molecule

which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 158 through SEQ ID NO: 159.

The present invention also provides a substantially purified maize putative
5 dehydroquinate dehydratase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 158 through SEQ ID NO: 159.

The present invention also provides a substantially purified maize or soybean shikimate dehydrogenase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule
10 having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 158 through SEQ ID NO: 159 and SEQ ID NO: 160.

The present invention also provides a substantially purified maize or soybean shikimate dehydrogenase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 158 through SEQ ID NO: 159 and SEQ ID NO: 160.

The present invention also provides a substantially purified maize or soybean shikimate
15 kinase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 161 through SEQ ID NO: 179 and SEQ ID NO: 180 through SEQ ID NO: 183.

20 The present invention also provides a substantially purified maize or soybean shikimate kinase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 161 through SEQ ID NO: 179 and SEQ ID NO: 180 through SEQ ID NO: 183.

The present invention also provides a substantially purified maize enolpyruvylshikimate-P-synthase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 184 through SEQ ID NO: 198.

The present invention also provides a substantially purified maize enolpyruvylshikimate-P-synthase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 184 through SEQ ID NO: 198.

The present invention also provides a substantially purified maize or soybean chorismate synthase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 199 through SEQ ID NO: 231 and SEQ ID NO: 232 through SEQ ID NO: 255.

The present invention also provides a substantially purified maize or soybean chorismate synthase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 199 through SEQ ID NO: 231 and SEQ ID NO: 232 through SEQ ID NO: 255.

The present invention also provides a substantially purified maize or soybean chorismate mutase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 256 through SEQ ID NO: 277 and SEQ ID NO: 278 through SEQ ID NO: 284.

The present invention also provides a substantially purified maize or soybean chorismate mutase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 256 through SEQ ID NO: 277 and SEQ ID NO: 278 through SEQ ID NO: 284.

5 The present invention also provides a substantially purified maize tyrosine transaminase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 285 through SEQ ID NO: 286.

10 The present invention also provides a substantially purified maize tyrosine transaminase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 285 through SEQ ID NO: 286.

15 The present invention also provides a substantially purified maize or soybean putative tyrosine transaminase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 287 through SEQ ID NO: 292 and SEQ ID NO: 293 through SEQ ID NO: 300.

20 The present invention also provides a substantially purified maize or soybean putative tyrosine transaminase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 287 through SEQ ID NO: 292 and SEQ ID NO: 293 through SEQ ID NO: 300.

 The present invention also provides a substantially purified maize or soybean transaminase A enzyme or fragment thereof encoded by a first nucleic acid molecule which

specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 301 through SEQ ID NO: 474 and SEQ ID NO: 475 through SEQ ID NO: 581.

5 The present invention also provides a substantially purified maize or soybean transaminase A enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 301 through SEQ ID NO: 474 and SEQ ID NO: 475 through SEQ ID NO: 581.

The present invention also provides a substantially purified soybean putative transaminase A enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 582 through SEQ ID NO: 597.

15 The present invention also provides a substantially purified soybean putative transaminase A enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 582 through SEQ ID NO: 597.

The present invention also provides a substantially purified maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 598 through SEQ ID NO: 600 and SEQ ID NO: 601 through SEQ ID NO: 607.

20

The present invention also provides a substantially purified maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment thereof encoded by a nucleic acid

sequence selected from the group consisting of SEQ ID NO: 598 through SEQ ID NO: 600 and SEQ ID NO: 601 through SEQ ID NO: 607.

5 The present invention also provides a substantially purified maize or soybean homogentisic acid dioxygenase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 608 through SEQ ID NO: 615 and SEQ ID NO: 616 through SEQ ID NO: 621.

6 The present invention also provides a substantially purified maize or soybean homogentisic acid dioxygenase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 608 through SEQ ID NO: 615 and SEQ ID NO: 616 through SEQ ID NO: 621.

7 The present invention also provides a substantially purified maize or soybean geranylgeranylpyrophosphate synthase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 622 through SEQ ID NO: 624 and SEQ ID NO: 625 through SEQ ID NO: 627.

8 The present invention also provides a substantially purified maize or soybean
20 geranylgeranylpyrophosphate synthase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 622 through SEQ ID NO: 624 and SEQ ID NO: 625 through SEQ ID NO: 627.

The present invention also provides a purified antibody or fragment thereof which is capable of specifically binding to a maize or soybean tocopherol synthesis pathway enzyme or fragment thereof, wherein the maize or soybean tocopherol synthesis pathway enzyme or fragment thereof is encoded by a nucleic acid molecule comprising a nucleic acid sequence
5 selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a substantially purified maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the
10 second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 1 through SEQ ID NO: 97 and SEQ ID NO: 100 through SEQ ID NO: 146.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize or soybean putative
15 deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 98 through SEQ ID NO: 99 and SEQ ID NO: 147 through SEQ ID NO: 152.

20 The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize dehydroquate synthase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically

hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence consisting of a complement of SEQ ID NO: 153 through SEQ ID NO: 157.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a soybean dehydroquinase
5 dehydratase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 160.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize putative
10 dehydroquinase dehydratase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 158 through SEQ ID NO: 159.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize or soybean shikimate
15 dehydrogenase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID
20 NO: 158 through SEQ ID NO: 159 and SEQ ID NO: 160.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize or soybean shikimate kinase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically

hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 161 through SEQ ID NO: 179 and SEQ ID NO: 180 through SEQ ID NO: 183.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize enolpyruvylshikimate-P-synthase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 184 through SEQ ID NO: 198.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize or soybean chorismate synthase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 199 through SEQ ID NO: 231 and SEQ ID NO: 232 through SEQ ID NO: 255.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize or soybean chorismate mutase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 256 through SEQ ID NO: 277 and SEQ ID NO: 278 through SEQ ID NO: 284.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize tyrosine transaminase

enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 285 through SEQ ID NO: 286.

5 The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize or soybean putative tyrosine transaminase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 287 through SEQ ID NO: 292 and SEQ ID NO: 293 through SEQ ID NO: 300.

10 The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize or soybean transaminase A enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 301 through SEQ ID NO: 474 and SEQ ID NO: 475 through SEQ ID NO: 581.

15 The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a soybean putative transaminase A enzyme or fragment thereof encoded by a first nucleic acid molecule which
20 specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 582 through SEQ ID NO: 597.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a
5 complement of SEQ ID NO: 598 through SEQ ID NO: 600 and SEQ ID NO: 601 through SEQ ID NO: 607.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize or soybean
10 homogentisic acid dioxygenase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 608 through SEQ ID NO: 615 and SEQ ID NO: 616 through SEQ ID NO: 621.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize or soybean
15 geranylgeranylpyrophosphate synthase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a
20 complement of SEQ ID NO: 622 through SEQ ID NO: 624 and SEQ ID NO: 625 through SEQ ID NO: 627.

The present invention also provides a transformed plant having a nucleic acid molecule which comprises: (A) an exogenous promoter region which functions in a plant cell to cause the

production of a mRNA molecule; (B) a structural nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of (a) a nucleic acid sequence which encodes for a deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; (b) a nucleic acid sequence which encodes for a putative deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; (c) a nucleic acid sequence which encodes for a dehydroquinase synthase enzyme or fragment thereof; (d) a nucleic acid sequence which encodes for a dehydroquinase dehydratase enzyme or fragment thereof; (e) a nucleic acid sequence which encodes for a putative dehydroquinase dehydratase enzyme or fragment thereof; (f) a nucleic acid sequence which encodes for a shikimate dehydrogenase enzyme or fragment thereof; (g) a nucleic acid sequence which encodes for a shikimate kinase enzyme or fragment thereof; (h) a nucleic acid sequence which encodes for an enolpyruvylshikimate-P-synthase enzyme or fragment thereof; (i) a nucleic acid sequence which encodes for a chorismate synthase enzyme or fragment thereof; (j) a nucleic acid sequence which encodes for a chorismate mutase enzyme or fragment thereof; (k) a nucleic acid sequence which encodes for a tyrosine transaminase enzyme or fragment thereof; (l) a nucleic acid sequence which encodes for a putative Tyrosine transaminase enzyme or fragment thereof; (m) a nucleic acid sequence which encodes for a transaminase A enzyme or fragment thereof; (n) a nucleic acid sequence which encodes for a putative transaminase A enzyme or fragment thereof; (o) a nucleic acid sequence which encodes for a 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment thereof; (p) a nucleic acid sequence which encodes for a homogentisic acid dioxygenase enzyme or fragment thereof; (q) a nucleic acid sequence which encodes for a geranylgeranylpyrophosphate synthase enzyme or fragment thereof; and (r) a nucleic acid sequence which is complementary to any of the nucleic acid sequences of (a) through (q); and (C) a 3' non-translated sequence that functions in the plant cell

to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of the mRNA molecule.

The present invention also provides a transformed plant having a nucleic acid molecule which comprises: (A) an exogenous promoter region which functions in a plant cell to cause the
5 production of a mRNA molecule; which is linked to (B) a structural nucleic acid molecule, wherein the structural nucleic acid molecule encodes a plant tocopherol synthesis pathway enzyme or fragment thereof, the structural nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627 or
10 fragment thereof; which is linked to (C) a 3' non-translated sequence that functions in the plant cell to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of the mRNA molecule.

The present invention also provides a transformed plant having a nucleic acid molecule which comprises: (A) an exogenous promoter region which functions in a plant cell to cause the
15 production of a mRNA molecule; which is linked to (B) a structural nucleic acid molecule, wherein the structural nucleic acid molecule is selected from the group consisting of a nucleic acid molecule that encodes for a deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes for a putative deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes for a dehydroquininate
20 dehydratase enzyme or fragment thereof; a nucleic acid molecule that encodes for a putative dehydroquininate dehydratase enzyme or fragment thereof; a nucleic acid molecule that encodes for a shikimate dehydrogenase enzyme or fragment thereof; a nucleic acid molecule that encodes for a shikimate kinase enzyme or fragment thereof; a nucleic acid molecule that encodes for an

enolpyruvylshikimate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes for a chorismate synthase enzyme or fragment thereof; a nucleic acid molecule that encodes for a chorismate mutase enzyme or fragment thereof; a nucleic acid molecule that encodes for a tyrosine transaminase enzyme or fragment thereof; a nucleic acid molecule that encodes for a putative Tyrosine transaminase enzyme or fragment thereof; a nucleic acid molecule that encodes for a transaminase A enzyme or fragment thereof; a nucleic acid molecule that encodes for a putative transaminase A enzyme or fragment thereof; a nucleic acid molecule that encodes for a 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment thereof; a nucleic acid molecule that encodes for a homogentisic acid dioxygenase enzyme or fragment thereof; a nucleic acid molecule that encodes for a geranylgeranylpyrophosphate synthase enzyme or fragment thereof; which is linked to (C) a 3' non-translated sequence that functions in the plant cell to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of the mRNA molecule.

The present invention also provides a transformed plant having a nucleic acid molecule which comprises: (A) an exogenous promoter region which functions in a plant cell to cause the production of a mRNA molecule; which is linked to (B) a transcribed nucleic acid molecule with a transcribed strand and a non-transcribed strand, wherein the transcribed strand is complementary to a nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627 or fragment thereof; which is linked to (C) a 3' non-translated sequence that functions in plant cells to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of the mRNA molecule.

The present invention also provides a transformed plant having a nucleic acid molecule which comprises: (A) an exogenous promoter region which functions in a plant cell to cause the

production of a mRNA molecule; which is linked to: (B) a transcribed nucleic acid molecule with a transcribed strand and a non-transcribed strand, wherein a transcribed mRNA of the transcribed strand is complementary to an endogenous mRNA molecule having a nucleic acid sequence selected from the group consisting of an endogenous mRNA molecule that encodes a

5 maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize dehydroquate synthase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a soybean dehydroquate dehydratase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize putative dehydroquate dehydratase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean shikimate dehydrogenase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean shikimate kinase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize enolpyruvylshikimate-P-synthase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean chorismate synthase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean chorismate mutase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize tyrosine transaminase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean putative tyrosine transaminase enzyme or fragment thereof; an endogenous

20 mRNA molecule that encodes a maize or soybean transaminase A enzyme or fragment thereof; an endogenous mRNA molecule that encodes a soybean putative transaminase A enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment thereof; an endogenous mRNA

molecule that encodes a maize or soybean homogentisic acid dioxygenase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean geranylgeranylpyrophosphate synthase enzyme or fragment thereof; which is linked to (C) a 3' non-translated sequence that functions in the plant cell to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of the mRNA molecule.

The present invention also provides a method for determining a level or pattern of a plant tocopherol synthesis pathway enzyme in a plant cell or plant tissue comprising: (A) incubating, under conditions permitting nucleic acid hybridization, a marker nucleic acid molecule, the marker nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof or fragment of either, with a complementary nucleic acid molecule obtained from the plant cell or plant tissue, wherein nucleic acid hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant cell or plant tissue permits the detection of the plant tocopherol synthesis pathway enzyme; (B) permitting hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant cell or plant tissue; and (C) detecting the level or pattern of the complementary nucleic acid, wherein the detection of the complementary nucleic acid is predictive of the level or pattern of the plant tocopherol synthesis pathway enzyme.

The present invention also provides a method for determining a level or pattern of a plant tocopherol synthesis pathway enzyme in a plant cell or plant tissue comprising: (A) incubating, under conditions permitting nucleic acid hybridization, a marker nucleic acid molecule, the marker nucleic acid molecule comprising a nucleic acid molecule that encodes a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or complement thereof or fragment of

either; a nucleic acid molecule that encodes a maize or soybean putative deoxyarabiono-
heptulosonate-P-synthase enzyme or complement thereof or fragment of either; a nucleic acid
molecule that encodes a maize dehydroquinase synthase enzyme or complement thereof or
fragment of either; a nucleic acid molecule that encodes a soybean dehydroquinase dehydratase
5 enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a
maize putative dehydroquinase dehydratase enzyme or complement thereof or fragment of either;
a nucleic acid molecule that encodes a maize or soybean shikimate dehydrogenase enzyme or
complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or
soybean shikimate kinase enzyme or complement thereof or fragment of either; a nucleic acid
10 molecule that encodes a maize enolpyruvylshikimate-P-synthase enzyme or complement thereof
or fragment of either; a nucleic acid molecule that encodes a maize or soybean chorismate
synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that
encodes a maize or soybean chorismate mutase enzyme or complement thereof or fragment of
either; a nucleic acid molecule that encodes a maize tyrosine transaminase enzyme or
15 complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or
soybean putative tyrosine transaminase enzyme or complement thereof or fragment of either; a
nucleic acid molecule that encodes a maize or soybean transaminase A enzyme or complement
thereof or fragment of either; a nucleic acid molecule that encodes a soybean putative
transaminase A enzyme or complement thereof or fragment of either; a nucleic acid molecule
20 that encodes a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or complement
thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean
homogentisic acid dioxygenase enzyme or complement thereof or fragment of either; a nucleic
acid molecule that encodes a maize or soybean geranylgeranylpyrophosphate synthase enzyme or

complement thereof or fragment of either, with a complementary nucleic acid molecule obtained from the plant cell or plant tissue, wherein nucleic acid hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant cell or plant tissue permits the detection of the plant tocopherol synthesis pathway enzyme; (B) permitting
5 hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant cell or plant tissue; and (C) detecting the level or pattern of the complementary nucleic acid, wherein the detection of the complementary nucleic acid is predictive of the level or pattern of the plant tocopherol synthesis pathway enzyme.

10 The present invention also provides a method for determining a level or pattern of a plant tocopherol synthesis pathway enzyme in a plant cell or plant tissue under evaluation which comprises assaying the concentration of a molecule, whose concentration is dependent upon the expression of a gene, the gene specifically hybridizes to a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof, in comparison to the concentration of that molecule present in a reference
15 plant cell or a reference plant tissue with a known level or pattern of the plant tocopherol synthesis pathway enzyme, wherein the assayed concentration of the molecule is compared to the assayed concentration of the molecule in the reference plant cell or reference plant tissue with the known level or pattern of the plant tocopherol synthesis pathway enzyme.

20 The present invention also provides a method for determining a level or pattern of a plant tocopherol synthesis pathway enzyme in a plant cell or plant tissue under evaluation which comprises assaying the concentration of a molecule, whose concentration is dependent upon the expression of a gene, the gene specifically hybridizes to a nucleic acid molecule selected from the group consisting of a nucleic acid molecule that encodes a maize or soybean deoxyarabiono-

6
10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100

heptulosonate-P-synthase enzyme; a nucleic acid molecule that encodes a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or complement thereof; a nucleic acid molecule that encodes a maize dehydroquinase synthase enzyme or complement thereof; a nucleic acid molecule that encodes a soybean dehydroquinase dehydratase enzyme or complement thereof; a nucleic acid molecule that encodes a maize putative dehydroquinase dehydratase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean shikimate dehydrogenase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean shikimate kinase enzyme or complement thereof; a nucleic acid molecule that encodes a maize enolpyruvylshikimate-P-synthase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean chorismate synthase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean chorismate mutase enzyme or complement thereof; a nucleic acid molecule that encodes a maize tyrosine transaminase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean putative tyrosine transaminase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean transaminase A enzyme or complement thereof; a nucleic acid molecule that encodes a soybean putative transaminase A enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean homogentisic acid dioxygenase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean geranylgeranylpyrophosphate synthase enzyme or complement thereof, in comparison to the concentration of that molecule present in a reference plant cell or a reference plant tissue with a known level or pattern of the plant tocopherol synthesis pathway enzyme, wherein the assayed concentration of the molecule is compared to the assayed

concentration of the molecule in the reference plant cell or the reference plant tissue with the known level or pattern of the plant tocopherol synthesis pathway enzyme.

The present invention provides a method of determining a mutation in a plant whose presence is predictive of a mutation affecting a level or pattern of a protein comprising the steps:

5 (A) incubating, under conditions permitting nucleic acid hybridization, a marker nucleic acid, the marker nucleic acid selected from the group of marker nucleic acid molecules which specifically hybridize to a nucleic acid molecule having a nucleic acid sequence selected from the group of SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof or fragment of either and a complementary nucleic acid molecule obtained from the plant, wherein nucleic acid

10 hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant permits the detection of a polymorphism whose presence is predictive of a mutation affecting the level or pattern of the protein in the plant; (B) permitting hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant; and (C) detecting the presence of the polymorphism, wherein the detection of the polymorphism is predictive of the mutation.

The present invention also provides a method for determining a mutation in a plant whose presence is predictive of a mutation affecting the level or pattern of a plant tocopherol synthesis pathway enzyme comprising the steps: (A) incubating, under conditions permitting nucleic acid hybridization, a marker nucleic acid molecule, the marker nucleic acid molecule comprising a

20 nucleic acid molecule that is linked to a gene, the gene specifically hybridizes to a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof and a complementary nucleic acid molecule obtained from the plant, wherein nucleic acid hybridization between the marker nucleic acid

molecule and the complementary nucleic acid molecule obtained from the plant permits the detection of a polymorphism whose presence is predictive of a mutation affecting the level or pattern of the plant tocopherol synthesis pathway enzyme in the plant; (B) permitting hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant; and (C) detecting the presence of the polymorphism, wherein the detection of the polymorphism is predictive of the mutation.

The present invention also provides a method for determining a mutation in a plant whose presence is predictive of a mutation affecting the level or pattern of a plant tocopherol synthesis pathway enzyme comprising the steps: (A) incubating, under conditions permitting nucleic acid hybridization, a marker nucleic acid molecule, the marker nucleic acid molecule comprising a nucleic acid molecule that is linked to a gene, the gene specifically hybridizes to a nucleic acid molecule selected from the group consisting of a nucleic acid molecule that encodes a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or complement thereof; a nucleic acid molecule that encodes a maize dehydroquinase synthase enzyme or complement thereof; a nucleic acid molecule that encodes a soybean dehydroquinase dehydratase enzyme or complement thereof; a nucleic acid molecule that encodes a maize putative dehydroquinase dehydratase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean shikimate dehydrogenase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean shikimate kinase enzyme or complement thereof; a nucleic acid molecule that encodes a maize enolpyruvylshikimate-P-synthase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean chorismate synthase enzyme or complement thereof; a nucleic acid molecule that

encodes a maize or soybean chorismate mutase enzyme or complement thereof; a nucleic acid molecule that encodes a maize tyrosine transaminase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean putative tyrosine transaminase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean transaminase A enzyme or complement thereof; a nucleic acid molecule that encodes a soybean putative transaminase A enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean homogentisic acid dioxygenase enzyme or complement thereof; a nucleic acid molecule that encodes a maize or soybean geranylgeranylpyrophosphate synthase enzyme or complement thereof, and a complementary nucleic acid molecule obtained from the plant, wherein nucleic acid hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant permits the detection of a polymorphism whose presence is predictive of a mutation affecting the level or pattern of the plant tocopherol synthesis pathway enzyme in the plant; (B) permitting hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant; and (C) detecting the presence of the polymorphism, wherein the detection of the polymorphism is predictive of the mutation.

The present invention also provides a method of producing a plant containing an overexpressed protein comprising: (A) transforming the plant with a functional nucleic acid molecule, wherein the functional nucleic acid molecule comprises a promoter region, wherein the promoter region is linked to a structural region, wherein the structural region has a nucleic acid sequence selected from group consisting of SEQ ID NO: 1 through SEQ ID NO: 627; wherein the structural region is linked to a 3' non-translated sequence that functions in the plant to cause

termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and wherein the functional nucleic acid molecule results in overexpression of the protein; and (B) growing the transformed plant.

5 The present invention also provides a method of producing a plant containing an overexpressed plant tocopherol synthesis pathway enzyme comprising: (A) transforming the plant with a functional nucleic acid molecule, wherein the functional nucleic acid molecule comprises a promoter region, wherein the promoter region is linked to a structural region, wherein the structural region comprises a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627 or fragment thereof; wherein the structural region is linked to a 3' non-translated sequence that functions in the plant to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and wherein the functional nucleic acid molecule results in overexpression of the plant tocopherol synthesis pathway enzyme; and (B) growing the transformed plant.

10 The present invention also provides a method of producing a plant containing an overexpressed plant tocopherol synthesis pathway enzyme comprising: (A) transforming the plant with a functional nucleic acid molecule, wherein the functional nucleic acid molecule comprises a promoter region, wherein the promoter region is linked to a structural region, wherein the structural region comprises a nucleic acid molecule selected from the group
15 consisting of a nucleic acid molecule that encodes a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize dehydroquinate synthase enzyme or fragment
20

thereof; a nucleic acid molecule that encodes a soybean dehydroquinase dehydratase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize putative dehydroquinase dehydratase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean shikimate dehydrogenase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean shikimate kinase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize enolpyruvylshikimate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean chorismate synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean chorismate mutase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize tyrosine transaminase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean putative tyrosine transaminase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean transaminase A enzyme or fragment thereof; a nucleic acid molecule that encodes a soybean putative transaminase A enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean homogentisic acid dioxygenase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean geranylgeranylpyrophosphate synthase enzyme or fragment thereof; wherein the structural region is linked to a 3' non-translated sequence that functions in the plant to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and wherein the functional nucleic acid molecule results in overexpression of the plant tocopherol synthesis pathway enzyme; and (B) growing the transformed plant.

The present invention also provides a method of producing a plant containing reduced levels of a plant tocopherol synthesis pathway enzyme comprising: (A) transforming the plant

with a functional nucleic acid molecule, wherein the functional nucleic acid molecule comprises a promoter region, wherein the promoter region is linked to a structural region, wherein the structural region comprises a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627; wherein the structural region is
5 linked to a 3' non-translated sequence that functions in the plant to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and wherein the functional nucleic acid molecule results in co-suppression of the plant tocopherol synthesis pathway enzyme; and (B) growing the transformed plant.

The present invention also provides a method of producing a plant containing reduced
10 levels of a plant tocopherol synthesis pathway enzyme comprising: (A) transforming the plant with a functional nucleic acid molecule, wherein the functional nucleic acid molecule comprises a promoter region, wherein the promoter region is linked to a structural region, wherein the structural region comprises a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a nucleic acid molecule that encodes a maize or soybean deoxyarabiono-
15 heptulosonate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize dehydroquinate synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a soybean dehydroquinate dehydratase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize putative dehydroquinate
20 dehydratase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean shikimate dehydrogenase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean shikimate kinase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize enolpyruvylshikimate-P-synthase enzyme or fragment thereof; a

nucleic acid molecule that encodes a maize or soybean chorismate synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean chorismate mutase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize tyrosine transaminase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean putative tyrosine
5 transaminase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean transaminase A enzyme or fragment thereof; a nucleic acid molecule that encodes a soybean putative transaminase A enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean homogentisic acid dioxygenase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean
10 geranylgeranylpyrophosphate synthase enzyme or fragment thereof, wherein the structural region is linked to a 3' non-translated sequence that functions in the plant to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and wherein the functional nucleic acid molecule results in co-suppression of the plant
15 tocopherol synthesis pathway enzyme; and (B) growing the transformed plant.

The present invention also provides a method for reducing expression of a plant tocopherol synthesis pathway enzyme in a plant comprising: (A) transforming the plant with a nucleic acid molecule, the nucleic acid molecule having an exogenous promoter region which functions in a plant cell to cause the production of a mRNA molecule, wherein the exogenous
20 promoter region is linked to a transcribed nucleic acid molecule having a transcribed strand and a non-transcribed strand, wherein the transcribed strand is complementary to a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof or fragments of either and the transcribed

strand is complementary to an endogenous mRNA molecule; and wherein the transcribed nucleic acid molecule is linked to a 3' non-translated sequence that functions in the plant cell to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and (B) growing the transformed plant.

5 The present invention also provides a method for reducing expression of a plant tocopherol synthesis pathway enzyme in a plant comprising: (A) transforming the plant with a nucleic acid molecule, the nucleic acid molecule having an exogenous promoter region which functions in a plant cell to cause the production of a mRNA molecule, wherein the exogenous promoter region is linked to a transcribed nucleic acid molecule having a transcribed strand and a non-transcribed strand, wherein a transcribed mRNA of the transcribed strand is complementary to a nucleic acid molecule selected from the group consisting of an endogenous mRNA molecule that encodes a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize dehydroquate synthase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a soybean dehydroquate dehydratase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize putative dehydroquate dehydratase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean shikimate dehydrogenase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean shikimate kinase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize enolpyruvylshikimate-P-synthase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean chorismate synthase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or

soybean chorismate mutase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize tyrosine transaminase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean putative tyrosine transaminase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean transaminase A enzyme or fragment thereof; an endogenous mRNA molecule that encodes a soybean putative transaminase A enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment thereof; an endogenous mRNA molecule that encodes a maize or soybean homogentisic acid dioxygenase enzyme or fragment thereof; and an endogenous mRNA molecule that encodes a maize or soybean geranylgeranylpyrophosphate synthase enzyme or fragment thereof; and wherein the transcribed nucleic acid molecule is linked to a 3' non-translated sequence that functions in the plant cell to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and (B) growing the transformed plant.

The present invention also provides a method of determining an association between a polymorphism and a plant trait comprising: (A) hybridizing a nucleic acid molecule specific for the polymorphism to genetic material of a plant, wherein the nucleic acid molecule has a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof or fragment of either; and (B) calculating the degree of association between the polymorphism and the plant trait.

The present invention also provides a method of determining an association between a polymorphism and a plant trait comprising: (A) hybridizing a nucleic acid molecule specific for the polymorphism to genetic material of a plant, wherein the nucleic acid molecule is selected from the group consisting of a nucleic acid molecule that encodes a maize or soybean

deoxyarabiono-heptulosonate-P-synthase enzyme or complement thereof or fragment of either; a
 nucleic acid molecule that encodes a maize or soybean putative deoxyarabiono-heptulosonate-P-
 synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that
 encodes a maize dehydroquinase synthase enzyme or complement thereof or fragment of either; a
 5 nucleic acid molecule that encodes a soybean dehydroquinase dehydratase enzyme or
 complement thereof or fragment of either; a nucleic acid molecule that encodes a maize putative
 dehydroquinase dehydratase enzyme or complement thereof or fragment of either; a nucleic acid
 molecule that encodes a maize or soybean shikimate dehydrogenase enzyme or complement
 thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean shikimate
 10 kinase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes
 a maize enolpyruvylshikimate-P-synthase enzyme or complement thereof or fragment of either; a
 nucleic acid molecule that encodes a maize or soybean chorismate synthase enzyme or
 complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or
 soybean chorismate mutase enzyme or complement thereof or fragment of either; a nucleic acid
 15 molecule that encodes a maize tyrosine transaminase enzyme or complement thereof or fragment
 of either; a nucleic acid molecule that encodes a maize or soybean putative tyrosine transaminase
 enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a
 maize or soybean transaminase A enzyme or complement thereof or fragment of either; a nucleic
 acid molecule that encodes a soybean putative transaminase A enzyme or complement thereof or
 20 fragment of either; a nucleic acid molecule that encodes a maize or soybean 4-
 hydroxyphenylpyruvate dioxygenase enzyme or complement thereof or fragment of either; a
 nucleic acid molecule that encodes a maize or soybean homogentisic acid dioxygenase enzyme
 or complement thereof or fragment of either; and a nucleic acid molecule that encodes a maize or

soybean geranylgeranylpyrophosphate synthase enzyme or complement thereof or fragment of either; and (B) calculating the degree of association between the polymorphism and the plant trait.

5 The present invention also provides a method of isolating a nucleic acid that encodes a plant tocopherol synthesis pathway enzyme or fragment thereof comprising: (A) incubating under conditions permitting nucleic acid hybridization, a first nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof or fragment of either with a complementary second nucleic acid molecule obtained from a plant cell or plant tissue; (B) permitting hybridization between the first nucleic acid molecule and the second nucleic acid molecule obtained from the plant cell or plant tissue; and (C) isolating the second nucleic acid molecule.

10 The present invention also provides a method of isolating a nucleic acid molecule that encodes a plant tocopherol synthesis pathway enzyme or fragment thereof comprising: (A) incubating under conditions permitting nucleic acid hybridization, a first nucleic acid molecule selected from the group consisting of a nucleic acid molecule that encodes a maize or soybean copalyl diphosphate synthase enzyme or complement thereof or fragment of either, a nucleic acid molecule that encodes a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize dehydroquinase synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a soybean dehydroquinase dehydratase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize putative dehydroquinase dehydratase enzyme or

15
20

complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean shikimate dehydrogenase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean shikimate kinase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize

5 enolpyruvylshikimate-P-synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean chorismate synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean chorismate mutase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize tyrosine transaminase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean putative tyrosine transaminase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean transaminase A enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a soybean putative transaminase A enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean 4-

15 hydroxyphenylpyruvate dioxygenase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean homogentisic acid dioxygenase enzyme or complement thereof or fragment of either; and a nucleic acid molecule that encodes a maize or soybean geranylgeranylpyrophosphate synthase enzyme or complement thereof or fragment of either, with a complementary second nucleic acid molecule obtained from a plant cell or plant
20 tissue; (B) permitting hybridization between the plant tocopherol synthesis pathway enzyme nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant cell or plant tissue; and (C) isolating the second nucleic acid molecule.

DETAILED DESCRIPTION OF THE INVENTION

Definitions and Agents of the Present Invention

Definitions:

As used herein, a tocopherol synthesis pathway enzyme is any enzyme that is associated
5 with the synthesis or degradation of tocopherol.

As used herein, a tocopherol synthesis enzyme is any enzyme that is associated with the
synthesis of tocopherol.

As used herein, a tocopherol degradation enzyme is any enzyme that is associated with
the degradation of tocopherol.

As used herein, deoxyarabinoheptulosonate phosphate synthase (DAHP synthase) is any
10 enzyme that catalyzes the formation of deoxyarabinoheptulosonate phosphate from erythrose
phosphate.

As used herein, dehydroquinate synthase is any enzyme that catalyzes the formation of
dehydroquinate from erythrose phosphate via an NAD-dependent reaction.

As used herein, dehydroquinate dehydratase is any enzyme that catalyzes the
15 stereospecific syn-dehydration of dehydroquinate to dehydroshikimate.

As used herein, shikimate dehydrogenase is any enzyme that catalyzes the NADPH-
dependent conversion of dehydroshikimate to shikimate.

As used herein, shikimate kinase is any enzyme that catalyzes the phosphorylation of
20 shikimate to shikimate-3-phosphate.

As used herein, enolpyruvylshikimatephosphate synthase (EPSPS) is any enzyme that
catalyzes the reversible transfer of the carboxyvinyl moiety of phosphoenolpyruvate to
shikimatephosphate, yielding enolpyruvylshikimate phosphate.

As used herein, chorismate synthase is any enzyme that catalyzes the conversion of enolpyruvylshikimate phosphate to chorismic acid with the introduction of a double bond of the aromatic ring in a trans-1,4-elimination of inorganic phosphorous.

As used herein, chorismate mutase is any enzyme that catalyzes the reaction that converts
5 chorismic acid to prephenic acid.

As used herein, prephenate dehydrogenase is any enzyme that catalyzes the formation of 4-hydroxyphenylpyruvate from prephenate via an NAD-dependent or an NADP-dependent reaction.

As used herein, tyrosine transaminase is any enzyme that catalyzes the pyridoxal-phosphate dependent conversion of L-tyrosine to 4-hydroxyphenylpyruvate.

As used herein, L-amino-acid oxidase is any enzyme that catalyzes the reaction to convert tyrosine to 4-hydroxyphenylpyruvate.

As used herein, aromatic amino acid amino transferase is any enzyme that catalyzes the reaction that converts 4-hydroxyphenylpyruvate to tyrosine.

As used herein, an aspartic acid amino transferase or transaminase A is any enzyme that catalyzes the formation of phenylalanine and tyrosine using phenylpyruvate or p-hydroxy phenylpyruvate, respectively.

As used herein, hydroxyphenylpyruvate dioxygenase is any enzyme that catalyzes the formation of homogentisate from hydroxyphenylpyruvate.

As used herein, geranylgeranyl-pyrophosphate synthase is any enzyme that catalyzes the
20 formation of geranylgeranylpyrophosphate by prenyltransferring isoprene moiety from isopentenylpyrophosphate to farnesylpyrophosphate.

As used herein, geranylgeranylpyrophosphate (GGPP) hydrogenase is any enzyme that catalyzes the reaction to convert geranylgeranylpyrophosphate to phytylpyrophosphate via an NADPH-dependent reaction.

As used herein, homogentisate:phytyl transferase is any enzyme that catalyzes the
5 reaction to convert homogentisic acid to 2-methyl-6-phytylbenzoquinol.

As used herein, tocopherol cyclase is any enzyme that catalyzes the cyclization of 2,3-dimethyl-6-phytylbenzoquinol to form γ -tocopherol.

As used herein, tocopherol methyltransferase is any enzyme that catalyzes the reaction that forms α -tocopherol from other tocopherols via an S-adenosyl methionine (SAM)-dependent
10 reaction.

As used herein, homogentisic acid dioxygenase is any enzyme that catalyzes the reaction to convert homogentisic acid to fumarylacetoacetate.

Agents

(a) Nucleic Acid Molecules

Agents of the present invention include plant nucleic acid molecules and more preferably include maize and soybean nucleic acid molecules and more preferably include nucleic acid molecules of the maize genotypes B73 (Illinois Foundation Seeds, Champaign, Illinois U.S.A.), B73 x Mo17 (Illinois Foundation Seeds, Champaign, Illinois U.S.A.), DK604 (Dekalb Genetics, Dekalb, Illinois U.S.A.), H99 (Illinois Foundation Seeds, Champaign, Illinois U.S.A.), RX601
15 (Asgrow Seed Company, Des Moines, Iowa), Mo17 (Illinois Foundation Seeds, Champaign, Illinois U.S.A.), and soybean types Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa), C1944 (United States Department of Agriculture (USDA) Soybean Germplasm Collection,
20

Urbana, Illinois U.S.A.), Cristalina (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.), FT108 (Monsoy, Brazil), Hartwig (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.), BW211S Null (Tohoku University, Morioka, Japan), PI507354 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.), Asgrow A4922 (Asgrow Seed Company, Des Moines, Iowa U.S.A.), PI227687 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.), PI229358 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) and Asgrow A3237 (Asgrow Seed Company, Des Moines, Iowa U.S.A.).

A subset of the nucleic acid molecules of the present invention includes nucleic acid molecules that are marker molecules. Another subset of the nucleic acid molecules of the present invention include nucleic acid molecules that encode a protein or fragment thereof. Another subset of the nucleic acid molecules of the present invention are EST molecules.

Fragment nucleic acid molecules may encode significant portion(s) of, or indeed most of, these nucleic acid molecules. Alternatively, the fragments may comprise smaller oligonucleotides (having from about 15 to about 250 nucleotide residues and more preferably, about 15 to about 30 nucleotide residues).

As used herein, an agent, be it a naturally occurring molecule or otherwise may be "substantially purified," if desired, such that one or more molecules that is or may be present in a naturally occurring preparation containing that molecule will have been removed or will be present at a lower concentration than that at which it would normally be found.

The agents of the present invention will preferably be "biologically active" with respect to either a structural attribute, such as the capacity of a nucleic acid to hybridize to another nucleic acid molecule, or the ability of a protein to be bound by an antibody (or to compete with

another molecule for such binding). Alternatively, such an attribute may be catalytic and thus involve the capacity of the agent to mediate a chemical reaction or response.

The agents of the present invention may also be recombinant. As used herein, the term recombinant means any agent (e.g. DNA, peptide etc.), that is, or results, however indirect, from
5 human manipulation of a nucleic acid molecule.

It is understood that the agents of the present invention may be labeled with reagents that facilitate detection of the agent (e.g. fluorescent labels, Prober *et al.*, *Science* 238:336-340 (1987); Albarella *et al.*, EP 144914; chemical labels, Sheldon *et al.*, U.S. Patent No. 4,582,789; Albarella *et al.*, U.S. Patent No. 4,563,417; modified bases, Miyoshi *et al.*, EP 119448, all of
10 which are hereby incorporated by reference in their entirety).

It is further understood, that the present invention provides recombinant bacterial, mammalian, microbial, insect, fungal and plant cells and viral constructs comprising the agents of the present invention. (See, for example, Uses of the Agents of the Invention, Section (a) Plant Constructs and Plant Transformants; Section (b) Fungal Constructs and Fungal
15 Transformants; Section (c) Mammalian Constructs and Transformed Mammalian Cells; Section (d) Insect Constructs and Transformed Insect Cells; and Section (e) Bacterial Constructs and Transformed Bacterial Cells)

Nucleic acid molecules or fragments thereof of the present invention are capable of specifically hybridizing to other nucleic acid molecules under certain circumstances. As used
20 herein, two nucleic acid molecules are said to be capable of specifically hybridizing to one another if the two molecules are capable of forming an anti-parallel, double-stranded nucleic acid structure. A nucleic acid molecule is said to be the "complement" of another nucleic acid molecule if they exhibit complete complementarity. As used herein, molecules are said to

exhibit "complete complementarity" when every nucleotide of one of the molecules is complementary to a nucleotide of the other. Two molecules are said to be "minimally complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under at least conventional "low-stringency" conditions.

- 5 Similarly, the molecules are said to be "complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under conventional "high-stringency" conditions. Conventional stringency conditions are described by Sambrook *et al.*, *Molecular Cloning*, A Laboratory Manual, 2nd Ed., Cold Spring Harbor Press, Cold Spring Harbor, New York (1989) and by Haymes *et al.*, *Nucleic Acid Hybridization, A Practical Approach*, IRL Press, Washington, DC (1985), the entirety of which is herein incorporated by reference. Departures from complete complementarity are therefore permissible, as long as such departures do not completely preclude the capacity of the molecules to form a double-stranded structure. Thus, in order for a nucleic acid molecule to serve as a primer or probe it need only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed.

Appropriate stringency conditions which promote DNA hybridization, for example, 6.0 X sodium chloride/sodium citrate (SSC) at about 45°C, followed by a wash of 2.0 X SSC at 50°C, are known to those skilled in the art or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, the salt concentration in the wash
20 step can be selected from a low stringency of about 2.0 X SSC at 50°C to a high stringency of about 0.2 X SSC at 50°C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22°C, to high stringency conditions at

about 65°C. Both temperature and salt may be varied, or either the temperature or the salt concentration may be held constant while the other variable is changed.

In a preferred embodiment, a nucleic acid of the present invention will specifically hybridize to one or more of the nucleic acid molecules set forth in SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof under moderately stringent conditions, for example at about 2.0 X SSC and about 65°C.

In a particularly preferred embodiment, a nucleic acid of the present invention will include those nucleic acid molecules that specifically hybridize to one or more of the nucleic acid molecules set forth in SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof under high stringency conditions such as 0.2 X SSC and about 65°C.

In one aspect of the present invention, the nucleic acid molecules of the present invention have one or more of the nucleic acid sequences set forth in SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof. In another aspect of the present invention, one or more of the nucleic acid molecules of the present invention share between 100% and 90% sequence identity with one or more of the nucleic acid sequences set forth in SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof. In a further aspect of the present invention, one or more of the nucleic acid molecules of the present invention share between 100% and 95% sequence identity with one or more of the nucleic acid sequences set forth in SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof. In a more preferred aspect of the present invention, one or more of the nucleic acid molecules of the present invention share between 100% and 98% sequence identity with one or more of the nucleic acid sequences set forth in SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof. In an even more preferred aspect of the present invention, one or more of the nucleic acid molecules of the present invention share between 100% and 99%

sequence identity with one or more of the sequences set forth in SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof.

In a further more preferred aspect of the present invention, one or more of the nucleic acid molecules of the present invention exhibit 100% sequence identity with a nucleic acid molecule present within MONN01, SATMON001, SATMON003 through SATMON014, SATMON016, SATMON017, SATMON019 through SATMON031, SATMON033, SATMON034, SATMON~001, SATMONN01, SATMONN04 through SATMONN06, CMz029 through CMz031, CMz033 through CMz037, CMz039 through CMz042, CMz044 through CMz045, CMz047 through CMz050, SOYMON001 through SOYMON038, Soy51 through Soy56, Soy58 through Soy62, Soy65 through Soy71, Soy 73 and Soy76 through Soy77 (Monsanto Company, St. Louis, Missouri U.S.A.).

(i) Nucleic Acid Molecules Encoding Proteins or Fragments Thereof

Nucleic acid molecules of the present invention can comprise sequences that encode a tocopherol synthesis pathway enzyme or fragment thereof. Such tocopherol synthesis pathway enzymes or fragments thereof include homologues of known tocopherol synthesis pathway enzymes in other organisms.

In a preferred embodiment of the present invention, a maize or soybean tocopherol synthesis pathway enzyme or fragment thereof of the present invention is a homologue of another plant tocopherol synthesis pathway enzyme. In another preferred embodiment of the present invention, a maize or soybean tocopherol synthesis pathway enzyme or fragment thereof of the present invention is a homologue of a fungal tocopherol synthesis pathway enzyme. In another preferred embodiment of the present invention, a maize or soybean tocopherol synthesis pathway enzyme or fragment thereof of the present invention is a homologue of a bacterial

tocopherol synthesis pathway enzyme. In another preferred embodiment of the present invention, a soybean tocopherol synthesis pathway enzyme or fragment thereof of the present invention is a homologue of a maize tocopherol synthesis pathway enzyme. In another preferred embodiment of the present invention, a maize tocopherol synthesis pathway enzyme homologue or fragment thereof of the present invention is a homologue of a soybean tocopherol synthesis pathway enzyme. In another preferred embodiment of the present invention, a maize or soybean tocopherol synthesis pathway enzyme homologue or fragment thereof of the present invention is a homologue of an *Arabidopsis thaliana* tocopherol synthesis pathway enzyme.

In a preferred embodiment of the present invention, the nucleic molecule of the present invention encodes a maize or soybean tocopherol synthesis pathway enzyme or fragment thereof where a maize or soybean tocopherol synthesis pathway enzyme exhibits a BLAST probability score of greater than 1E-12, preferably a BLAST probability score of between about 1E-30 and about 1E-12, even more preferably a BLAST probability score of greater than 1E-30 with its homologue.

In another preferred embodiment of the present invention, the nucleic acid molecule encoding a maize or soybean tocopherol synthesis pathway enzyme or fragment thereof exhibits a % identity with its homologue of between about 25% and about 40%, more preferably of between about 40 and about 70%, even more preferably of between about 70% and about 90% and even more preferably between about 90% and 99%. In another preferred embodiment of the present invention, a maize or soybean tocopherol synthesis pathway enzyme or fragments thereof exhibits a % identity with its homologue of 100%.

In a preferred embodiment of the present invention, the nucleic molecule of the present invention encodes a maize or soybean tocopherol synthesis pathway enzyme or fragment thereof

where a maize or soybean tocopherol synthesis pathway enzyme exhibits a BLAST score of greater than 120, preferably a BLAST score of between about 1450 and about 120, even more preferably a BLAST score of greater than 1450 with its homologue.

Nucleic acid molecules of the present invention also include non-maize, non-soybean
5 homologues. Preferred non-maize, non-soybean homologues are selected from the group consisting of alfalfa, *Arabidopsis*, barley, *Brassica*, broccoli, cabbage, citrus, cotton, garlic, oat, oilseed rape, onion, canola, flax, an ornamental plant, pea, peanut, pepper, potato, rice, rye, sorghum, strawberry, sugarcane, sugarbeet, tomato, wheat, poplar, pine, fir, eucalyptus, apple, lettuce, lentils, grape, banana, tea, turf grasses, sunflower, oil palm and *Phaseolus*.

In a preferred embodiment, nucleic acid molecules having SEQ ID NO: 1 through SEQ
10 ID NO: 627 or complements and fragments of either can be utilized to obtain such homologues.

The degeneracy of the genetic code, which allows different nucleic acid sequences to code for the same protein or peptide, is known in the literature. (U.S. Patent No. 4,757,006, the entirety of which is herein incorporated by reference).

In an aspect of the present invention, one or more of the nucleic acid molecules of the
15 present invention differ in nucleic acid sequence from those encoding a maize or soybean tocopherol synthesis pathway enzyme or fragment thereof in SEQ ID NO: 1 through SEQ ID NO: 627 due to the degeneracy in the genetic code in that they encode the same tocopherol synthesis pathway enzyme but differ in nucleic acid sequence.

20 In another further aspect of the present invention, one or more of the nucleic acid molecules of the present invention differ in nucleic acid sequence from those encoding a maize or soybean tocopherol synthesis pathway enzyme or fragment thereof in SEQ ID NO: 1 through SEQ ID NO: 627 due to fact that the different nucleic acid sequence encodes a tocopherol

synthesis pathway enzyme having one or more conservative amino acid residue. Examples of conservative substitutions are set forth in Table 1. It is understood that codons capable of coding for such conservative substitutions are known in the art.

Table 1

<u>Original Residue</u>	<u>Conservative Substitutions</u>
Ala	Ser
Arg	Lys
Asn	Gln; His
Asp	Glu
Cys	Ser; Ala
Gln	Asn
Glu	Asp
Gly	Pro
His	Asn; Gln
Ile	Leu; Val
Leu	Ile; Val
Lys	Arg; Gln; Glu
Met	Leu; Ile
Phe	Met; Leu; Tyr
Ser	Thr
Thr	Ser
Trp	Tyr

Tyr

Trp; Phe

Val

Ile; Leu

In a further aspect of the present invention, one or more of the nucleic acid molecules of the present invention differ in nucleic acid sequence from those encoding a maize or soybean tocopherol synthesis pathway enzyme or fragment thereof set forth in SEQ ID NO: 1 through
5 SEQ ID NO: 627 or fragment thereof due to the fact that one or more codons encoding an amino acid has been substituted for a codon that encodes a nonessential substitution of the amino acid originally encoded.

Agents of the present invention include nucleic acid molecules that encode a maize or soybean tocopherol synthesis pathway enzyme or fragment thereof and particularly substantially
10 purified nucleic acid molecules selected from the group consisting of a nucleic acid molecule that encodes a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize dehydroquate synthase enzyme or fragment thereof; a nucleic acid molecule that
15 encodes a soybean dehydroquate dehydratase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize putative dehydroquate dehydratase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean shikimate dehydrogenase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean shikimate kinase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize enolpyruvylshikimate-
20 P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean chorismate synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or

soybean chorismate mutase enzyme or fragment thereof; a nucleic acid molecule that encodes a
maize tyrosine transaminase enzyme or fragment thereof; a nucleic acid molecule that encodes a
maize or soybean putative tyrosine transaminase enzyme or fragment thereof; a nucleic acid
molecule that encodes a maize or soybean transaminase A enzyme or fragment thereof; a nucleic
5 acid molecule that encodes a soybean putative transaminase A enzyme or fragment thereof; a
nucleic acid molecule that encodes a maize or soybean 4-hydroxyphenylpyruvate dioxygenase
enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean
homogentisic acid dioxygenase enzyme or fragment thereof; a nucleic acid molecule that
encodes a maize or soybean geranylgeranylpyrophosphate synthase enzyme or fragment thereof.

Non-limiting examples of such nucleic acid molecules of the present invention are
nucleic acid molecules comprising: SEQ ID NO: 1 through SEQ ID NO: 627 or fragment thereof
that encode for a plant tocopherol synthesis pathway enzyme or fragment thereof, SEQ ID NO: 1
through SEQ ID NO: 97 and SEQ ID NO: 100 through SEQ ID NO: 146 or fragment thereof that
encodes for a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or fragment
thereof; SEQ ID NO: 98 through SEQ ID NO: 99 and SEQ ID NO: 147 through SEQ ID NO:
15 152 or fragment thereof that encodes for a maize or soybean putative deoxyarabiono-
heptulosonate-P-synthase enzyme or fragment thereof; SEQ ID NO: 153 through SEQ ID NO:
157 or fragment thereof that encodes for a maize dehydroquinase synthase enzyme or fragment
thereof; SEQ ID NO: 160 or fragment thereof that encodes for a soybean dehydroquinase
20 dehydratase enzyme or fragment thereof; SEQ ID NO: 158 through SEQ ID NO: 159 or fragment
thereof that encodes for a maize putative dehydroquinase dehydratase enzyme or fragment
thereof; SEQ ID NO: 158 through SEQ ID NO: 159 and SEQ ID NO: 160 or fragment thereof
that encodes for a maize or soybean shikimate dehydrogenase enzyme or fragment thereof; SEQ

ID NO: 161 through SEQ ID NO: 179 and SEQ ID NO: 180 through SEQ ID NO: 183 or
fragment thereof that encodes for a maize or soybean shikimate kinase enzyme or fragment
thereof; SEQ ID NO: 184 through SEQ ID NO: 198 or fragment thereof that encodes for a maize
enolpyruvylshikimate-P-synthase enzyme or fragment thereof; SEQ ID NO: 199 through SEQ ID
5 NO: 231 and SEQ ID NO: 232 through SEQ ID NO: 255 or fragment thereof that encodes for a
maize or soybean chorismate synthase enzyme or fragment thereof; SEQ ID NO: 256 through
SEQ ID NO: 277 and SEQ ID NO: 278 through SEQ ID NO: 284 or fragment thereof that
encodes for a maize or soybean chorismate mutase enzyme or fragment thereof; SEQ ID NO:
285 through SEQ ID NO: 286 or fragment thereof that encodes for a maize tyrosine transaminase
enzyme or fragment thereof; SEQ ID NO: 287 through SEQ ID NO: 292 and SEQ ID NO: 293
through SEQ ID NO: 300 or fragment thereof that encodes for a maize or soybean putative
tyrosine transaminase enzyme or fragment thereof; SEQ ID NO: 301 through SEQ ID NO: 474
and SEQ ID NO: 475 through SEQ ID NO: 581 or fragment thereof that encodes for a maize or
soybean transaminase A enzyme or fragment thereof; SEQ ID NO: 582 through SEQ ID NO:
15 597 or fragment thereof that encodes for a soybean putative transaminase A enzyme or fragment
thereof; SEQ ID NO: 598 through SEQ ID NO: 600 and SEQ ID NO: 601 through SEQ ID NO:
607 or fragment thereof that encodes for a maize or soybean 4-hydroxyphenylpyruvate
dioxygenase enzyme or fragment thereof; SEQ ID NO: 608 through SEQ ID NO: 615 and SEQ
ID NO: 616 through SEQ ID NO: 621 or fragment thereof that encodes for a maize or soybean
20 homogentisic acid dioxygenase enzyme or fragment thereof; SEQ ID NO: 622 through SEQ ID
NO: 624 and SEQ ID NO: 625 through SEQ ID NO: 627 or fragment thereof that encodes for a
maize or soybean geranylgeranylpyrophosphate synthase enzyme or fragment thereof.

A nucleic acid molecule of the present invention can also encode an homologue of a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a maize dehydroquinase synthase enzyme or fragment thereof; a soybean dehydroquinase
5 dehydratase enzyme or fragment thereof; a maize putative dehydroquinase dehydratase enzyme or fragment thereof; a maize or soybean shikimate dehydrogenase enzyme or fragment thereof; a maize or soybean shikimate kinase enzyme or fragment thereof; a maize enolpyruvylshikimate-P-synthase enzyme or fragment thereof; a maize or soybean chorismate synthase enzyme or fragment thereof; a maize or soybean chorismate mutase enzyme or fragment thereof; a maize tyrosine transaminase enzyme or fragment thereof; a maize or soybean putative tyrosine
10 transaminase enzyme or fragment thereof; a maize or soybean transaminase A enzyme or fragment thereof; a soybean putative transaminase A enzyme or fragment thereof; a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment thereof; a maize or soybean homogentisic acid dioxygenase enzyme or fragment thereof; and a maize or soybean geranylgeranylpyrophosphate synthase enzyme or fragment thereof. As used herein a
15 homologue protein molecule or fragment thereof is a counterpart protein molecule or fragment thereof in a second species (*e.g.*, maize copalyl diphosphate synthase is a homologue of *Arabidopsis* copalyl diphosphate synthase).

(ii) Nucleic Acid Molecule Markers and Probes

20 One aspect of the present invention concerns markers that include nucleic acid molecules SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof or fragments of either that can act as markers or other nucleic acid molecules of the present invention that can act as markers. Genetic markers of the present invention include "dominant" or "codominant" markers

“Codominant markers” reveal the presence of two or more alleles (two per diploid individual) at a locus. “Dominant markers” reveal the presence of only a single allele per locus. The presence of the dominant marker phenotype (e.g., a band of DNA) is an indication that one allele is present in either the homozygous or heterozygous condition. The absence of the dominant marker phenotype (e.g. absence of a DNA band) is merely evidence that “some other” undefined allele is present. In the case of populations where individuals are predominantly homozygous and loci are predominately dimorphic, dominant and codominant markers can be equally valuable. As populations become more heterozygous and multi-allelic, codominant markers often become more informative of the genotype than dominant markers. Marker molecules can be, for example, capable of detecting polymorphisms such as single nucleotide polymorphisms (SNPs).

SNPs are single base changes in genomic DNA sequence. They occur at greater frequency and are spaced with a greater uniformity throughout a genome than other reported forms of polymorphism. The greater frequency and uniformity of SNPs means that there is greater probability that such a polymorphism will be found near or in a genetic locus of interest than would be the case for other polymorphisms. SNPs are located in protein-coding regions and noncoding regions of a genome. Some of these SNPs may result in defective or variant protein expression (e.g., as a results of mutations or defective splicing). Analysis (genotyping) of characterized SNPs can require only a plus/minus assay rather than a lengthy measurement, permitting easier automation.

SNPs can be characterized using any of a variety of methods. Such methods include the direct or indirect sequencing of the site, the use of restriction enzymes (Botstein *et al.*, *Am. J. Hum. Genet.* 32:314-331 (1980), the entirety of which is herein incorporated reference;

Konieczny and Ausubel, *Plant J.* 4:403-410 (1993), the entirety of which is herein incorporated by reference), enzymatic and chemical mismatch assays (Myers *et al.*, *Nature* 313:495-498 (1985), the entirety of which is herein incorporated by reference), allele-specific PCR (Newton *et al.*, *Nucl. Acids Res.* 17:2503-2516 (1989), the entirety of which is herein incorporated by reference; Wu *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:2757-2760 (1989), the entirety of which is herein incorporated by reference), ligase chain reaction (Barany, *Proc. Natl. Acad. Sci. (U.S.A.)* 88:189-193 (1991), the entirety of which is herein incorporated by reference), single-strand conformation polymorphism analysis (Labrune *et al.*, *Am. J. Hum. Genet.* 48: 1115-1120 (1991), the entirety of which is herein incorporated by reference), primer-directed nucleotide incorporation assays (Kuppuswami *et al.*, *Proc. Natl. Acad. Sci. USA* 88:1143-1147 (1991), the entirety of which is herein incorporated by reference), dideoxy fingerprinting (Sarkar *et al.*, *Genomics* 13:441-443 (1992), the entirety of which is herein incorporated by reference), solid-phase ELISA-based oligonucleotide ligation assays (Nikiforov *et al.*, *Nucl. Acids Res.* 22:4167-4175 (1994), the entirety of which is herein incorporated by reference), oligonucleotide fluorescence-quenching assays (Livak *et al.*, *PCR Methods Appl.* 4:357-362 (1995), the entirety of which is herein incorporated by reference), 5'-nuclease allele-specific hybridization TaqMan assay (Livak *et al.*, *Nature Genet.* 9:341-342 (1995), the entirety of which is herein incorporated by reference), template-directed dye-terminator incorporation (TDI) assay (Chen and Kwok, *Nucl. Acids Res.* 25:347-353 (1997), the entirety of which is herein incorporated by reference), allele-specific molecular beacon assay (Tyagi *et al.*, *Nature Biotech.* 16: 49-53 (1998), the entirety of which is herein incorporated by reference), PinPoint assay (Haff and Smirnov, *Genome Res.* 7: 378-388 (1997), the entirety of which is herein incorporated by reference) and

dCAPS analysis (Neff *et al.*, *Plant J.* 14:387-392 (1998), the entirety of which is herein incorporated by reference).

Additional markers, such as AFLP markers, RFLP markers and RAPD markers, can be utilized (Walton, *Seed World* 22-29 (July, 1993), the entirety of which is herein incorporated by reference; Burow and Blake, *Molecular Dissection of Complex Traits*, 13-29, Paterson (ed.), CRC Press, New York (1988), the entirety of which is herein incorporated by reference). DNA markers can be developed from nucleic acid molecules using restriction endonucleases, the PCR and/or DNA sequence information. RFLP markers result from single base changes or insertions/deletions. These codominant markers are highly abundant in plant genomes, have a medium level of polymorphism and are developed by a combination of restriction endonuclease digestion and Southern blotting hybridization. CAPS are similarly developed from restriction nuclease digestion but only of specific PCR products. These markers are also codominant, have a medium level of polymorphism and are highly abundant in the genome. The CAPS result from single base changes and insertions/deletions.

Another marker type, RAPDs, are developed from DNA amplification with random primers and result from single base changes and insertions/deletions in plant genomes. They are dominant markers with a medium level of polymorphisms and are highly abundant. AFLP markers require using the PCR on a subset of restriction fragments from extended adapter primers. These markers are both dominant and codominant are highly abundant in genomes and exhibit a medium level of polymorphism.

SSRs require DNA sequence information. These codominant markers result from repeat length changes, are highly polymorphic and do not exhibit as high a degree of abundance in the genome as CAPS, AFLPs and RAPDs SNPs also require DNA sequence information. These

codominant markers result from single base substitutions. They are highly abundant and exhibit a medium of polymorphism (Rafalski *et al.*, In: *Nonmammalian Genomic Analysis*, Birren and Lai (ed.), Academic Press, San Diego, CA, pp. 75-134 (1996), the entirety of which is herein incorporated by reference). It is understood that a nucleic acid molecule of the present invention
5 may be used as a marker.

A PCR probe is a nucleic acid molecule capable of initiating a polymerase activity while in a double-stranded structure to with another nucleic acid. Various methods for determining the structure of PCR probes and PCR techniques exist in the art. Computer generated searches using programs such as Primer3 (www-genome.wi.mit.edu/cgi-bin/primer/primer3.cgi), STSPipeline (www-genome.wi.mit.edu/cgi-bin/www-STSPipeline), or GeneUp (Pesole *et al.*, *BioTechniques* 25:112-123 (1998) the entirety of which is herein incorporated by reference), for example, can be used to identify potential PCR primers.

It is understood that a fragment of one or more of the nucleic acid molecules of the present invention may be a probe and specifically a PCR probe.

(b) Protein and Peptide Molecules

A class of agents comprises one or more of the protein or fragments thereof or peptide molecules encoded by SEQ ID NO: 1 through SEQ ID NO: 627 or one or more of the protein or fragment thereof and peptide molecules encoded by other nucleic acid agents of the present invention. As used herein, the term "protein molecule" or "peptide molecule" includes any
20 molecule that comprises five or more amino acids. It is well known in the art that proteins may undergo modification, including post-translational modifications, such as, but not limited to, disulfide bond formation, glycosylation, phosphorylation, or oligomerization. Thus, as used herein, the term "protein molecule" or "peptide molecule" includes any protein molecule that is

modified by any biological or non-biological process. The terms "amino acid" and "amino acids" refer to all naturally occurring L-amino acids. This definition is meant to include norleucine, ornithine, homocysteine and homoserine.

Non-limiting examples of the protein or fragment thereof of the present invention include

5 a maize or soybean tocopherol synthesis pathway enzyme or fragment thereof; a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a maize dehydroquate synthase enzyme or fragment thereof; a soybean dehydroquate dehydratase enzyme or fragment thereof; a maize putative dehydroquate dehydratase enzyme or fragment thereof; a maize or soybean shikimate dehydrogenase enzyme or fragment thereof; a maize or soybean shikimate kinase enzyme or fragment thereof; a maize enolpyruvylshikimate-P-synthase enzyme or fragment thereof; a maize or soybean chorismate synthase enzyme or fragment thereof; a maize or soybean chorismate mutase enzyme or fragment thereof; a maize tyrosine transaminase enzyme or fragment thereof; a maize or soybean putative tyrosine transaminase enzyme or fragment thereof; a maize or soybean transaminase A enzyme or fragment thereof; a soybean putative transaminase A enzyme or fragment thereof; a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment thereof; a maize or soybean homogentisic acid dioxygenase enzyme or fragment thereof; and a maize or soybean geranylgeranylpyrophosphate synthase enzyme or fragment thereof.

20 Non-limiting examples of the protein or fragment molecules of the present invention are a tocopherol synthesis pathway enzyme or fragment thereof encoded by: SEQ ID NO: 1 through SEQ ID NO: 627 or fragment thereof that encode for a tocopherol synthesis pathway enzyme or fragment thereof, SEQ ID NO: 1 through SEQ ID NO: 97 and SEQ ID NO: 100 through SEQ ID

NO: 146 or fragment thereof that encodes for a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; SEQ ID NO: 98 through SEQ ID NO: 99 and SEQ ID NO: 147 through SEQ ID NO: 152 or fragment thereof that encodes for a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; SEQ ID NO: 153 through SEQ ID NO: 157 or fragment thereof that encodes for a maize dehydroquinase synthase enzyme or fragment thereof; SEQ ID NO: 160 or fragment thereof that encodes for a soybean dehydroquinase dehydratase enzyme or fragment thereof; SEQ ID NO: 158 through SEQ ID NO: 159 or fragment thereof that encodes for a maize putative dehydroquinase dehydratase enzyme or fragment thereof; SEQ ID NO: 158 through SEQ ID NO: 159 and SEQ ID NO: 160 or fragment thereof that encodes for a maize or soybean shikimate dehydrogenase enzyme or fragment thereof; SEQ ID NO: 161 through SEQ ID NO: 179 and SEQ ID NO: 180 through SEQ ID NO: 183 or fragment thereof that encodes for a maize or soybean shikimate kinase enzyme or fragment thereof; SEQ ID NO: 184 through SEQ ID NO: 198 or fragment thereof that encodes for a maize enolpyruvylshikimate-P-synthase enzyme or fragment thereof; SEQ ID NO: 199 through SEQ ID NO: 231 and SEQ ID NO: 232 through SEQ ID NO: 255 or fragment thereof that encodes for a maize or soybean chorismate synthase enzyme or fragment thereof; SEQ ID NO: 256 through SEQ ID NO: 277 and SEQ ID NO: 278 through SEQ ID NO: 284 or fragment thereof that encodes for a maize or soybean chorismate mutase enzyme or fragment thereof; SEQ ID NO: 285 through SEQ ID NO: 286 or fragment thereof that encodes for a maize tyrosine transaminase enzyme or fragment thereof; SEQ ID NO: 287 through SEQ ID NO: 292 and SEQ ID NO: 293 through SEQ ID NO: 300 or fragment thereof that encodes for a maize or soybean putative tyrosine transaminase enzyme or fragment thereof; SEQ ID NO: 301 through SEQ ID NO: 474 and SEQ ID NO: 475 through SEQ ID NO: 581 or fragment thereof that encodes for a

maize or soybean transaminase A enzyme or fragment thereof; SEQ ID NO: 582 through SEQ ID NO: 597 or fragment thereof that encodes for a soybean putative transaminase A enzyme or fragment thereof; SEQ ID NO: 598 through SEQ ID NO: 600 and SEQ ID NO: 601 through SEQ ID NO: 607 or fragment thereof that encodes for a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment thereof; SEQ ID NO: 608 through SEQ ID NO: 615 and SEQ ID NO: 616 through SEQ ID NO: 621 or fragment thereof that encodes for a maize or soybean homogentisic acid dioxygenase enzyme or fragment thereof; SEQ ID NO: 622 through SEQ ID NO: 624 and SEQ ID NO: 625 through SEQ ID NO: 627 or fragment thereof that encodes for a maize or soybean geranylgeranylpyrophosphate synthase enzyme or fragment thereof.

One or more of the protein or fragment of peptide molecules may be produced via chemical synthesis, or more preferably, by expressing in a suitable bacterial or eucaryotic host. Suitable methods for expression are described by Sambrook *et al.*, (In: *Molecular Cloning, A Laboratory Manual, 2nd Edition, Cold Spring Harbor Press, Cold Spring Harbor, New York* (1989)), or similar texts. For example, the protein may be expressed in, for example, Uses of the Agents of the Invention, Section (a) Plant Constructs and Plant Transformants; Section (b) Fungal Constructs and Fungal Transformants; Section (c) Mammalian Constructs and Transformed Mammalian Cells; Section (d) Insect Constructs and Transformed Insect Cells; and Section (e) Bacterial Constructs and Transformed Bacterial Cells.

A "protein fragment" is a peptide or polypeptide molecule whose amino acid sequence comprises a subset of the amino acid sequence of that protein. A protein or fragment thereof that comprises one or more additional peptide regions not derived from that protein is a "fusion" protein. Such molecules may be derivatized to contain carbohydrate or other moieties (such as

keyhole limpet hemocyanin, etc.). Fusion protein or peptide molecules of the present invention are preferably produced via recombinant means.

Another class of agents comprise protein or peptide molecules or fragments or fusions thereof encoded by SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof in which
5 conservative, non-essential or non-relevant amino acid residues have been added, replaced or deleted. Computerized means for designing modifications in protein structure are known in the art (Dahiyat and Mayo, *Science* 278:82-87 (1997), the entirety of which is herein incorporated by reference).

The protein molecules of the present invention include plant homologue proteins. An example of such a homologue is a homologue protein of a non-maize or non-soybean plant species, that include but not limited to alfalfa, *Arabidopsis*, barley, *Brassica*, broccoli, cabbage, citrus, cotton, garlic, oat, oilseed rape, onion, canola, flax, an ornamental plant, pea, peanut, pepper, potato, rice, rye, sorghum, strawberry, sugarcane, sugarbeet, tomato, wheat, poplar, pine, fir, eucalyptus, apple, lettuce, lentils, grape, banana, tea, turf grasses, sunflower, oil palm, *Phaseolus* etc. Particularly preferred non-maize or non-soybean for use for the isolation of
10 homologs would include, *Arabidopsis*, barley, cotton, oat, oilseed rape, rice, canola, ornamentals, sugarcane, sugarbeet, tomato, potato, wheat and turf grasses. Such a homologue can be obtained by any of a variety of methods. Most preferably, as indicated above, one or more of the disclosed sequences (SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof) will be
20 used to define a pair of primers that may be used to isolate the homologue-encoding nucleic acid molecules from any desired species. Such molecules can be expressed to yield homologues by recombinant means.

(c) Antibodies

One aspect of the present invention concerns antibodies, single-chain antigen binding molecules, or other proteins that specifically bind to one or more of the protein or peptide molecules of the present invention and their homologues, fusions or fragments. Such antibodies
5 may be used to quantitatively or qualitatively detect the protein or peptide molecules of the present invention. As used herein, an antibody or peptide is said to "specifically bind" to a protein or peptide molecule of the present invention if such binding is not competitively inhibited by the presence of non-related molecules.

Nucleic acid molecules that encode all or part of the protein of the present invention can be expressed, via recombinant means, to yield protein or peptides that can in turn be used to elicit antibodies that are capable of binding the expressed protein or peptide. Such antibodies may be used in immunoassays for that protein. Such protein-encoding molecules, or their fragments may be a "fusion" molecule (i.e., a part of a larger nucleic acid molecule) such that, upon expression, a fusion protein is produced. It is understood that any of the nucleic acid molecules of the present invention may be expressed, via recombinant means, to yield proteins or peptides encoded by these nucleic acid molecules.

The antibodies that specifically bind proteins and protein fragments of the present invention may be polyclonal or monoclonal and may comprise intact immunoglobulins, or antigen binding portions of immunoglobulins fragments (such as $(F(ab'))_2$), or single-chain
20 immunoglobulins producible, for example, via recombinant means. It is understood that practitioners are familiar with the standard resource materials which describe specific conditions and procedures for the construction, manipulation and isolation of antibodies (*see*, for example,

Harlow and Lane, In: *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, New York (1988), the entirety of which is herein incorporated by reference).

Murine monoclonal antibodies are particularly preferred. BALB/c mice are preferred for this purpose, however, equivalent strains may also be used. The animals are preferably

5 immunized with approximately 25 μ g of purified protein (or fragment thereof) that has been emulsified in a suitable adjuvant (such as TiterMax adjuvant (Vaxcel, Norcross, GA)).

Immunization is preferably conducted at two intramuscular sites, one intraperitoneal site and one subcutaneous site at the base of the tail. An additional i.v. injection of approximately 25 μ g of antigen is preferably given in normal saline three weeks later. After approximately 11 days following the second injection, the mice may be bled and the blood screened for the presence of anti-protein or peptide antibodies. Preferably, a direct binding Enzyme-Linked Immunoassay (ELISA) is employed for this purpose.

More preferably, the mouse having the highest antibody titer is given a third i.v. injection of approximately 25 μ g of the same protein or fragment. The splenic leukocytes from this animal may be recovered 3 days later and then permitted to fuse, most preferably, using polyethylene glycol, with cells of a suitable myeloma cell line (such as, for example, the P3X63Ag8.653 myeloma cell line). Hybridoma cells are selected by culturing the cells under "HAT" (hypoxanthine-aminopterin-thymine) selection for about one week. The resulting clones may then be screened for their capacity to produce monoclonal antibodies ("mAbs"), preferably
20 by direct ELISA.

In one embodiment, anti-protein or peptide monoclonal antibodies are isolated using a fusion of a protein or peptide of the present invention, or conjugate of a protein or peptide of the present invention, as immunogens. Thus, for example, a group of mice can be immunized using

a fusion protein emulsified in Freund's complete adjuvant (*e.g.* approximately 50 μ g of antigen per immunization). At three week intervals, an identical amount of antigen is emulsified in Freund's incomplete adjuvant and used to immunize the animals. Ten days following the third immunization, serum samples are taken and evaluated for the presence of antibody. If antibody
5 titers are too low, a fourth booster can be employed. Polysera capable of binding the protein or peptide can also be obtained using this method.

In a preferred procedure for obtaining monoclonal antibodies, the spleens of the above-described immunized mice are removed, disrupted and immune splenocytes are isolated over a ficoll gradient. The isolated splenocytes are fused, using polyethylene glycol with BALB/c-derived HGPRT (hypoxanthine guanine phosphoribosyl transferase) deficient P3x63xAg8.653 plasmacytoma cells. The fused cells are plated into 96 well microtiter plates and screened for hybridoma fusion cells by their capacity to grow in culture medium supplemented with hypothanthine, aminopterin and thymidine for approximately 2-3 weeks.

Hybridoma cells that arise from such incubation are preferably screened for their capacity to produce an immunoglobulin that binds to a protein of interest. An indirect ELISA may be used for this purpose. In brief, the supernatants of hybridomas are incubated in microtiter wells that contain immobilized protein. After washing, the titer of bound immunoglobulin can be determined using, for example, a goat anti-mouse antibody conjugated to horseradish peroxidase. After additional washing, the amount of immobilized enzyme is determined (for example
20 through the use of a chromogenic substrate). Such screening is performed as quickly as possible after the identification of the hybridoma in order to ensure that a desired clone is not overgrown by non-secreting neighbor cells. Desirably, the fusion plates are screened several times since the rates of hybridoma growth vary. In a preferred sub-embodiment, a different antigenic form may

be used to screen the hybridoma. Thus, for example, the splenocytes may be immunized with one immunogen, but the resulting hybridomas can be screened using a different immunogen. It is understood that any of the protein or peptide molecules of the present invention may be used to raise antibodies.

5 As discussed below, such antibody molecules or their fragments may be used for diagnostic purposes. Where the antibodies are intended for diagnostic purposes, it may be desirable to derivatize them, for example with a ligand group (such as biotin) or a detectable marker group (such as a fluorescent group, a radioisotope or an enzyme).

10 The ability to produce antibodies that bind the protein or peptide molecules of the present invention permits the identification of mimetic compounds of those molecules. A "mimetic compound" is a compound that is not that compound, or a fragment of that compound, but which nonetheless exhibits an ability to specifically bind to antibodies directed against that compound.

15 It is understood that any of the agents of the present invention can be substantially purified and/or be biologically active and/or recombinant.

Uses of the Agents of the Invention

20 Nucleic acid molecules and fragments thereof of the present invention may be employed to obtain other nucleic acid molecules from the same species (e.g., ESTs or fragment thereof from maize may be utilized to obtain other nucleic acid molecules from maize). Such nucleic acid molecules include the nucleic acid molecules that encode the complete coding sequence of a protein and promoters and flanking sequences of such molecules. In addition, such nucleic acid molecules include nucleic acid molecules that encode for other isozymes or gene family members. Such molecules can be readily obtained by using the above-described nucleic acid

molecules or fragments thereof to screen cDNA or genomic libraries obtained from maize or soybean. Methods for forming such libraries are well known in the art.

5 Nucleic acid molecules and fragments thereof of the present invention may also be employed to obtain nucleic acid homologues. Such homologues include the nucleic acid molecule of other plants or other organisms (*e.g.*, alfalfa, *Arabidopsis*, barley, *Brassica*, broccoli, cabbage, citrus, cotton, garlic, oat, oilseed rape, onion, canola, flax, an ornamental plant, pea, peanut, pepper, potato, rice, rye, sorghum, strawberry, sugarcane, sugarbeet, tomato, wheat, poplar, pine, fir, eucalyptus, apple, lettuce, lentils, grape, banana, tea, turf grasses, sunflower, oil palm, *Phaseolus*, etc.) including the nucleic acid molecules that encode, in whole or in part, protein homologues of other plant species or other organisms, sequences of genetic elements such as promoters and transcriptional regulatory elements. Such molecules can be readily obtained by using the above-described nucleic acid molecules or fragments thereof to screen cDNA or genomic libraries obtained from such plant species. Methods for forming such libraries are well known in the art. Such homologue molecules may differ in their nucleotide sequences from those found in one or more of SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof because complete complementarity is not needed for stable hybridization. The nucleic acid molecules of the present invention therefore also include molecules that, although capable of specifically hybridizing with the nucleic acid molecules may lack "complete complementarity."

20 Any of a variety of methods may be used to obtain one or more of the above-described nucleic acid molecules (Zamechik *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 83:4143-4146 (1986), the entirety of which is herein incorporated by reference; Goodchild *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 85:5507-5511 (1988), the entirety of which is herein incorporated by reference; Wickstrom *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 85:1028-1032 (1988), the entirety of which is

herein incorporated by reference; Holt *et al.*, *Molec. Cell. Biol.* 8:963-973 (1988), the entirety of which is herein incorporated by reference; Gerwitz *et al.*, *Science* 242:1303-1306 (1988), the entirety of which is herein incorporated by reference; Anfossi *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:3379-3383 (1989), the entirety of which is herein incorporated by reference; Becker
5 *et al.*, *EMBO J.* 8:3685-3691 (1989); the entirety of which is herein incorporated by reference).

Automated nucleic acid synthesizers may be employed for this purpose. In lieu of such synthesis, the disclosed nucleic acid molecules may be used to define a pair of primers that can be used with the polymerase chain reaction (Mullis *et al.*, *Cold Spring Harbor Symp. Quant. Biol.* 51:263-273 (1986); Erlich *et al.*, European Patent 50,424; European Patent 84,796; European Patent 258,017; European Patent 237,362; Mullis, European Patent 201,184; Mullis *et al.*, U.S. Patent No. 4,683,202; Erlich, U.S. Patent No. 4,582,788; and Saiki *et al.*, U.S. Patent No. 4,683,194, all of which are herein incorporated by reference in their entirety) to amplify and obtain any desired nucleic acid molecule or fragment.

Promoter sequence(s) and other genetic elements, including but not limited to transcriptional regulatory flanking sequences, associated with one or more of the disclosed nucleic acid sequences can also be obtained using the disclosed nucleic acid sequence provided herein. In one embodiment, such sequences are obtained by incubating EST nucleic acid molecules or preferably fragments thereof with members of genomic libraries (*e.g.* maize and soybean) and recovering clones that hybridize to the EST nucleic acid molecule or fragment
20 thereof. In a second embodiment, methods of "chromosome walking," or inverse PCR may be used to obtain such sequences (Frohman *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 85:8998-9002 (1988); Ohara *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:5673-5677 (1989); Pang *et al.*, *Biotechniques* 22:1046-1048 (1977); Huang *et al.*, *Methods Mol. Biol.* 69:89-96 (1997); Huang

et al., *Method Mol. Biol.* 67:287-294 (1997); Benkel *et al.*, *Genet. Anal.* 13:123-127 (1996); Hartl *et al.*, *Methods Mol. Biol.* 58:293-301 (1996), all of which are herein incorporated by reference in their entirety).

The nucleic acid molecules of the present invention may be used to isolate promoters of cell enhanced, cell specific, tissue enhanced, tissue specific, developmentally or environmentally regulated expression profiles. Isolation and functional analysis of the 5' flanking promoter sequences of these genes from genomic libraries, for example, using genomic screening methods and PCR techniques would result in the isolation of useful promoters and transcriptional regulatory elements. These methods are known to those of skill in the art and have been described (See, for example, Birren *et al.*, *Genome Analysis: Analyzing DNA*, 1, (1997), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., the entirety of which is herein incorporated by reference). Promoters obtained utilizing the nucleic acid molecules of the present invention could also be modified to affect their control characteristics. Examples of such modifications would include but are not limited to enhanced sequences as reported in Uses of the Agents of the Invention, Section (a) Plant Constructs and Plant Transformants. Such genetic elements could be used to enhance gene expression of new and existing traits for crop improvements.

In one sub-aspect, such an analysis is conducted by determining the presence and/or identity of polymorphism(s) by one or more of the nucleic acid molecules of the present invention and more preferably one or more of the EST nucleic acid molecule or fragment thereof which are associated with a phenotype, or a predisposition to that phenotype.

Any of a variety of molecules can be used to identify such polymorphism(s). In one embodiment, one or more of the EST nucleic acid molecules (or a sub-fragment thereof) may be

employed as a marker nucleic acid molecule to identify such polymorphism(s). Alternatively, such polymorphisms can be detected through the use of a marker nucleic acid molecule or a marker protein that is genetically linked to (i.e., a polynucleotide that co-segregates with) such polymorphism(s).

5 In an alternative embodiment, such polymorphisms can be detected through the use of a marker nucleic acid molecule that is physically linked to such polymorphism(s). For this purpose, marker nucleic acid molecules comprising a nucleotide sequence of a polynucleotide located within 1mb of the polymorphism(s) and more preferably within 100kb of the polymorphism(s) and most preferably within 10kb of the polymorphism(s) can be employed.

10 The genomes of animals and plants naturally undergo spontaneous mutation in the course of their continuing evolution (Gusella, *Ann. Rev. Biochem.* 55:831-854 (1986)). A “polymorphism” is a variation or difference in the sequence of the gene or its flanking regions that arises in some of the members of a species. The variant sequence and the “original” sequence co-exist in the species’ population. In some instances, such co-existence is in stable or
15 quasi-stable equilibrium.

A polymorphism is thus said to be “allelic,” in that, due to the existence of the polymorphism, some members of a species may have the original sequence (i.e., the original “allele”) whereas other members may have the variant sequence (i.e., the variant “allele”). In the simplest case, only one variant sequence may exist and the polymorphism is thus said to be di-
20 allelic. In other cases, the species’ population may contain multiple alleles and the polymorphism is termed tri-allelic, etc. A single gene may have multiple different unrelated polymorphisms. For example, it may have a di-allelic polymorphism at one site and a multi-allelic polymorphism at another site.

The variation that defines the polymorphism may range from a single nucleotide variation to the insertion or deletion of extended regions within a gene. In some cases, the DNA sequence variations are in regions of the genome that are characterized by short tandem repeats (STRs) that include tandem di- or tri-nucleotide repeated motifs of nucleotides. Polymorphisms characterized by such tandem repeats are referred to as "variable number tandem repeat" ("VNTR") polymorphisms. VNTRs have been used in identity analysis (Weber, U.S. Patent No. 5,075,217; Armour *et al.*, *FEBS Lett.* 307:113-115 (1992); Jones *et al.*, *Eur. J. Haematol.* 39:144-147 (1987); Horn *et al.*, PCT Patent Application WO91/14003; Jeffreys, European Patent Application 370,719; Jeffreys, U.S. Patent No. 5,175,082; Jeffreys *et al.*, *Amer. J. Hum. Genet.* 39:11-24 (1986); Jeffreys *et al.*, *Nature* 316:76-79 (1985); Gray *et al.*, *Proc. R. Acad. Soc. Lond.* 243:241-253 (1991); Moore *et al.*, *Genomics* 10:654-660 (1991); Jeffreys *et al.*, *Anim. Genet.* 18:1-15 (1987); Hillel *et al.*, *Anim. Genet.* 20:145-155 (1989); Hillel *et al.*, *Genet.* 124:783-789 (1990), all of which are herein incorporated by reference in their entirety).

The detection of polymorphic sites in a sample of DNA may be facilitated through the use of nucleic acid amplification methods. Such methods specifically increase the concentration of polynucleotides that span the polymorphic site, or include that site and sequences located either distal or proximal to it. Such amplified molecules can be readily detected by gel electrophoresis or other means.

The most preferred method of achieving such amplification employs the polymerase chain reaction ("PCR") (Mullis *et al.*, *Cold Spring Harbor Symp. Quant. Biol.* 51:263-273 (1986); Erlich *et al.*, European Patent Appln. 50,424; European Patent Appln. 84,796; European Patent Application 258,017; European Patent Appln. 237,362; Mullis, European Patent Appln. 201,184; Mullis *et al.*, U.S. Patent No. 4,683,202; Erlich, U.S. Patent No. 4,582,788; and Saiki *et*

al., U.S. Patent No. 4,683,194), using primer pairs that are capable of hybridizing to the proximal sequences that define a polymorphism in its double-stranded form.

In lieu of PCR, alternative methods, such as the "Ligase Chain Reaction" ("LCR") may be used (Barany, *Proc. Natl. Acad. Sci. (U.S.A.)* 88:189-193 (1991), the entirety of which is
5 herein incorporated by reference). LCR uses two pairs of oligonucleotide probes to exponentially amplify a specific target. The sequences of each pair of oligonucleotides is selected to permit the pair to hybridize to abutting sequences of the same strand of the target. Such hybridization forms a substrate for a template-dependent ligase. As with PCR, the resulting products thus serve as a template in subsequent cycles and an exponential amplification of the desired sequence is obtained.

LCR can be performed with oligonucleotides having the proximal and distal sequences of the same strand of a polymorphic site. In one embodiment, either oligonucleotide will be designed to include the actual polymorphic site of the polymorphism. In such an embodiment, the reaction conditions are selected such that the oligonucleotides can be ligated together only if the target molecule either contains or lacks the specific nucleotide that is complementary to the polymorphic site present on the oligonucleotide. Alternatively, the oligonucleotides may be selected such that they do not include the polymorphic site (see, Segev, PCT Application WO 90/01069, the entirety of which is herein incorporated by reference).

The "Oligonucleotide Ligation Assay" ("OLA") may alternatively be employed
20 (Landegren *et al.*, *Science* 241:1077-1080 (1988), the entirety of which is herein incorporated by reference). The OLA protocol uses two oligonucleotides which are designed to be capable of hybridizing to abutting sequences of a single strand of a target. OLA, like LCR, is particularly

suited for the detection of point mutations. Unlike LCR, however, OLA results in "linear" rather than exponential amplification of the target sequence.

Nickerson *et al.*, have described a nucleic acid detection assay that combines attributes of PCR and OLA (Nickerson *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 87:8923-8927 (1990), the
5 entirety of which is herein incorporated by reference). In this method, PCR is used to achieve the exponential amplification of target DNA, which is then detected using OLA. In addition to requiring multiple and separate, processing steps, one problem associated with such combinations is that they inherit all of the problems associated with PCR and OLA.

Schemes based on ligation of two (or more) oligonucleotides in the presence of nucleic
10 acid having the sequence of the resulting "di-oligonucleotide", thereby amplifying the di-oligonucleotide, are also known (Wu *et al.*, *Genomics* 4:560-569 (1989), the entirety of which is herein incorporated by reference) and may be readily adapted to the purposes of the present invention.

Other known nucleic acid amplification procedures, such as allele-specific oligomers,
15 branched DNA technology, transcription-based amplification systems, or isothermal amplification methods may also be used to amplify and analyze such polymorphisms (Malek *et al.*, U.S. Patent No. 5,130,238; Davey *et al.*, European Patent Application 329,822; Schuster *et al.*, U.S. Patent No. 5,169,766; Miller *et al.*, PCT Patent Application WO 89/06700; Kwoh *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:1173-1177 (1989); Gingeras *et al.*, PCT Patent Application
20 WO 88/10315; Walker *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 89:392-396 (1992), all of which are herein incorporated by reference in their entirety).

The identification of a polymorphism can be determined in a variety of ways. By correlating the presence or absence of it in a plant with the presence or absence of a phenotype, it

is possible to predict the phenotype of that plant. If a polymorphism creates or destroys a restriction endonuclease cleavage site, or if it results in the loss or insertion of DNA (e.g., a VNTR polymorphism), it will alter the size or profile of the DNA fragments that are generated by digestion with that restriction endonuclease. As such, individuals that possess a variant
5 sequence can be distinguished from those having the original sequence by restriction fragment analysis. Polymorphisms that can be identified in this manner are termed "restriction fragment length polymorphisms" ("RFLPs"). RFLPs have been widely used in human and plant genetic analyses (Glassberg, UK Patent Application 2135774; Skolnick *et al.*, *Cytogen. Cell Genet.* 32:58-67 (1982); Botstein *et al.*, *Ann. J. Hum. Genet.* 32:314-331 (1980); Fischer *et al.*, (PCT Application WO90/13668); Uhlen, PCT Application WO90/11369).

Polymorphisms can also be identified by Single Strand Conformation Polymorphism (SSCP) analysis. SSCP is a method capable of identifying most sequence variations in a single strand of DNA, typically between 150 and 250 nucleotides in length (Elles, *Methods in Molecular Medicine: Molecular Diagnosis of Genetic Diseases*, Humana Press (1996), the
15 entirety of which is herein incorporated by reference); Orita *et al.*, *Genomics* 5:874-879 (1989), the entirety of which is herein incorporated by reference). Under denaturing conditions a single strand of DNA will adopt a conformation that is uniquely dependent on its sequence conformation. This conformation usually will be different, even if only a single base is changed. Most conformations have been reported to alter the physical configuration or size sufficiently to
20 be detectable by electrophoresis. A number of protocols have been described for SSCP including, but not limited to, Lee *et al.*, *Anal. Biochem.* 205:289-293 (1992), the entirety of which is herein incorporated by reference; Suzuki *et al.*, *Anal. Biochem.* 192:82-84 (1991), the entirety of which is herein incorporated by reference; Lo *et al.*, *Nucleic Acids Research* 20:1005-

1009 (1992), the entirety of which is herein incorporated by reference; Sarkar *et al.*, *Genomics* 13:441-443 (1992), the entirety of which is herein incorporated by reference. It is understood that one or more of the nucleic acids of the present invention, may be utilized as markers or probes to detect polymorphisms by SSCP analysis.

5 Polymorphisms may also be found using a DNA fingerprinting technique called amplified fragment length polymorphism (AFLP), which is based on the selective PCR amplification of restriction fragments from a total digest of genomic DNA to profile that DNA (Vos *et al.*, *Nucleic Acids Res.* 23:4407-4414 (1995), the entirety of which is herein incorporated by reference). This method allows for the specific co-amplification of high numbers of
10 restriction fragments, which can be visualized by PCR without knowledge of the nucleic acid sequence.

AFLP employs basically three steps. Initially, a sample of genomic DNA is cut with restriction enzymes and oligonucleotide adapters are ligated to the restriction fragments of the DNA. The restriction fragments are then amplified using PCR by using the adapter and
15 restriction sequence as target sites for primer annealing. The selective amplification is achieved by the use of primers that extend into the restriction fragments, amplifying only those fragments in which the primer extensions match the nucleotide flanking the restriction sites. These amplified fragments are then visualized on a denaturing polyacrylamide gel.

AFLP analysis has been performed on *Salix* (Beismann *et al.*, *Mol. Ecol.* 6:989-993
20 (1997), the entirety of which is herein incorporated by reference), *Acinetobacter* (Janssen *et al.*, *Int. J. Syst. Bacteriol.* 47:1179-1187 (1997), the entirety of which is herein incorporated by reference), *Aeromonas popoffi* (Huys *et al.*, *Int. J. Syst. Bacteriol.* 47:1165-1171 (1997), the entirety of which is herein incorporated by reference), rice (McCouch *et al.*, *Plant Mol. Biol.*

35:89-99 (1997), the entirety of which is herein incorporated by reference; Nandi *et al.*, *Mol. Gen. Genet.* 255:1-8 (1997), the entirety of which is herein incorporated by reference; Cho *et al.*, *Genome* 39:373-378 (1996), the entirety of which is herein incorporated by reference), barley (*Hordeum vulgare*)(Simons *et al.*, *Genomics* 44:61-70 (1997), the entirety of which is herein incorporated by reference; Waugh *et al.*, *Mol. Gen. Genet.* 255:311-321 (1997), the entirety of which is herein incorporated by reference; Qi *et al.*, *Mol. Gen. Genet.* 254:330-336 (1997), the entirety of which is herein incorporated by reference; Becker *et al.*, *Mol. Gen. Genet.* 249:65-73 (1995), the entirety of which is herein incorporated by reference), potato (Van der Voort *et al.*, *Mol. Gen. Genet.* 255:438-447 (1997), the entirety of which is herein incorporated by reference; Meksem *et al.*, *Mol. Gen. Genet.* 249:74-81 (1995), the entirety of which is herein incorporated by reference), *Phytophthora infestans* (Van der Lee *et al.*, *Fungal Genet. Biol.* 21:278-291 (1997), the entirety of which is herein incorporated by reference), *Bacillus anthracis* (Keim *et al.*, *J. Bacteriol.* 179:818-824 (1997), the entirety of which is herein incorporated by reference), *Astragalus cremnophylax* (Travis *et al.*, *Mol. Ecol.* 5:735-745 (1996), the entirety of which is herein incorporated by reference), *Arabidopsis* (Cnops *et al.*, *Mol. Gen. Genet.* 253:32-41 (1996), the entirety of which is herein incorporated by reference), *Escherichia coli* (Lin *et al.*, *Nucleic Acids Res.* 24:3649-3650 (1996), the entirety of which is herein incorporated by reference), *Aeromonas* (Huys *et al.*, *Int. J. Syst. Bacteriol.* 46:572-580 (1996), the entirety of which is herein incorporated by reference), nematode (Folkertsma *et al.*, *Mol. Plant Microbe Interact.* 9:47-54 (1996), the entirety of which is herein incorporated by reference), tomato (Thomas *et al.*, *Plant J.* 8:785-794 (1995), the entirety of which is herein incorporated by reference) and human (Latorra *et al.*, *PCR Methods Appl.* 3:351-358 (1994), the entirety of which is herein incorporated by reference). AFLP analysis has also been used for fingerprinting mRNA (Money *et al.*, *Nucleic*

Acids Res. 24:2616-2617 (1996), the entirety of which is herein incorporated by reference;

Bachem *et al.*, *Plant J.* 9:745-753 (1996), the entirety of which is herein incorporated by

reference). It is understood that one or more of the nucleic acids of the present invention, may be utilized as markers or probes to detect polymorphisms by AFLP analysis or for fingerprinting

5 RNA.

Polymorphisms may also be found using random amplified polymorphic DNA (RAPD) (Williams *et al.*, *Nucl. Acids Res.* 18:6531-6535 (1990), the entirety of which is herein incorporated by reference) and cleaveable amplified polymorphic sequences (CAPS) (Lyamichev *et al.*, *Science* 260:778-783 (1993), the entirety of which is herein incorporated by reference). It is understood that one or more of the nucleic acid molecules of the present invention, may be utilized as markers or probes to detect polymorphisms by RAPD or CAPS analysis.

Through genetic mapping, a fine scale linkage map can be developed using DNA markers and, then, a genomic DNA library of large-sized fragments can be screened with molecular markers linked to the desired trait. Molecular markers are advantageous for agronomic traits that are otherwise difficult to tag, such as resistance to pathogens, insects and nematodes, tolerance to abiotic stress, quality parameters and quantitative traits such as high yield potential.

The essential requirements for marker-assisted selection in a plant breeding program are: (1) the marker(s) should co-segregate or be closely linked with the desired trait; (2) an efficient means of screening large populations for the molecular marker(s) should be available; and (3) the screening technique should have high reproducibility across laboratories and preferably be economical to use and be user-friendly.

The genetic linkage of marker molecules can be established by a gene mapping model such as, without limitation, the flanking marker model reported by Lander and Botstein, *Genetics*

121:185-199 (1989) and the interval mapping, based on maximum likelihood methods described by Lander and Botstein, *Genetics* 121:185-199 (1989) and implemented in the software package MAPMAKER/QTL (Lincoln and Lander, *Mapping Genes Controlling Quantitative Traits Using MAPMAKER/QTL*, Whitehead Institute for Biomedical Research, Massachusetts, (1990).

- 5 Additional software includes Qgene, Version 2.23 (1996), Department of Plant Breeding and Biometry, 266 Emerson Hall, Cornell University, Ithaca, NY, the manual of which is herein incorporated by reference in its entirety). Use of Qgene software is a particularly preferred approach.

A maximum likelihood estimate (MLE) for the presence of a marker is calculated, together with an MLE assuming no QTL effect, to avoid false positives. A \log_{10} of an odds ratio (LOD) is then calculated as: $LOD = \log_{10}(\text{MLE for the presence of a QTL} / \text{MLE given no linked QTL})$.

The LOD score essentially indicates how much more likely the data are to have arisen assuming the presence of a QTL than in its absence. The LOD threshold value for avoiding a false positive with a given confidence, say 95%, depends on the number of markers and the length of the genome. Graphs indicating LOD thresholds are set forth in Lander and Botstein, *Genetics* 121:185-199 (1989) the entirety of which is herein incorporated by reference and further described by Arús and Moreno-González, *Plant Breeding*, Hayward *et al.*, (eds.) Chapman & Hall, London, pp. 314-331 (1993), the entirety of which is herein incorporated by
20 reference.

Additional models can be used. Many modifications and alternative approaches to interval mapping have been reported, including the use non-parametric methods (Kruglyak and Lander, *Genetics* 139:1421-1428 (1995), the entirety of which is herein incorporated by

reference). Multiple regression methods or models can be also be used, in which the trait is regressed on a large number of markers (Jansen, *Biometrics in Plant Breeding*, van Oijen and Jansen (eds.), Proceedings of the Ninth Meeting of the Eucarpia Section Biometrics in Plant Breeding, The Netherlands, pp. 116-124 (1994); Weber and Wricke, *Advances in Plant Breeding*, Blackwell, Berlin, 16 (1994), both of which is herein incorporated by reference in their entirety).

5 Procedures combining interval mapping with regression analysis, whereby the phenotype is regressed onto a single putative QTL at a given marker interval and at the same time onto a number of markers that serve as 'cofactors,' have been reported by Jansen and Stam, *Genetics* 136:1447-1455 (1994), the entirety of which is herein incorporated by reference and Zeng, *Genetics* 136:1457-1468 (1994) the entirety of which is herein incorporated by reference.

10 Generally, the use of cofactors reduces the bias and sampling error of the estimated QTL positions (Utz and Melchinger, *Biometrics in Plant Breeding*, van Oijen and Jansen (eds.) Proceedings of the Ninth Meeting of the Eucarpia Section Biometrics in Plant Breeding, The Netherlands, pp.195-204 (1994), the entirety of which is herein incorporated by reference,

15 thereby improving the precision and efficiency of QTL mapping (Zeng, *Genetics* 136:1457-1468 (1994)). These models can be extended to multi-environment experiments to analyze genotype-environment interactions (Jansen *et al.*, *Theo. Appl. Genet.* 91:33-37 (1995), the entirety of which is herein incorporated by reference).

Selection of an appropriate mapping populations is important to map construction. The

20 choice of appropriate mapping population depends on the type of marker systems employed (Tanksley *et al.*, *Molecular mapping plant chromosomes. Chromosome structure and function: Impact of new concepts*, Gustafson and Appels (eds.), Plenum Press, New York, pp 157-173 (1988), the entirety of which is herein incorporated by reference). Consideration must be given

to the source of parents (adapted vs. exotic) used in the mapping population. Chromosome pairing and recombination rates can be severely disturbed (suppressed) in wide crosses (adapted x exotic) and generally yield greatly reduced linkage distances. Wide crosses will usually provide segregating populations with a relatively large array of polymorphisms when compared to progeny in a narrow cross (adapted x adapted).

An F_2 population is the first generation of selfing after the hybrid seed is produced. Usually a single F_1 plant is selfed to generate a population segregating for all the genes in Mendelian (1:2:1) fashion. Maximum genetic information is obtained from a completely classified F_2 population using a codominant marker system (Mather, *Measurement of Linkage in Heredity*, Methuen and Co., (1938), the entirety of which is herein incorporated by reference). In the case of dominant markers, progeny tests (e.g. F_3 , BCF_2) are required to identify the heterozygotes, thus making it equivalent to a completely classified F_2 population. However, this procedure is often prohibitive because of the cost and time involved in progeny testing. Progeny testing of F_2 individuals is often used in map construction where phenotypes do not consistently reflect genotype (e.g. disease resistance) or where trait expression is controlled by a QTL. Segregation data from progeny test populations (e.g. F_3 or BCF_2) can be used in map construction. Marker-assisted selection can then be applied to cross progeny based on marker-trait map associations (F_2 , F_3), where linkage groups have not been completely disassociated by recombination events (i.e., maximum disequilibrium).

Recombinant inbred lines (RIL) (genetically related lines; usually $>F_5$, developed from continuously selfing F_2 lines towards homozygosity) can be used as a mapping population. Information obtained from dominant markers can be maximized by using RIL because all loci are homozygous or nearly so. Under conditions of tight linkage (i.e., about $<10\%$

recombination), dominant and co-dominant markers evaluated in RIL populations provide more information per individual than either marker type in backcross populations (Reiter *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 89:1477-1481 (1992), the entirety of which is herein incorporated by reference). However, as the distance between markers becomes larger (i.e., loci become more independent), the information in RIL populations decreases dramatically when compared to codominant markers.

Backcross populations (e.g., generated from a cross between a successful variety (recurrent parent) and another variety (donor parent) carrying a trait not present in the former) can be utilized as a mapping population. A series of backcrosses to the recurrent parent can be made to recover most of its desirable traits. Thus a population is created consisting of individuals nearly like the recurrent parent but each individual carries varying amounts or mosaic of genomic regions from the donor parent. Backcross populations can be useful for mapping dominant markers if all loci in the recurrent parent are homozygous and the donor and recurrent parent have contrasting polymorphic marker alleles (Reiter *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 89:1477-1481 (1992)). Information obtained from backcross populations using either codominant or dominant markers is less than that obtained from F_2 populations because one, rather than two, recombinant gametes are sampled per plant. Backcross populations, however, are more informative (at low marker saturation) when compared to RILs as the distance between linked loci increases in RIL populations (i.e. about 15% recombination). Increased recombination can be beneficial for resolution of tight linkages, but may be undesirable in the construction of maps with low marker saturation.

Near-isogenic lines (NIL) created by many backcrosses to produce an array of individuals that are nearly identical in genetic composition except for the trait or genomic region under

interrogation can be used as a mapping population. In mapping with NILs, only a portion of the polymorphic loci are expected to map to a selected region.

Bulk segregant analysis (BSA) is a method developed for the rapid identification of linkage between markers and traits of interest (Michelmore *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 88:9828-9832 (1991), the entirety of which is herein incorporated by reference). In BSA, two bulked DNA samples are drawn from a segregating population originating from a single cross. These bulks contain individuals that are identical for a particular trait (resistant or susceptible to particular disease) or genomic region but arbitrary at unlinked regions (i.e. heterozygous). Regions unlinked to the target region will not differ between the bulked samples of many individuals in BSA.

It is understood that one or more of the nucleic acid molecules of the present invention may be used as molecular markers. It is also understood that one or more of the protein molecules of the present invention may be used as molecular markers.

In accordance with this aspect of the present invention, a sample nucleic acid is obtained from plants cells or tissues. Any source of nucleic acid may be used. Preferably, the nucleic acid is genomic DNA. The nucleic acid is subjected to restriction endonuclease digestion. For example, one or more nucleic acid molecule or fragment thereof of the present invention can be used as a probe in accordance with the above-described polymorphic methods. The polymorphism obtained in this approach can then be cloned to identify the mutation at the coding region which alters the protein's structure or regulatory region of the gene which affects its expression level.

In an aspect of the present invention, one or more of the nucleic molecules of the present invention are used to determine the level (i.e., the concentration of mRNA in a sample, etc.) in a

plant (preferably maize or soybean) or pattern (i.e., the kinetics of expression, rate of decomposition, stability profile, etc.) of the expression of a protein encoded in part or whole by one or more of the nucleic acid molecule of the present invention (collectively, the "Expression Response" of a cell or tissue). As used herein, the Expression Response manifested by a cell or tissue is said to be "altered" if it differs from the Expression Response of cells or tissues of plants not exhibiting the phenotype. To determine whether an Expression Response is altered, the Expression Response manifested by the cell or tissue of the plant exhibiting the phenotype is compared with that of a similar cell or tissue sample of a plant not exhibiting the phenotype. As will be appreciated, it is not necessary to re-determine the Expression Response of the cell or tissue sample of plants not exhibiting the phenotype each time such a comparison is made; rather, the Expression Response of a particular plant may be compared with previously obtained values of normal plants. As used herein, the phenotype of the organism is any of one or more characteristics of an organism (e.g. disease resistance, pest tolerance, environmental tolerance such as tolerance to abiotic stress, male sterility, quality improvement or yield etc.). A change in genotype or phenotype may be transient or permanent. Also as used herein, a tissue sample is any sample that comprises more than one cell. In a preferred aspect, a tissue sample comprises cells that share a common characteristic (e.g. derived from root, seed, flower, leaf, stem or pollen etc.).

In one aspect of the present invention, an evaluation can be conducted to determine whether a particular mRNA molecule is present. One or more of the nucleic acid molecules of the present invention, preferably one or more of the EST nucleic acid molecules or fragments thereof of the present invention are utilized to detect the presence or quantity of the mRNA species. Such molecules are then incubated with cell or tissue extracts of a plant under

conditions sufficient to permit nucleic acid hybridization. The detection of double-stranded probe-mRNA hybrid molecules is indicative of the presence of the mRNA; the amount of such hybrid formed is proportional to the amount of mRNA. Thus, such probes may be used to ascertain the level and extent of the mRNA production in a plant's cells or tissues. Such nucleic acid hybridization may be conducted under quantitative conditions (thereby providing a numerical value of the amount of the mRNA present). Alternatively, the assay may be conducted as a qualitative assay that indicates either that the mRNA is present, or that its level exceeds a user set, predefined value.

A principle of *in situ* hybridization is that a labeled, single-stranded nucleic acid probe will hybridize to a complementary strand of cellular DNA or RNA and, under the appropriate conditions, these molecules will form a stable hybrid. When nucleic acid hybridization is combined with histological techniques, specific DNA or RNA sequences can be identified within a single cell. An advantage of *in situ* hybridization over more conventional techniques for the detection of nucleic acids is that it allows an investigator to determine the precise spatial population (Angerer *et al.*, *Dev. Biol.* 101:477-484 (1984), the entirety of which is herein incorporated by reference; Angerer *et al.*, *Dev. Biol.* 112:157-166 (1985), the entirety of which is herein incorporated by reference; Dixon *et al.*, *EMBO J.* 10:1317-1324 (1991), the entirety of which is herein incorporated by reference). *In situ* hybridization may be used to measure the steady-state level of RNA accumulation. It is a sensitive technique and RNA sequences present in as few as 5-10 copies per cell can be detected (Hardin *et al.*, *J. Mol. Biol.* 202:417-431 (1989), the entirety of which is herein incorporated by reference). A number of protocols have been devised for *in situ* hybridization, each with tissue preparation, hybridization and washing conditions (Meyerowitz, *Plant Mol. Biol. Rep.* 5:242-250 (1987), the entirety of which is herein

incorporated by reference; Cox and Goldberg, In: *Plant Molecular Biology: A Practical Approach*, Shaw (ed.), pp 1-35, IRL Press, Oxford (1988), the entirety of which is herein incorporated by reference; Raikhel *et al.*, *In situ RNA hybridization in plant tissues*, In: *Plant Molecular Biology Manual*, vol. B9:1-32, Kluwer Academic Publisher, Dordrecht, Belgium
5 (1989), the entirety of which is herein incorporated by reference).

In situ hybridization also allows for the localization of proteins within a tissue or cell (Wilkinson, *In Situ Hybridization*, Oxford University Press, Oxford (1992), the entirety of which is herein incorporated by reference; Langdale, *In Situ Hybridization In: The Maize Handbook*, Freeling and Walbot (eds.), pp 165-179, Springer-Verlag, New York (1994), the entirety of which is herein incorporated by reference). It is understood that one or more of the molecules of the present invention, preferably one or more of the EST nucleic acid molecules or fragments thereof of the present invention or one or more of the antibodies of the present invention may be utilized to detect the level or pattern of a tocopherol synthesis pathway enzyme or mRNA thereof by *in situ* hybridization.

Fluorescent *in situ* hybridization allows the localization of a particular DNA sequence along a chromosome which is useful, among other uses, for gene mapping, following chromosomes in hybrid lines or detecting chromosomes with translocations, transversions or deletions. *In situ* hybridization has been used to identify chromosomes in several plant species (Griffor *et al.*, *Plant Mol. Biol.* 17:101-109 (1991), the entirety of which is herein incorporated
20 by reference; Gustafson *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 87:1899-1902 (1990), herein incorporated by reference; Mukai and Gill, *Genome* 34:448-452 (1991), the entirety of which is herein incorporated by reference; Schwarzacher and Heslop-Harrison, *Genome* 34:317-323 (1991); Wang *et al.*, *Jpn. J. Genet.* 66:313-316 (1991), the entirety of which is herein

incorporated by reference; Parra and Windle, *Nature Genetics* 5:17-21 (1993), the entirety of which is herein incorporated by reference). It is understood that the nucleic acid molecules of the present invention may be used as probes or markers to localize sequences along a chromosome.

Another method to localize the expression of a molecule is tissue printing. Tissue printing provides a way to screen, at the same time on the same membrane many tissue sections from different plants or different developmental stages. Tissue-printing procedures utilize films designed to immobilize proteins and nucleic acids. In essence, a freshly cut section of a tissue is pressed gently onto nitrocellulose paper, nylon membrane or polyvinylidene difluoride membrane. Such membranes are commercially available (e.g. Millipore, Bedford, Massachusetts U.S.A.). The contents of the cut cell transfer onto the membrane and the contents and are immobilized to the membrane. The immobilized contents form a latent print that can be visualized with appropriate probes. When a plant tissue print is made on nitrocellulose paper, the cell walls leave a physical print that makes the anatomy visible without further treatment (Varner and Taylor, *Plant Physiol.* 91:31-33 (1989), the entirety of which is herein incorporated by reference).

Tissue printing on substrate films is described by Daoust, *Exp. Cell Res.* 12:203-211 (1957), the entirety of which is herein incorporated by reference, who detected amylase, protease, ribonuclease and deoxyribonuclease in animal tissues using starch, gelatin and agar films. These techniques can be applied to plant tissues (Yomo and Taylor, *Planta* 112:35-43 (1973); the entirety of which is herein incorporated by reference; Harris and Chrispeels, *Plant Physiol.* 56:292-299 (1975), the entirety of which is herein incorporated by reference). Advances in membrane technology have increased the range of applications of Daoust's tissue-printing techniques allowing (Cassab and Varner, *J. Cell. Biol.* 105:2581-2588 (1987), the entirety of

which is herein incorporated by reference) the histochemical localization of various plant enzymes and deoxyribonuclease on nitrocellulose paper and nylon (Spruce *et al.*, *Phytochemistry* 26:2901-2903 (1987), the entirety of which is herein incorporated by reference; Barres *et al.*, *Neuron* 5:527-544 (1990), the entirety of which is herein incorporated by reference; Reid and Pont-Lezica, *Tissue Printing: Tools for the Study of Anatomy, Histochemistry and Gene Expression*, Academic Press, New York, New York (1992), the entirety of which is herein incorporated by reference; Reid *et al.*, *Plant Physiol.* 93:160-165 (1990), the entirety of which is herein incorporated by reference; Ye *et al.*, *Plant J.* 1:175-183 (1991), the entirety of which is herein incorporated by reference).

It is understood that one or more of the molecules of the present invention, preferably one or more of the EST nucleic acid molecules or fragments thereof of the present invention or one or more of the antibodies of the present invention may be utilized to detect the presence or quantity of a tocopherol synthesis pathway enzyme by tissue printing.

Further it is also understood that any of the nucleic acid molecules of the present invention may be used as marker nucleic acids and or probes in connection with methods that require probes or marker nucleic acids. As used herein, a probe is an agent that is utilized to determine an attribute or feature (e.g. presence or absence, location, correlation, etc.) of a molecule, cell, tissue or plant. As used herein, a marker nucleic acid is a nucleic acid molecule that is utilized to determine an attribute or feature (e.g., presence or absence, location, correlation, etc.) or a molecule, cell, tissue or plant.

A microarray-based method for high-throughput monitoring of plant gene expression may be utilized to measure gene-specific hybridization targets. This 'chip'-based approach involves using microarrays of nucleic acid molecules as gene-specific hybridization targets to

quantitatively measure expression of the corresponding plant genes (Schena *et al.*, *Science* 270:467-470 (1995), the entirety of which is herein incorporated by reference; Shalon, Ph.D. Thesis, Stanford University (1996), the entirety of which is herein incorporated by reference). Every nucleotide in a large sequence can be queried at the same time. Hybridization can be used
5 to efficiently analyze nucleotide sequences.

Several microarray methods have been described. One method compares the sequences to be analyzed by hybridization to a set of oligonucleotides representing all possible subsequences (Bains and Smith, *J. Theor. Biol.* 135:303-307 (1989), the entirety of which is herein incorporated by reference). A second method hybridizes the sample to an array of
10 oligonucleotide or cDNA molecules. An array consisting of oligonucleotides complementary to subsequences of a target sequence can be used to determine the identity of a target sequence, measure its amount and detect differences between the target and a reference sequence. Nucleic acid molecule microarrays may also be screened with protein molecules or fragments thereof to determine nucleic acid molecules that specifically bind protein molecules or fragments thereof.

The microarray approach may be used with polypeptide targets (U.S. Patent No. 5,445,934; U.S. Patent No. 5,143,854; U.S. Patent No. 5,079,600; U.S. Patent No. 4,923,901, all of which are herein incorporated by reference in their entirety). Essentially, polypeptides are synthesized on a substrate (microarray) and these polypeptides can be screened with either
15 protein molecules or fragments thereof or nucleic acid molecules in order to screen for either
20 protein molecules or fragments thereof or nucleic acid molecules that specifically bind the target polypeptides. (Fodor *et al.*, *Science* 251:767-773 (1991), the entirety of which is herein incorporated by reference). It is understood that one or more of the nucleic acid molecules or

protein or fragments thereof of the present invention may be utilized in a microarray based method.

In a preferred embodiment of the present invention microarrays may be prepared that comprise nucleic acid molecules where such nucleic acid molecules encode at least one,

5 preferably at least two, more preferably at least three tocopherol synthesis pathway enzymes, more preferably at least four tocopherol synthesis pathway enzymes, more preferably at least five tocopherol synthesis pathway enzymes, more preferably at least six tocopherol synthesis pathway enzymes, more preferably at least seven tocopherol synthesis pathway enzymes, more preferably at least eight tocopherol synthesis pathway enzymes, more preferably at least nine tocopherol synthesis pathway enzymes, more preferably at least ten tocopherol synthesis pathway enzymes, more preferably at least eleven tocopherol synthesis pathway enzymes, more preferably at least twelve tocopherol synthesis pathway enzymes, more preferably at least thirteen tocopherol synthesis pathway enzymes, more preferably at least fourteen tocopherol synthesis pathway enzymes, more preferably at least fifteen tocopherol synthesis pathway enzymes, more preferably at least sixteen tocopherol synthesis pathway enzymes, and even more preferably at least seventeen tocopherol synthesis pathway enzymes. In a preferred embodiment the nucleic acid molecules are selected from the group consisting of a nucleic acid molecule that encodes a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize dehydroquininate synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a soybean dehydroquininate dehydratase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize putative dehydroquininate dehydratase enzyme or fragment thereof; a nucleic acid molecule

that encodes a maize or soybean shikimate dehydrogenase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean shikimate kinase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize enolpyruvylshikimate-P-synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean chorismate synthase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean chorismate mutase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize tyrosine transaminase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean putative tyrosine transaminase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean transaminase A enzyme or fragment thereof; a nucleic acid molecule that encodes a soybean putative transaminase A enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or fragment thereof; a nucleic acid molecule that encodes a maize or soybean homogentisic acid dioxygenase enzyme or fragment thereof; and a nucleic acid molecule that encodes a maize or soybean geranylgeranylpyrophosphate synthase enzyme or fragment thereof.

Site directed mutagenesis may be utilized to modify nucleic acid sequences, particularly as it is a technique that allows one or more of the amino acids encoded by a nucleic acid molecule to be altered (e.g. a threonine to be replaced by a methionine). Three basic methods for site directed mutagenesis are often employed. These are cassette mutagenesis (Wells *et al.*, *Gene* 34:315-323 (1985), the entirety of which is herein incorporated by reference), primer extension (Gilliam *et al.*, *Gene* 12:129-137 (1980), the entirety of which is herein incorporated by reference; Zoller and Smith, *Methods Enzymol.* 100:468-500 (1983), the entirety of which is herein incorporated by reference; Dalbadie-McFarland *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 79:6409-6413 (1982), the entirety of which is herein incorporated by reference) and methods

based upon PCR (Scharf *et al.*, *Science* 233:1076-1078 (1986), the entirety of which is herein incorporated by reference; Higuchi *et al.*, *Nucleic Acids Res.* 16:7351-7367 (1988), the entirety of which is herein incorporated by reference). Site directed mutagenesis approaches are also described in European Patent 0 385 962, the entirety of which is herein incorporated by reference; European Patent 0 359 472, the entirety of which is herein incorporated by reference; and PCT Patent Application WO 93/07278, the entirety of which is herein incorporated by reference.

Site directed mutagenesis strategies have been applied to plants for both *in vitro* as well as *in vivo* site directed mutagenesis (Lanz *et al.*, *J. Biol. Chem.* 266:9971-9976 (1991), the entirety of which is herein incorporated by reference; Kovgan and Zhdanov, *Biotekhnologiya* 5:148-154; No. 207160n, Chemical Abstracts 110:225 (1989), the entirety of which is herein incorporated by reference; Ge *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:4037-4041 (1989), the entirety of which is herein incorporated by reference; Zhu *et al.*, *J. Biol. Chem.* 271:18494-18498 (1996), the entirety of which is herein incorporated by reference; Chu *et al.*, *Biochemistry* 33:6150-6157 (1994), the entirety of which is herein incorporated by reference; Small *et al.*, *EMBO J.* 11:1291-1296 (1992), the entirety of which is herein incorporated by reference; Cho *et al.*, *Mol. Biotechnol.* 8:13-16 (1997), the entirety of which is herein incorporated by reference; Kita *et al.*, *J. Biol. Chem.* 271:26529-26535 (1996), the entirety of which is herein incorporated by reference; Jin *et al.*, *Mol. Microbiol.* 7:555-562 (1993), the entirety of which is herein incorporated by reference; Hatfield and Vierstra, *J. Biol. Chem.* 267:14799-14803 (1992), the entirety of which is herein incorporated by reference; Zhao *et al.*, *Biochemistry* 31:5093-5099 (1992), the entirety of which is herein incorporated by reference).

Any of the nucleic acid molecules of the present invention may either be modified by site directed mutagenesis or used as, for example, nucleic acid molecules that are used to target other nucleic acid molecules for modification. It is understood that mutants with more than one altered nucleotide can be constructed using techniques that practitioners are familiar with such as isolating restriction fragments and ligating such fragments into an expression vector (see, for example, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (1989)).

Sequence-specific DNA-binding proteins play a role in the regulation of transcription. The isolation of recombinant cDNAs encoding these proteins facilitates the biochemical analysis of their structural and functional properties. Genes encoding such DNA-binding proteins have been isolated using classical genetics (Vollbrecht *et al.*, *Nature* 350: 241-243 (1991), the entirety of which is herein incorporated by reference) and molecular biochemical approaches, including the screening of recombinant cDNA libraries with antibodies (Landschulz *et al.*, *Genes Dev.* 2:786-800 (1988), the entirety of which is herein incorporated by reference) or DNA probes (Bodner *et al.*, *Cell* 55:505-518 (1988), the entirety of which is herein incorporated by reference). In addition, an *in situ* screening procedure has been used and has facilitated the isolation of sequence-specific DNA-binding proteins from various plant species (Gilmartin *et al.*, *Plant Cell* 4:839-849 (1992), the entirety of which is herein incorporated by reference; Schindler *et al.*, *EMBO J.* 11:1261-1273 (1992), the entirety of which is herein incorporated by reference). An *in situ* screening protocol does not require the purification of the protein of interest (Vinson *et al.*, *Genes Dev.* 2:801-806 (1988), the entirety of which is herein incorporated by reference; Singh *et al.*, *Cell* 52:415-423 (1988), the entirety of which is herein incorporated by reference).

Two steps may be employed to characterize DNA-protein interactions. The first is to identify promoter fragments that interact with DNA-binding proteins, to titrate binding activity, to determine the specificity of binding and to determine whether a given DNA-binding activity can interact with related DNA sequences (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989)). Electrophoretic mobility-shift assay is a widely used assay. The assay provides a rapid and sensitive method for detecting DNA-binding proteins based on the observation that the mobility of a DNA fragment through a nondenaturing, low-ionic strength polyacrylamide gel is retarded upon association with a DNA-binding protein (Fried and Crother, *Nucleic Acids Res.* 9:6505-6525 (1981), the entirety of which is herein incorporated by reference). When one or more specific binding activities have been identified, the exact sequence of the DNA bound by the protein may be determined. Several procedures for characterizing protein/DNA-binding sites are used, including methylation and ethylation interference assays (Maxam and Gilbert, *Methods Enzymol.* 65:499-560 (1980), the entirety of which is herein incorporated by reference; Wissman and Hillen, *Methods Enzymol.* 208:365-379 (1991), the entirety of which is herein incorporated by reference), footprinting techniques employing DNase I (Galas and Schmitz, *Nucleic Acids Res.* 5:3157-3170 (1978), the entirety of which is herein incorporated by reference), 1,10-phenanthroline-copper ion methods (Sigman *et al.*, *Methods Enzymol.* 208:414-433 (1991), the entirety of which is herein incorporated by reference) and hydroxyl radicals methods (Dixon *et al.*, *Methods Enzymol.* 208:414-433 (1991), the entirety of which is herein incorporated by reference). It is understood that one or more of the nucleic acid molecules of the present invention may be utilized to identify a protein or fragment thereof that specifically binds to a

nucleic acid molecule of the present invention. It is also understood that one or more of the protein molecules or fragments thereof of the present invention may be utilized to identify a nucleic acid molecule that specifically binds to it.

A two-hybrid system is based on the fact that many cellular functions are carried out by proteins, such as transcription factors, that interact (physically) with one another. Two-hybrid systems have been used to probe the function of new proteins (Chien *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 88:9578-9582 (1991) the entirety of which is herein incorporated by reference; Durfee *et al.*, *Genes Dev.* 7:555-569 (1993) the entirety of which is herein incorporated by reference; Choi *et al.*, *Cell* 78:499-512 (1994), the entirety of which is herein incorporated by reference; Kranz *et al.*, *Genes Dev.* 8:313-327 (1994), the entirety of which is herein incorporated by reference).

Interaction mating techniques have facilitated a number of two-hybrid studies of protein-protein interaction. Interaction mating has been used to examine interactions between small sets of tens of proteins (Finley and Brent, *Proc. Natl. Acad. Sci. (U.S.A.)* 91:12098-12984 (1994), the entirety of which is herein incorporated by reference), larger sets of hundreds of proteins (Bendixen *et al.*, *Nucl. Acids Res.* 22:1778-1779 (1994), the entirety of which is herein incorporated by reference) and to comprehensively map proteins encoded by a small genome (Bartel *et al.*, *Nature Genetics* 12:72-77 (1996), the entirety of which is herein incorporated by reference). This technique utilizes proteins fused to the DNA-binding domain and proteins fused to the activation domain. They are expressed in two different haploid yeast strains of opposite mating type and the strains are mated to determine if the two proteins interact. Mating occurs when haploid yeast strains come into contact and result in the fusion of the two haploids into a diploid yeast strain. An interaction can be determined by the activation of a two-hybrid reporter gene in the diploid strain. An advantage of this technique is that it reduces the number of yeast

transformations needed to test individual interactions. It is understood that the protein-protein interactions of protein or fragments thereof of the present invention may be investigated using the two-hybrid system and that any of the nucleic acid molecules of the present invention that encode such proteins or fragments thereof may be used to transform yeast in the two-hybrid system.

(a) Plant Constructs and Plant Transformants

One or more of the nucleic acid molecules of the present invention may be used in plant transformation or transfection. Exogenous genetic material may be transferred into a plant cell and the plant cell regenerated into a whole, fertile or sterile plant. Exogenous genetic material is any genetic material, whether naturally occurring or otherwise, from any source that is capable of being inserted into any organism. Such genetic material may be transferred into either monocotyledons and dicotyledons including, but not limited to maize (pp 63-69), soybean (pp 50-60), *Arabidopsis* (p 45), phaseolus (pp 47-49), peanut (pp 49-50), alfalfa (p 60), wheat (pp 69-71), rice (pp 72-79), oat (pp 80-81), sorghum (p 83), rye (p 84), tritordeum (p 84), millet (p85), fescue (p 85), perennial ryegrass (p 86), sugarcane (p87), cranberry (p101), papaya (pp 101-102), banana (p 103), banana (p 103), muskmelon (p 104), apple (p 104), cucumber (p 105), dendrobium (p 109), gladiolus (p 110), chrysanthemum (p 110), liliacea (p 111), cotton (pp113-114), eucalyptus (p 115), sunflower (p 118), canola (p 118), turfgrass (p121), sugarbeet (p 122), coffee (p 122) and dioscorea (p 122) (Christou, In: *Particle Bombardment for Genetic Engineering of Plants*, Biotechnology Intelligence Unit. Academic Press, San Diego, California (1996), the entirety of which is herein incorporated by reference).

Transfer of a nucleic acid that encodes for a protein can result in overexpression of that protein in a transformed cell or transgenic plant. One or more of the proteins or fragments

thereof encoded by nucleic acid molecules of the present invention may be overexpressed in a transformed cell or transformed plant. Particularly, any of the tocopherol synthesis pathway enzymes or fragments thereof may be overexpressed in a transformed cell or transgenic plant. Such overexpression may be the result of transient or stable transfer of the exogenous genetic material.

Exogenous genetic material may be transferred into a plant cell and the plant cell by the use of a DNA vector or construct designed for such a purpose. Design of such a vector is generally within the skill of the art (See, *Plant Molecular Biology: A Laboratory Manual*, Clark (ed.), Springer, New York (1997), the entirety of which is herein incorporated by reference).

A construct or vector may include a plant promoter to express the protein or protein fragment of choice. A number of promoters which are active in plant cells have been described in the literature. These include the nopaline synthase (NOS) promoter (Ebert *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 84:5745-5749 (1987), the entirety of which is herein incorporated by reference), the octopine synthase (OCS) promoter (which are carried on tumor-inducing plasmids of *Agrobacterium tumefaciens*), the caulimovirus promoters such as the cauliflower mosaic virus (CaMV) 19S promoter (Lawton *et al.*, *Plant Mol. Biol.* 9:315-324 (1987), the entirety of which is herein incorporated by reference) and the CAMV 35S promoter (Odell *et al.*, *Nature* 313:810-812 (1985), the entirety of which is herein incorporated by reference), the figwort mosaic virus 35S-promoter, the light-inducible promoter from the small subunit of ribulose-1,5-bis-phosphate carboxylase (ssRUBISCO), the Adh promoter (Walker *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 84:6624-6628 (1987), the entirety of which is herein incorporated by reference), the sucrose synthase promoter (Yang *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 87:4144-4148 (1990), the entirety of which is herein incorporated by reference), the R gene complex promoter (Chandler *et al.*, *The*

Plant Cell 1:1175-1183 (1989), the entirety of which is herein incorporated by reference) and the chlorophyll a/b binding protein gene promoter, etc. These promoters have been used to create DNA constructs which have been expressed in plants; *see, e.g.*, PCT publication WO 84/02913, herein incorporated by reference in its entirety.

5 Promoters which are known or are found to cause transcription of DNA in plant cells can be used in the present invention. Such promoters may be obtained from a variety of sources such as plants and plant viruses. It is preferred that the particular promoter selected should be capable of causing sufficient expression to result in the production of an effective amount of the tocopherol synthesis pathway enzyme to cause the desired phenotype. In addition to promoters
10 that are known to cause transcription of DNA in plant cells, other promoters may be identified for use in the current invention by screening a plant cDNA library for genes which are selectively or preferably expressed in the target tissues or cells.

For the purpose of expression in source tissues of the plant, such as the leaf, seed, root or stem, it is preferred that the promoters utilized in the present invention have relatively high
15 expression in these specific tissues. For this purpose, one may choose from a number of promoters for genes with tissue- or cell-specific or -enhanced expression. Examples of such promoters reported in the literature include the chloroplast glutamine synthetase GS2 promoter from pea (Edwards *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 87:3459-3463 (1990), herein incorporated by reference in its entirety), the chloroplast fructose-1,6-biphosphatase (FBPase)
20 promoter from wheat (Lloyd *et al.*, *Mol. Gen. Genet.* 225:209-216 (1991), herein incorporated by reference in its entirety), the nuclear photosynthetic ST-LS1 promoter from potato (Stockhaus *et al.*, *EMBO J.* 8:2445-2451 (1989), herein incorporated by reference in its entirety), the serine/threonine kinase (PAL) promoter and the glucoamylase (CHS) promoter from *Arabidopsis*

thaliana. Also reported to be active in photosynthetically active tissues are the ribulose-1,5-bisphosphate carboxylase (RbcS) promoter from eastern larch (*Larix laricina*), the promoter for the *cab* gene, *cab6*, from pine (Yamamoto *et al.*, *Plant Cell Physiol.* 35:773-778 (1994), herein incorporated by reference in its entirety), the promoter for the Cab-1 gene from wheat (Fejes *et al.*, *Plant Mol. Biol.* 15:921-932 (1990), herein incorporated by reference in its entirety), the promoter for the CAB-1 gene from spinach (Lubberstedt *et al.*, *Plant Physiol.* 104:997-1006 (1994), herein incorporated by reference in its entirety), the promoter for the *cab1R* gene from rice (Luan *et al.*, *Plant Cell.* 4:971-981 (1992), the entirety of which is herein incorporated by reference), the pyruvate, orthophosphate dikinase (PPDK) promoter from maize (Matsuoka *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 90: 9586-9590 (1993), herein incorporated by reference in its entirety), the promoter for the tobacco *Lhcb1*2* gene (Cerdan *et al.*, *Plant Mol. Biol.* 33:245-255 (1997), herein incorporated by reference in its entirety), the *Arabidopsis thaliana* SUC2 sucrose-H⁺ symporter promoter (Truernit *et al.*, *Planta.* 196:564-570 (1995), herein incorporated by reference in its entirety) and the promoter for the thylakoid membrane proteins from spinach (*psaD*, *psaF*, *psaE*, *PC*, *FNR*, *atpC*, *atpD*, *cab*, *rbcS*). Other promoters for the chlorophyll a/b-binding proteins may also be utilized in the present invention, such as the promoters for *Lhcb* gene and *PsbP* gene from white mustard (*Sinapis alba*; Kretsch *et al.*, *Plant Mol. Biol.* 28:219-229 (1995), the entirety of which is herein incorporated by reference).

For the purpose of expression in sink tissues of the plant, such as the tuber of the potato plant, the fruit of tomato, or the seed of maize, wheat, rice and barley, it is preferred that the promoters utilized in the present invention have relatively high expression in these specific tissues. A number of promoters for genes with tuber-specific or -enhanced expression are known, including the class I patatin promoter (Bevan *et al.*, *EMBO J.* 8:1899-1906 (1986);

Jefferson *et al.*, *Plant Mol. Biol.* 14:995-1006 (1990), both of which are herein incorporated by reference in its entirety), the promoter for the potato tuber ADPGPP genes, both the large and small subunits, the sucrose synthase promoter (Salanoubat and Belliard, *Gene*. 60:47-56 (1987), Salanoubat and Belliard, *Gene*. 84:181-185 (1989), both of which are incorporated by reference
5 in their entirety), the promoter for the major tuber proteins including the 22 kd protein complexes and proteinase inhibitors (Hannapel, *Plant Physiol.* 101:703-704 (1993), herein incorporated by reference in its entirety), the promoter for the granule bound starch synthase gene (GBSS) (Visser *et al.*, *Plant Mol. Biol.* 17:691-699 (1991), herein incorporated by reference in its entirety) and other class I and II patatins promoters (Koster-Topfer *et al.*, *Mol Gen Genet.* 219:390-396 (1989); Mignery *et al.*, *Gene*. 62:27-44 (1988), both of which are herein
10 incorporated by reference in their entirety).

Other promoters can also be used to express a tocopherol synthesis pathway enzyme or fragment thereof in specific tissues, such as seeds or fruits. The promoter for β -conglycinin (Chen *et al.*, *Dev. Genet.* 10: 112-122 (1989), herein incorporated by reference in its entirety) or
15 other seed-specific promoters such as the napin and phaseolin promoters, can be used. The zeins are a group of storage proteins found in maize endosperm. Genomic clones for zein genes have been isolated (Pedersen *et al.*, *Cell* 29:1015-1026 (1982), herein incorporated by reference in its entirety) and the promoters from these clones, including the 15 kD, 16 kD, 19 kD, 22 kD, 27 kD and γ genes, could also be used. Other promoters known to function, for example, in maize
20 include the promoters for the following genes: *waxy*, *Brittle*, *Shrunken 2*, Branching enzymes I and II, starch synthases, debranching enzymes, oleosins, glutelins and sucrose synthases. A particularly preferred promoter for maize endosperm expression is the promoter for the glutelin

gene from rice, more particularly the Osgt-1 promoter (Zheng *et al.*, *Mol. Cell Biol.* 13:5829-5842 (1993), herein incorporated by reference in its entirety). Examples of promoters suitable for expression in wheat include those promoters for the ADPglucose pyrosynthase (ADPGPP) subunits, the granule bound and other starch synthase, the branching and debranching enzymes, the embryogenesis-abundant proteins, the gliadins and the glutenins. Examples of such promoters in rice include those promoters for the ADPGPP subunits, the granule bound and other starch synthase, the branching enzymes, the debranching enzymes, sucrose synthases and the glutelins. A particularly preferred promoter is the promoter for rice glutelin, Osgt-1. Examples of such promoters for barley include those for the ADPGPP subunits, the granule bound and other starch synthase, the branching enzymes, the debranching enzymes, sucrose synthases, the hordeins, the embryo globulins and the aleurone specific proteins.

Root specific promoters may also be used. An example of such a promoter is the promoter for the acid chitinase gene (Samac *et al.*, *Plant Mol. Biol.* 25:587-596 (1994), the entirety of which is herein incorporated by reference). Expression in root tissue could also be accomplished by utilizing the root specific subdomains of the CaMV35S promoter that have been identified (Lam *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:7890-7894 (1989), herein incorporated by reference in its entirety). Other root cell specific promoters include those reported by Conkling *et al.* (Conkling *et al.*, *Plant Physiol.* 93:1203-1211 (1990), the entirety of which is herein incorporated by reference).

Additional promoters that may be utilized are described, for example, in U.S. Patent Nos. 5,378,619; 5,391,725; 5,428,147; 5,447,858; 5,608,144; 5,608,144; 5,614,399; 5,633,441; 5,633,435; and 4,633,436, all of which are herein incorporated in their entirety. In addition, a

tissue specific enhancer may be used (Fromm *et al.*, *The Plant Cell* 1:977-984 (1989), the entirety of which is herein incorporated by reference).

Constructs or vectors may also include with the coding region of interest a nucleic acid sequence that acts, in whole or in part, to terminate transcription of that region. For example, such sequences have been isolated including the Tr7 3' sequence and the NOS 3' sequence (Ingelbrecht *et al.*, *The Plant Cell* 1:671-680 (1989), the entirety of which is herein incorporated by reference; Bevan *et al.*, *Nucleic Acids Res.* 11:369-385 (1983), the entirety of which is herein incorporated by reference), or the like.

A vector or construct may also include regulatory elements. Examples of such include the Adh intron 1 (Callis *et al.*, *Genes and Develop.* 1:1183-1200 (1987), the entirety of which is herein incorporated by reference), the sucrose synthase intron (Vasil *et al.*, *Plant Physiol.* 91:1575-1579 (1989), the entirety of which is herein incorporated by reference) and the TMV omega element (Gallie *et al.*, *The Plant Cell* 1:301-311 (1989), the entirety of which is herein incorporated by reference). These and other regulatory elements may be included when appropriate.

A vector or construct may also include a selectable marker. Selectable markers may also be used to select for plants or plant cells that contain the exogenous genetic material. Examples of such include, but are not limited to, a neo gene (Potrykus *et al.*, *Mol. Gen. Genet.* 199:183-188 (1985), the entirety of which is herein incorporated by reference) which codes for kanamycin resistance and can be selected for using kanamycin, G418, etc.; a bar gene which codes for bialaphos resistance; a mutant EPSP synthase gene (Hinchee *et al.*, *Bio/Technology* 6:915-922 (1988), the entirety of which is herein incorporated by reference) which encodes glyphosate resistance; a nitrilase gene which confers resistance to bromoxynil (Stalker *et al.*, *J. Biol. Chem.*

263:6310-6314 (1988), the entirety of which is herein incorporated by reference); a mutant acetolactate synthase gene (ALS) which confers imidazolinone or sulphonylurea resistance (European Patent Application 154,204 (Sept. 11, 1985), the entirety of which is herein incorporated by reference); and a methotrexate resistant DHFR gene (Thillet *et al.*, *J. Biol. Chem.* 263:12500-12508 (1988), the entirety of which is herein incorporated by reference).

A vector or construct may also include a transit peptide. Incorporation of a suitable chloroplast transit peptide may also be employed (European Patent Application Publication Number 0218571, the entirety of which is herein incorporated by reference). Translational enhancers may also be incorporated as part of the vector DNA. DNA constructs could contain one or more 5' non-translated leader sequences which may serve to enhance expression of the gene products from the resulting mRNA transcripts. Such sequences may be derived from the promoter selected to express the gene or can be specifically modified to increase translation of the mRNA. Such regions may also be obtained from viral RNAs, from suitable eukaryotic genes, or from a synthetic gene sequence. For a review of optimizing expression of transgenes, see Koziel *et al.*, *Plant Mol. Biol.* 32:393-405 (1996), the entirety of which is herein incorporated by reference.

A vector or construct may also include a screenable marker. Screenable markers may be used to monitor expression. Exemplary screenable markers include a β -glucuronidase or uidA gene (GUS) which encodes an enzyme for which various chromogenic substrates are known (Jefferson, *Plant Mol. Biol. Rep.* 5:387-405 (1987), the entirety of which is herein incorporated by reference; Jefferson *et al.*, *EMBO J.* 6:3901-3907 (1987), the entirety of which is herein incorporated by reference); an R-locus gene, which encodes a product that regulates the

production of anthocyanin pigments (red color) in plant tissues (Dellaporta *et al.*, Stadler Symposium 11:263-282 (1988), the entirety of which is herein incorporated by reference); a β -lactamase gene (Sutcliffe *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 75:3737-3741 (1978), the entirety of which is herein incorporated by reference), a gene which encodes an enzyme for which various chromogenic substrates are known (e.g., PADAC, a chromogenic cephalosporin); a luciferase gene (Ow *et al.*, *Science* 234:856-859 (1986), the entirety of which is herein incorporated by reference); a xylE gene (Zukowsky *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 80:1101-1105 (1983), the entirety of which is herein incorporated by reference) which encodes a catechol dioxygenase that can convert chromogenic catechols; an α -amylase gene (Ikata *et al.*, *Bio/Technol.* 8:241-242 (1990), the entirety of which is herein incorporated by reference); a tyrosinase gene (Katz *et al.*, *J. Gen. Microbiol.* 129:2703-2714 (1983), the entirety of which is herein incorporated by reference) which encodes an enzyme capable of oxidizing tyrosine to DOPA and dopaquinone which in turn condenses to melanin; an α -galactosidase, which will turn a chromogenic α -galactose substrate.

Included within the terms "selectable or screenable marker genes" are also genes which encode a secretable marker whose secretion can be detected as a means of identifying or selecting for transformed cells. Examples include markers which encode a secretable antigen that can be identified by antibody interaction, or even secretable enzymes which can be detected catalytically. Secretable proteins fall into a number of classes, including small, diffusible proteins which are detectable, (e.g., by ELISA), small active enzymes which are detectable in extracellular solution (e.g., α -amylase, β -lactamase, phosphinothricin transferase), or proteins which are inserted or trapped in the cell wall (such as proteins which include a leader sequence

such as that found in the expression unit of extension or tobacco PR-S). Other possible selectable and/or screenable marker genes will be apparent to those of skill in the art.

There are many methods for introducing transforming nucleic acid molecules into plant cells. Suitable methods are believed to include virtually any method by which nucleic acid molecules may be introduced into a cell, such as by *Agrobacterium* infection or direct delivery of nucleic acid molecules such as, for example, by PEG-mediated transformation, by electroporation or by acceleration of DNA coated particles, etc (Potrykus, *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 42:205-225 (1991), the entirety of which is herein incorporated by reference; Vasil, *Plant Mol. Biol.* 25:925-937 (1994), the entirety of which is herein incorporated by reference). For example, electroporation has been used to transform maize protoplasts (Fromm *et al.*, *Nature* 312:791-793 (1986), the entirety of which is herein incorporated by reference).

Other vector systems suitable for introducing transforming DNA into a host plant cell include but are not limited to binary artificial chromosome (BIBAC) vectors (Hamilton *et al.*, *Gene* 200:107-116 (1997), the entirety of which is herein incorporated by reference); and transfection with RNA viral vectors (Della-Cioppa *et al.*, *Ann. N.Y. Acad. Sci.* (1996), 792 (Engineering Plants for Commercial Products and Applications), 57-61, the entirety of which is herein incorporated by reference). Additional vector systems also include plant selectable YAC vectors such as those described in Mullen *et al.*, *Molecular Breeding* 4:449-457 (1988), the entirety of which is herein incorporated by reference).

Technology for introduction of DNA into cells is well known to those of skill in the art. Four general methods for delivering a gene into cells have been described: (1) chemical methods (Graham and van der Eb, *Virology* 54:536-539 (1973), the entirety of which is herein

incorporated by reference); (2) physical methods such as microinjection (Capecchi, *Cell* 22:479-488 (1980), the entirety of which is herein incorporated by reference), electroporation (Wong and Neumann, *Biochem. Biophys. Res. Commun.* 107:584-587 (1982); Fromm *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 82:5824-5828 (1985); U.S. Patent No. 5,384,253, all of which are herein
5 incorporated in their entirety); and the gene gun (Johnston and Tang, *Methods Cell Biol.* 43:353-365 (1994), the entirety of which is herein incorporated by reference); (3) viral vectors (Clapp, *Clin. Perinatol.* 20:155-168 (1993); Lu *et al.*, *J. Exp. Med.* 178:2089-2096 (1993); Eglitis and Anderson, *Biotechniques* 6:608-614 (1988), all of which are herein incorporated in their entirety); and (4) receptor-mediated mechanisms (Curiel *et al.*, *Hum. Gen. Ther.* 3:147-154 (1992), Wagner *et al.*, *Proc. Natl. Acad. Sci. (USA)* 89:6099-6103 (1992), both of which are
10 incorporated by reference in their entirety).

Acceleration methods that may be used include, for example, microprojectile bombardment and the like. One example of a method for delivering transforming nucleic acid molecules to plant cells is microprojectile bombardment. This method has been reviewed by Yang and Christou (eds.), *Particle Bombardment Technology for Gene Transfer*, Oxford Press, Oxford, England (1994), the entirety of which is herein incorporated by reference). Non-
15 biological particles (microprojectiles) that may be coated with nucleic acids and delivered into cells by a propelling force. Exemplary particles include those comprised of tungsten, gold, platinum and the like.

20 A particular advantage of microprojectile bombardment, in addition to it being an effective means of reproducibly transforming monocots, is that neither the isolation of protoplasts (Cristou *et al.*, *Plant Physiol.* 87:671-674 (1988), the entirety of which is herein incorporated by reference) nor the susceptibility of *Agrobacterium* infection are required. An

illustrative embodiment of a method for delivering DNA into maize cells by acceleration is a biolistics α -particle delivery system, which can be used to propel particles coated with DNA through a screen, such as a stainless steel or Nytex screen, onto a filter surface covered with corn cells cultured in suspension. Gordon-Kamm *et al.*, describes the basic procedure for coating tungsten particles with DNA (Gordon-Kamm *et al.*, *Plant Cell* 2:603-618 (1990), the entirety of which is herein incorporated by reference). The screen disperses the tungsten nucleic acid particles so that they are not delivered to the recipient cells in large aggregates. A particle delivery system suitable for use with the present invention is the helium acceleration PDS-1000/He gun is available from Bio-Rad Laboratories (Bio-Rad, Hercules, California)(Sanford *et al.*, *Technique* 3:3-16 (1991), the entirety of which is herein incorporated by reference).

For the bombardment, cells in suspension may be concentrated on filters. Filters containing the cells to be bombarded are positioned at an appropriate distance below the microprojectile stopping plate. If desired, one or more screens are also positioned between the gun and the cells to be bombarded.

Alternatively, immature embryos or other target cells may be arranged on solid culture medium. The cells to be bombarded are positioned at an appropriate distance below the microprojectile stopping plate. If desired, one or more screens are also positioned between the acceleration device and the cells to be bombarded. Through the use of techniques set forth herein one may obtain up to 1000 or more foci of cells transiently expressing a marker gene. The number of cells in a focus which express the exogenous gene product 48 hours post-bombardment often range from one to ten and average one to three.

In bombardment transformation, one may optimize the pre-bombardment culturing conditions and the bombardment parameters to yield the maximum numbers of stable transformants. Both the physical and biological parameters for bombardment are important in this technology. Physical factors are those that involve manipulating the DNA/microprojectile precipitate or those that affect the flight and velocity of either the macro- or microprojectiles. Biological factors include all steps involved in manipulation of cells before and immediately after bombardment, the osmotic adjustment of target cells to help alleviate the trauma associated with bombardment and also the nature of the transforming DNA, such as linearized DNA or intact supercoiled plasmids. It is believed that pre-bombardment manipulations are especially important for successful transformation of immature embryos.

In another alternative embodiment, plastids can be stably transformed. Methods disclosed for plastid transformation in higher plants include the particle gun delivery of DNA containing a selectable marker and targeting of the DNA to the plastid genome through homologous recombination (Svab *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 87:8526-8530 (1990); Svab and Maliga, *Proc. Natl. Acad. Sci. (U.S.A.)* 90:913-917 (1993); Staub and Maliga, *EMBO J.* 12:601-606 (1993); U.S. Patent Nos. 5, 451,513 and 5,545,818, all of which are herein incorporated by reference in their entirety).

Accordingly, it is contemplated that one may wish to adjust various aspects of the bombardment parameters in small scale studies to fully optimize the conditions. One may particularly wish to adjust physical parameters such as gap distance, flight distance, tissue distance and helium pressure. One may also minimize the trauma reduction factors by modifying conditions which influence the physiological state of the recipient cells and which may therefore influence transformation and integration efficiencies. For example, the osmotic state, tissue

hydration and the subculture stage or cell cycle of the recipient cells may be adjusted for optimum transformation. The execution of other routine adjustments will be known to those of skill in the art in light of the present disclosure.

Agrobacterium-mediated transfer is a widely applicable system for introducing genes into
5 plant cells because the DNA can be introduced into whole plant tissues, thereby bypassing the need for regeneration of an intact plant from a protoplast. The use of *Agrobacterium*-mediated plant integrating vectors to introduce DNA into plant cells is well known in the art. See, for example the methods described by Fraley *et al.*, *Bio/Technology* 3:629-635 (1985) and Rogers *et al.*, *Methods Enzymol.* 153:253-277 (1987), both of which are herein incorporated by reference in their entirety. Further, the integration of the Ti-DNA is a relatively precise process resulting in few rearrangements. The region of DNA to be transferred is defined by the border sequences and intervening DNA is usually inserted into the plant genome as described (Spielmann *et al.*, *Mol. Gen. Genet.* 205:34 (1986), the entirety of which is herein incorporated by reference).

Modern *Agrobacterium* transformation vectors are capable of replication in *E. coli* as well as *Agrobacterium*, allowing for convenient manipulations as described (Klee *et al.*, In: *Plant DNA Infectious Agents*, Hohn and Schell (eds.), Springer-Verlag, New York, pp. 179-203 (1985), the entirety of which is herein incorporated by reference. Moreover, technological advances in vectors for *Agrobacterium*-mediated gene transfer have improved the arrangement of genes and restriction sites in the vectors to facilitate construction of vectors capable of expressing various
20 polypeptide coding genes. The vectors described have convenient multi-linker regions flanked by a promoter and a polyadenylation site for direct expression of inserted polypeptide coding genes and are suitable for present purposes (Rogers *et al.*, *Methods Enzymol.* 153:253-277 (1987)). In addition, *Agrobacterium* containing both armed and disarmed Ti genes can be used

for the transformations. In those plant strains where *Agrobacterium*-mediated transformation is efficient, it is the method of choice because of the facile and defined nature of the gene transfer.

A transgenic plant formed using *Agrobacterium* transformation methods typically contains a single gene on one chromosome. Such transgenic plants can be referred to as being
5 heterozygous for the added gene. More preferred is a transgenic plant that is homozygous for the added structural gene; *i.e.*, a transgenic plant that contains two added genes, one gene at the same locus on each chromosome of a chromosome pair. A homozygous transgenic plant can be obtained by sexually mating (selfing) an independent segregant transgenic plant that contains a single added gene, germinating some of the seed produced and analyzing the resulting plants
10 produced for the gene of interest.

It is also to be understood that two different transgenic plants can also be mated to produce offspring that contain two independently segregating added, exogenous genes. Selfing
15 of appropriate progeny can produce plants that are homozygous for both added, exogenous genes that encode a polypeptide of interest. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation.

Transformation of plant protoplasts can be achieved using methods based on calcium phosphate precipitation, polyethylene glycol treatment, electroporation and combinations of
20 these treatments (*See, for example*, Potrykus *et al.*, *Mol. Gen. Genet.* 205:193-200 (1986); Lorz *et al.*, *Mol. Gen. Genet.* 199:178 (1985); Fromm *et al.*, *Nature* 319:791 (1986); Uchimiya *et al.*, *Mol. Gen. Genet.* 204:204 (1986); Marcotte *et al.*, *Nature* 335:454-457 (1988), all of which are herein incorporated by reference in their entirety).

Application of these systems to different plant strains depends upon the ability to regenerate that particular plant strain from protoplasts. Illustrative methods for the regeneration

of cereals from protoplasts are described (Fujimura *et al.*, *Plant Tissue Culture Letters* 2:74 (1985); Toriyama *et al.*, *Theor Appl. Genet.* 205:34 (1986); Yamada *et al.*, *Plant Cell Rep.* 4:85 (1986); Abdullah *et al.*, *Biotechnolog* 4:1087 (1986), all of which are herein incorporated by reference in their entirety).

5 To transform plant strains that cannot be successfully regenerated from protoplasts, other ways to introduce DNA into intact cells or tissues can be utilized. For example, regeneration of cereals from immature embryos or explants can be effected as described (Vasil, *Biotechnology* 6:397 (1988), the entirety of which is herein incorporated by reference). In addition, "particle gun" or high-velocity microprojectile technology can be utilized (Vasil *et al.*, *Bio/Technology* 10:667 (1992), the entirety of which is herein incorporated by reference).

Using the latter technology, DNA is carried through the cell wall and into the cytoplasm on the surface of small metal particles as described (Klein *et al.*, *Nature* 328:70 (1987); Klein *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 85:8502-8505 (1988); McCabe *et al.*, *Bio/Technology* 6:923 (1988), all of which are herein incorporated by reference in their entirety). The metal particles penetrate through several layers of cells and thus allow the transformation of cells within tissue explants.

Other methods of cell transformation can also be used and include but are not limited to introduction of DNA into plants by direct DNA transfer into pollen (Zhou *et al.*, *Methods Enzymol.* 101:433 (1983); Hess *et al.*, *Intern Rev. Cytol.* 107:367 (1987); Luo *et al.*, *Plant Mol Biol. Reporter* 6:165 (1988), all of which are herein incorporated by reference in their entirety),
20 by direct injection of DNA into reproductive organs of a plant (Pena *et al.*, *Nature* 325:274 (1987), the entirety of which is herein incorporated by reference), or by direct injection of DNA into the cells of immature embryos followed by the rehydration of desiccated embryos (Neuhaus

et al., *Theor. Appl. Genet.* 75:30 (1987), the entirety of which is herein incorporated by reference).

The regeneration, development and cultivation of plants from single plant protoplast transformants or from various transformed explants is well known in the art (Weissbach and Weissbach, In: *Methods for Plant Molecular Biology*, Academic Press, San Diego, CA, (1988), the entirety of which is herein incorporated by reference). This regeneration and growth process typically includes the steps of selection of transformed cells, culturing those individualized cells through the usual stages of embryonic development through the rooted plantlet stage. Transgenic embryos and seeds are similarly regenerated. The resulting transgenic rooted shoots are thereafter planted in an appropriate plant growth medium such as soil.

The development or regeneration of plants containing the foreign, exogenous gene that encodes a protein of interest is well known in the art. Preferably, the regenerated plants are self-pollinated to provide homozygous transgenic plants. Otherwise, pollen obtained from the regenerated plants is crossed to seed-grown plants of agronomically important lines. Conversely, pollen from plants of these important lines is used to pollinate regenerated plants. A transgenic plant of the present invention containing a desired polypeptide is cultivated using methods well known to one skilled in the art.

There are a variety of methods for the regeneration of plants from plant tissue. The particular method of regeneration will depend on the starting plant tissue and the particular plant species to be regenerated.

Methods for transforming dicots, primarily by use of *Agrobacterium tumefaciens* and obtaining transgenic plants have been published for cotton (U.S. Patent No. 5,004,863; U.S. Patent No. 5,159,135; U.S. Patent No. 5,518,908, all of which are herein incorporated by

reference in their entirety); soybean (U.S. Patent No. 5,569,834; U.S. Patent No. 5,416,011; McCabe *et al.*, *Biotechnology* 6:923 (1988); Christou *et al.*, *Plant Physiol.* 87:671-674 (1988); all of which are herein incorporated by reference in their entirety); *Brassica* (U.S. Patent No. 5,463,174, the entirety of which is herein incorporated by reference); peanut (Cheng *et al.*, *Plant*
5 *Cell Rep.* 15:653-657 (1996), McKently *et al.*, *Plant Cell Rep.* 14:699-703 (1995), all of which are herein incorporated by reference in their entirety); papaya; and pea (Grant *et al.*, *Plant Cell Rep.* 15:254-258 (1995), the entirety of which is herein incorporated by reference).

Transformation of monocotyledons using electroporation, particle bombardment and *Agrobacterium* have also been reported. Transformation and plant regeneration have been achieved in asparagus (Bytebier *et al.*, *Proc. Natl. Acad. Sci. (USA)* 84:5354 (1987), the entirety of which is herein incorporated by reference); barley (Wan and Lemaux, *Plant Physiol* 104:37 (1994), the entirety of which is herein incorporated by reference); maize (Rhodes *et al.*, *Science* 240:204 (1988); Gordon-Kamm *et al.*, *Plant Cell* 2:603-618 (1990); Fromm *et al.*, *Bio/Technology* 8:833 (1990); Koziel *et al.*, *Bio/Technology* 11:194 (1993); Armstrong *et al.*, *Crop Science* 35:550-557 (1995); all of which are herein incorporated by reference in their
10
15
entirety); oat (Somers *et al.*, *Bio/Technology* 10:1589 (1992), the entirety of which is herein incorporated by reference); orchard grass (Horn *et al.*, *Plant Cell Rep.* 7:469 (1988), the entirety of which is herein incorporated by reference); rice (Toriyama *et al.*, *Theor Appl. Genet.* 205:34 (1986); Part *et al.*, *Plant Mol. Biol.* 32:1135-1148 (1996); Abedinia *et al.*, *Aust. J. Plant Physiol.*
20 24:133-141 (1997); Zhang and Wu, *Theor. Appl. Genet.* 76:835 (1988); Zhang *et al.*, *Plant Cell Rep.* 7:379 (1988); Battraw and Hall, *Plant Sci.* 86:191-202 (1992); Christou *et al.*, *Bio/Technology* 9:957 (1991), all of which are herein incorporated by reference in their entirety); rye (De la Pena *et al.*, *Nature* 325:274 (1987), the entirety of which is herein incorporated by

reference); sugarcane (Bower and Birch, *Plant J.* 2:409 (1992), the entirety of which is herein incorporated by reference); tall fescue (Wang *et al.*, *Bio/Technology* 10:691 (1992), the entirety of which is herein incorporated by reference) and wheat (Vasil *et al.*, *Bio/Technology* 10:667 (1992), the entirety of which is herein incorporated by reference; U.S. Patent No. 5,631,152, the
5 entirety of which is herein incorporated by reference.)

Assays for gene expression based on the transient expression of cloned nucleic acid constructs have been developed by introducing the nucleic acid molecules into plant cells by polyethylene glycol treatment, electroporation, or particle bombardment (Marcotte *et al.*, *Nature* 335:454-457 (1988), the entirety of which is herein incorporated by reference; Marcotte *et al.*, *Plant Cell* 1:523-532 (1989), the entirety of which is herein incorporated by reference; McCarty *et al.*, *Cell* 66:895-905 (1991), the entirety of which is herein incorporated by reference; Hattori *et al.*, *Genes Dev.* 6:609-618 (1992), the entirety of which is herein incorporated by reference; Goff *et al.*, *EMBO J.* 9:2517-2522 (1990), the entirety of which is herein incorporated by reference). Transient expression systems may be used to functionally dissect gene constructs (see generally, Mailga *et al.*, *Methods in Plant Molecular Biology*, Cold Spring Harbor Press (1995)).

Any of the nucleic acid molecules of the present invention may be introduced into a plant cell in a permanent or transient manner in combination with other genetic elements such as vectors, promoters, enhancers etc. Further, any of the nucleic acid molecules of the present
20 invention may be introduced into a plant cell in a manner that allows for overexpression of the protein or fragment thereof encoded by the nucleic acid molecule.

Cosuppression is the reduction in expression levels, usually at the level of RNA, of a particular endogenous gene or gene family by the expression of a homologous sense construct

that is capable of transcribing mRNA of the same strandedness as the transcript of the endogenous gene (Napoli *et al.*, *Plant Cell* 2:279-289 (1990), the entirety of which is herein incorporated by reference; van der Krol *et al.*, *Plant Cell* 2:291-299 (1990), the entirety of which is herein incorporated by reference). Cosuppression may result from stable transformation with a single copy nucleic acid molecule that is homologous to a nucleic acid sequence found with the cell (Prolls and Meyer, *Plant J.* 2:465-475 (1992), the entirety of which is herein incorporated by reference) or with multiple copies of a nucleic acid molecule that is homologous to a nucleic acid sequence found with the cell (Mittlesten *et al.*, *Mol. Gen. Genet.* 244:325-330 (1994), the entirety of which is herein incorporated by reference). Genes, even though different, linked to homologous promoters may result in the cosuppression of the linked genes (Vaucheret, *C.R. Acad. Sci. III* 316:1471-1483 (1993), the entirety of which is herein incorporated by reference).

This technique has, for example, been applied to generate white flowers from red petunia and tomatoes that do not ripen on the vine. Up to 50% of petunia transformants that contained a sense copy of the glucoamylase (CHS) gene produced white flowers or floral sectors; this was as a result of the post-transcriptional loss of mRNA encoding CHS (Flavell, *Proc. Natl. Acad. Sci. (U.S.A.)* 91:3490-3496 (1994), the entirety of which is herein incorporated by reference); van Blokland *et al.*, *Plant J.* 6:861-877 (1994), the entirety of which is herein incorporated by reference). Cosuppression may require the coordinate transcription of the transgene and the endogenous gene and can be reset by a developmental control mechanism (Jorgensen, *Trends Biotechnol.* 8:340-344 (1990), the entirety of which is herein incorporated by reference; Meins and Kunz, In: *Gene Inactivation and Homologous Recombination in Plants*, Paszkowski (ed.), pp. 335-348, Kluwer Academic, Netherlands (1994), the entirety of which is herein incorporated by reference).

It is understood that one or more of the nucleic acids of the present invention may be introduced into a plant cell and transcribed using an appropriate promoter with such transcription resulting in the cosuppression of an endogenous tocopherol synthesis pathway enzyme.

Antisense approaches are a way of preventing or reducing gene function by targeting the genetic material (Mol *et al.*, *FEBS Lett.* 268:427-430 (1990), the entirety of which is herein incorporated by reference). The objective of the antisense approach is to use a sequence complementary to the target gene to block its expression and create a mutant cell line or organism in which the level of a single chosen protein is selectively reduced or abolished. Antisense techniques have several advantages over other 'reverse genetic' approaches. The site of inactivation and its developmental effect can be manipulated by the choice of promoter for antisense genes or by the timing of external application or microinjection. Antisense can manipulate its specificity by selecting either unique regions of the target gene or regions where it shares homology to other related genes (Hiatt *et al.*, In: *Genetic Engineering*, Setlow (ed.), Vol. 11, New York: Plenum 49-63 (1989), the entirety of which is herein incorporated by reference).

The principle of regulation by antisense RNA is that RNA that is complementary to the target mRNA is introduced into cells, resulting in specific RNA:RNA duplexes being formed by base pairing between the antisense substrate and the target mRNA (Green *et al.*, *Annu. Rev. Biochem.* 55:569-597 (1986), the entirety of which is herein incorporated by reference). Under one embodiment, the process involves the introduction and expression of an antisense gene sequence. Such a sequence is one in which part or all of the normal gene sequences are placed under a promoter in inverted orientation so that the 'wrong' or complementary strand is transcribed into a noncoding antisense RNA that hybridizes with the target mRNA and interferes with its expression (Takayama and Inouye, *Crit. Rev. Biochem. Mol. Biol.* 25:155-184 (1990),

the entirety of which is herein incorporated by reference). An antisense vector is constructed by standard procedures and introduced into cells by transformation, transfection, electroporation, microinjection, infection, etc. The type of transformation and choice of vector will determine whether expression is transient or stable. The promoter used for the antisense gene may
5 influence the level, timing, tissue, specificity, or inducibility of the antisense inhibition.

It is understood that the activity of a tocopherol synthesis pathway enzyme in a plant cell may be reduced or depressed by growing a transformed plant cell containing a nucleic acid molecule whose non-transcribed strand encodes a tocopherol synthesis pathway enzyme or fragment thereof.

Antibodies have been expressed in plants (Hiatt *et al.*, *Nature* 342:76-78 (1989), the
10 entirety of which is herein incorporated by reference; Conrad and Fielder, *Plant Mol. Biol.* 26:1023-1030 (1994), the entirety of which is herein incorporated by reference). Cytoplasmic expression of a scFv (single-chain Fv antibodies) has been reported to delay infection by artichoke mottled crinkle virus. Transgenic plants that express antibodies directed against
15 endogenous proteins may exhibit a physiological effect (Philips *et al.*, *EMBO J.* 16:4489-4496 (1997), the entirety of which is herein incorporated by reference; Marion-Poll, *Trends in Plant Science* 2:447-448 (1997), the entirety of which is herein incorporated by reference). For example, expressed anti-abscisic antibodies have been reported to result in a general perturbation of seed development (Philips *et al.*, *EMBO J.* 16: 4489-4496 (1997)).

20 Antibodies that are catalytic may also be expressed in plants (abzymes). The principle behind abzymes is that since antibodies may be raised against many molecules, this recognition ability can be directed toward generating antibodies that bind transition states to force a chemical reaction forward (Persidas, *Nature Biotechnology* 15:1313-1315 (1997), the entirety of which is

herein incorporated by reference; Baca *et al.*, *Ann. Rev. Biophys. Biomol. Struct.* 26:461-493 (1997), the entirety of which is herein incorporated by reference). The catalytic abilities of abzymes may be enhanced by site directed mutagenesis. Examples of abzymes are, for example, set forth in U.S. Patent No. 5,658,753; U.S. Patent No. 5,632,990; U.S. Patent No. 5,631,137; 5 U.S. Patent 5,602,015; U.S. Patent No. 5,559,538; U.S. Patent No. 5,576,174; U.S. Patent No. 5,500,358; U.S. Patent No. 5,318,897; U.S. Patent No. 5,298,409; U.S. Patent No. 5,258,289 and U.S. Patent No. 5,194,585, all of which are herein incorporated in their entirety.

It is understood that any of the antibodies of the present invention may be expressed in plants and that such expression can result in a physiological effect. It is also understood that any of the expressed antibodies may be catalytic.

(b) Fungal Constructs and Fungal Transformants

The present invention also relates to a fungal recombinant vector comprising exogenous genetic material. The present invention also relates to a fungal cell comprising a fungal recombinant vector. The present invention also relates to methods for obtaining a recombinant fungal host cell comprising introducing into a fungal host cell exogenous genetic material.

Exogenous genetic material may be transferred into a fungal cell. In a preferred embodiment the exogenous genetic material includes a nucleic acid molecule of the present invention having a sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof or fragments of either or other nucleic acid molecule of the present invention. The fungal recombinant vector may be any vector which can be conveniently 20 subjected to recombinant DNA procedures. The choice of a vector will typically depend on the compatibility of the vector with the fungal host cell into which the vector is to be introduced. The vector may be a linear or a closed circular plasmid. The vector system may be a single

vector or plasmid or two or more vectors or plasmids which together contain the total DNA to be introduced into the genome of the fungal host.

The fungal vector may be an autonomously replicating vector, *i.e.*, a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, *e.g.*, a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one which, when introduced into the fungal cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. For integration, the vector may rely on the nucleic acid sequence of the vector for stable integration of the vector into the genome by homologous or nonhomologous recombination. Alternatively, the vector may contain additional nucleic acid sequences for directing integration by homologous recombination into the genome of the fungal host. The additional nucleic acid sequences enable the vector to be integrated into the host cell genome at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, there should be preferably two nucleic acid sequences which individually contain a sufficient number of nucleic acids, preferably 400bp to 1500bp, more preferably 800bp to 1000bp, which are highly homologous with the corresponding target sequence to enhance the probability of homologous recombination. These nucleic acid sequences may be any sequence that is homologous with a target sequence in the genome of the fungal host cell and, furthermore, may be non-encoding or encoding sequences.

For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. Examples of origin of replications for use in a yeast host cell are the 2 micron origin of replication and the combination

of CEN3 and ARS 1. Any origin of replication may be used which is compatible with the fungal host cell of choice.

The fungal vectors of the present invention preferably contain one or more selectable markers which permit easy selection of transformed cells. A selectable marker is a gene the product of which provides, for example biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs and the like. The selectable marker may be selected from the group including, but not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferase), *bar* (phosphinothricin acetyltransferase), *hygB* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase) and *sC* (sulfate adenylyltransferase) and *trpC* (anthranilate synthase). Preferred for use in an *Aspergillus* cell are the *amdS* and *pyrG* markers of *Aspergillus nidulans* or *Aspergillus oryzae* and the *bar* marker of *Streptomyces hygroscopicus*. Furthermore, selection may be accomplished by co-transformation, e.g., as described in WO 91/17243, the entirety of which is herein incorporated by reference. A nucleic acid sequence of the present invention may be operably linked to a suitable promoter sequence. The promoter sequence is a nucleic acid sequence which is recognized by the fungal host cell for expression of the nucleic acid sequence. The promoter sequence contains transcription and translation control sequences which mediate the expression of the protein or fragment thereof.

A promoter may be any nucleic acid sequence which shows transcriptional activity in the fungal host cell of choice and may be obtained from genes encoding polypeptides either homologous or heterologous to the host cell. Examples of suitable promoters for directing the transcription of a nucleic acid construct of the invention in a filamentous fungal host are promoters obtained from the genes encoding *Aspergillus oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable

alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (glaA), *Rhizomucor miehei* lipase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Aspergillus nidulans* acetamidase and hybrids thereof. In a yeast host, a useful promoter is the *Saccharomyces cerevisiae* enolase (eno-1) promoter. Particularly preferred
5 promoters are the TAKA amylase, NA2-tpi (a hybrid of the promoters from the genes encoding *Aspergillus niger* neutral alpha -amylase and *Aspergillus oryzae* triose phosphate isomerase) and glaA promoters.

A protein or fragment thereof encoding nucleic acid molecule of the present invention may also be operably linked to a terminator sequence at its 3' terminus. The terminator sequence may be native to the nucleic acid sequence encoding the protein or fragment thereof or may be obtained from foreign sources. Any terminator which is functional in the fungal host cell of choice may be used in the present invention, but particularly preferred terminators are obtained from the genes encoding *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* alpha-glucosidase and
10 *Saccharomyces cerevisiae* enolase.
15

A protein or fragment thereof encoding nucleic acid molecule of the present invention may also be operably linked to a suitable leader sequence. A leader sequence is a nontranslated region of a mRNA which is important for translation by the fungal host. The leader sequence is operably linked to the 5' terminus of the nucleic acid sequence encoding the protein or fragment
20 thereof. The leader sequence may be native to the nucleic acid sequence encoding the protein or fragment thereof or may be obtained from foreign sources. Any leader sequence which is functional in the fungal host cell of choice may be used in the present invention, but particularly

preferred leaders are obtained from the genes encoding *Aspergillus oryzae* TAKA amylase and *Aspergillus oryzae* triose phosphate isomerase.

A polyadenylation sequence may also be operably linked to the 3' terminus of the nucleic acid sequence of the present invention. The polyadenylation sequence is a sequence which when
5 transcribed is recognized by the fungal host to add polyadenosine residues to transcribed mRNA. The polyadenylation sequence may be native to the nucleic acid sequence encoding the protein or fragment thereof or may be obtained from foreign sources. Any polyadenylation sequence which is functional in the fungal host of choice may be used in the present invention, but particularly preferred polyadenylation sequences are obtained from the genes encoding *Aspergillus oryzae*
10 TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase and *Aspergillus niger* alpha-glucosidase.

To avoid the necessity of disrupting the cell to obtain the protein or fragment thereof and to minimize the amount of possible degradation of the expressed protein or fragment thereof within the cell, it is preferred that expression of the protein or fragment thereof gives rise to a
15 product secreted outside the cell. To this end, a protein or fragment thereof of the present invention may be linked to a signal peptide linked to the amino terminus of the protein or fragment thereof. A signal peptide is an amino acid sequence which permits the secretion of the protein or fragment thereof from the fungal host into the culture medium. The signal peptide may be native to the protein or fragment thereof of the invention or may be obtained from
20 foreign sources. The 5' end of the coding sequence of the nucleic acid sequence of the present invention may inherently contain a signal peptide coding region naturally linked in translation reading frame with the segment of the coding region which encodes the secreted protein or fragment thereof. Alternatively, the 5' end of the coding sequence may contain a signal peptide

coding region which is foreign to that portion of the coding sequence which encodes the secreted protein or fragment thereof. The foreign signal peptide may be required where the coding sequence does not normally contain a signal peptide coding region. Alternatively, the foreign signal peptide may simply replace the natural signal peptide to obtain enhanced secretion of the
5 desired protein or fragment thereof. The foreign signal peptide coding region may be obtained from a glucoamylase or an amylase gene from an *Aspergillus* species, a lipase or proteinase gene from *Rhizomucor miehei*, the gene for the alpha-factor from *Saccharomyces cerevisiae*, or the calf preprochymosin gene. An effective signal peptide for fungal host cells is the *Aspergillus oryzae* TAKA amylase signal, *Aspergillus niger* neutral amylase signal, the *Rhizomucor miehei* aspartic proteinase signal, the *Humicola lanuginosus* cellulase signal, or the *Rhizomucor miehei* lipase signal. However, any signal peptide capable of permitting secretion of the protein or fragment thereof in a fungal host of choice may be used in the present invention.

A protein or fragment thereof encoding nucleic acid molecule of the present invention may also be linked to a propeptide coding region. A propeptide is an amino acid sequence found at the amino terminus of a protein or proenzyme. Cleavage of the propeptide from the
10 proprotein yields a mature biochemically active protein. The resulting polypeptide is known as a propolypeptide or proenzyme (or a zymogen in some cases). Propolypeptides are generally inactive and can be converted to mature active polypeptides by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide or proenzyme. The propeptide coding region may be
15 native to the protein or fragment thereof or may be obtained from foreign sources. The foreign propeptide coding region may be obtained from the *Saccharomyces cerevisiae* alpha-factor gene or *Myceliophthora thermophila* laccase gene (WO 95/33836, the entirety of which is herein incorporated by reference).
20

The procedures used to ligate the elements described above to construct the recombinant expression vector of the present invention are well known to one skilled in the art (see, for example, Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual*, 2nd ed., Cold Spring Harbor, N.Y., (1989)).

5 The present invention also relates to recombinant fungal host cells produced by the methods of the present invention which are advantageously used with the recombinant vector of the present invention. The cell is preferably transformed with a vector comprising a nucleic acid sequence of the invention followed by integration of the vector into the host chromosome. The choice of fungal host cells will to a large extent depend upon the gene encoding the protein or fragment thereof and its source. The fungal host cell may, for example, be a yeast cell or a filamentous fungal cell.

10 "Yeast" as used herein includes *Ascosporogenous* yeast (*Endomycetales*), *Basidiosporogenous* yeast and yeast belonging to the *Fungi Imperfecti* (*Blastomycetes*). The *Ascosporogenous* yeasts are divided into the families *Spermophthoraceae* and *Saccharomycetaceae*. The latter is comprised of four subfamilies, *Schizosaccharomycoideae* (for example, genus *Schizosaccharomyces*), *Nadsonioideae*, *Lipomycoideae* and *Saccharomycoideae* (for example, genera *Pichia*, *Kluyveromyces* and *Saccharomyces*). The *Basidiosporogenous* yeasts include the genera *Leucosporidim*, *Rhodospordium*, *Sporidiobolus*, *Filobasidium* and *Filobasidiella*. Yeast belonging to the *Fungi Imperfecti* are divided into two families, *Sporobolomycetaceae* (for example, genera *Sorobolomyces* and *Bullera*) and *Cryptococcaceae* (for example, genus *Candida*). Since the classification of yeast may change in the future, for the purposes of this invention, yeast shall be defined as described in *Biology and Activities of Yeast* (Skinner *et al.*, *Soc. App. Bacteriol. Symposium Series* No. 9, (1980), the entirety of which is

15
20

herein incorporated by reference). The biology of yeast and manipulation of yeast genetics are well known in the art (see, for example, *Biochemistry and Genetics of Yeast*, Bacil *et al.* (ed.), 2nd edition, 1987; *The Yeasts*, Rose and Harrison (eds.), 2nd ed., (1987); and *The Molecular Biology of the Yeast Saccharomyces*, Strathern *et al.* (eds.), (1981), all of which are herein
5 incorporated by reference in their entirety).

"Fungi" as used herein includes the phyla *Ascomycota*, *Basidiomycota*, *Chytridiomycota* and *Zygomycota* (as defined by Hawksworth *et al.*, In: Ainsworth and Bisby's *Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK; the entirety of which is herein incorporated by reference) as well as the *Oomycota* (as cited in Hawksworth *et al.*, In: Ainsworth and Bisby's *Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK) and all mitosporic fungi (Hawksworth *et al.*, In: Ainsworth and Bisby's *Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK). Representative groups of *Ascomycota* include, for example, *Neurospora*, *Eupenicillium* (= *Penicillium*), *Emericella* (= *Aspergillus*), *Eurotium* (= *Aspergillus*) and the true yeasts listed above. Examples of *Basidiomycota* include mushrooms, rusts and smuts.
10 Representative groups of *Chytridiomycota* include, for example, *Allomyces*, *Blastocladiella*, *Coelomomyces* and aquatic fungi. Representative groups of *Oomycota* include, for example, *Saprolegniomycetous* aquatic fungi (water molds) such as *Achlya*. Examples of mitosporic fungi include *Aspergillus*, *Penicillium*, *Candida* and *Alternaria*. Representative groups of *Zygomycota*
15 include, for example, *Rhizopus* and *Mucor*.
20

"Filamentous fungi" include all filamentous forms of the subdivision *Eumycota* and *Oomycota* (as defined by Hawksworth *et al.*, In: Ainsworth and Bisby's *Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK). The filamentous fungi

are characterized by a vegetative mycelium composed of chitin, cellulose, glucan, chitosan, mannan and other complex polysaccharides. Vegetative growth is by hyphal elongation and carbon catabolism is obligately aerobic. In contrast, vegetative growth by yeasts such as *Saccharomyces cerevisiae* is by budding of a unicellular thallus and carbon catabolism may be fermentative.

In one embodiment, the fungal host cell is a yeast cell. In a preferred embodiment, the yeast host cell is a cell of the species of *Candida*, *Kluyveromyces*, *Saccharomyces*, *Schizosaccharomyces*, *Pichia* and *Yarrowia*. In a preferred embodiment, the yeast host cell is a *Saccharomyces cerevisiae* cell, a *Saccharomyces carlsbergensis*, *Saccharomyces diastaticus* cell, a *Saccharomyces douglasii* cell, a *Saccharomyces kluyveri* cell, a *Saccharomyces norbensis* cell, or a *Saccharomyces oviformis* cell. In another preferred embodiment, the yeast host cell is a *Kluyveromyces lactis* cell. In another preferred embodiment, the yeast host cell is a *Yarrowia lipolytica* cell.

In another embodiment, the fungal host cell is a filamentous fungal cell. In a preferred embodiment, the filamentous fungal host cell is a cell of the species of, but not limited to, *Acremonium*, *Aspergillus*, *Fusarium*, *Humicola*, *Myceliophthora*, *Mucor*, *Neurospora*, *Penicillium*, *Thielavia*, *Tolypocladium* and *Trichoderma*. In a preferred embodiment, the filamentous fungal host cell is an *Aspergillus* cell. In another preferred embodiment, the filamentous fungal host cell is an *Acremonium* cell. In another preferred embodiment, the filamentous fungal host cell is a *Fusarium* cell. In another preferred embodiment, the filamentous fungal host cell is a *Humicola* cell. In another preferred embodiment, the filamentous fungal host cell is a *Myceliophthora* cell. In another even preferred embodiment, the filamentous fungal host cell is a *Mucor* cell. In another preferred embodiment, the filamentous

fungal host cell is a *Neurospora* cell. In another preferred embodiment, the filamentous fungal host cell is a *Penicillium* cell. In another preferred embodiment, the filamentous fungal host cell is a *Thielavia* cell. In another preferred embodiment, the filamentous fungal host cell is a *Tolypocladiun* cell. In another preferred embodiment, the filamentous fungal host cell is a *Trichoderma* cell. In a preferred embodiment, the filamentous fungal host cell is an *Aspergillus oryzae* cell, an *Aspergillus niger* cell, an *Aspergillus foetidus* cell, or an *Aspergillus japonicus* cell. In another preferred embodiment, the filamentous fungal host cell is a *Fusarium oxysporum* cell or a *Fusarium graminearum* cell. In another preferred embodiment, the filamentous fungal host cell is a *Humicola insolens* cell or a *Humicola lanuginosus* cell. In another preferred embodiment, the filamentous fungal host cell is a *Myceliophthora thermophila* cell. In a most preferred embodiment, the filamentous fungal host cell is a *Mucor miehei* cell. In a most preferred embodiment, the filamentous fungal host cell is a *Neurospora crassa* cell. In a most preferred embodiment, the filamentous fungal host cell is a *Penicillium purpurogenum* cell. In another most preferred embodiment, the filamentous fungal host cell is a *Thielavia terrestris* cell. In another most preferred embodiment, the *Trichoderma* cell is a *Trichoderma reesei* cell, a *Trichoderma viride* cell, a *Trichoderma longibrachiatum* cell, a *Trichoderma harzianum* cell, or a *Trichoderma koningii* cell. In a preferred embodiment, the fungal host cell is selected from an *A. nidulans* cell, an *A. niger* cell, an *A. oryzae* cell and an *A. sojæ* cell. In a further preferred embodiment, the fungal host cell is an *A. nidulans* cell.

The recombinant fungal host cells of the present invention may further comprise one or more sequences which encode one or more factors that are advantageous in the expression of the protein or fragment thereof, for example, an activator (e.g., a trans-acting factor), a chaperone and a processing protease. The nucleic acids encoding one or more of these factors are

preferably not operably linked to the nucleic acid encoding the protein or fragment thereof. An activator is a protein which activates transcription of a nucleic acid sequence encoding a polypeptide (Kudla *et al.*, *EMBO* 9:1355-1364(1990); Jarai and Buxton, *Current Genetics* 26:2238-244(1994); Verdier, *Yeast* 6:271-297(1990), all of which are herein incorporated by reference in their entirety). The nucleic acid sequence encoding an activator may be obtained from the genes encoding *Saccharomyces cerevisiae* heme activator protein 1 (hap1), *Saccharomyces cerevisiae* galactose metabolizing protein 4 (gal4) and *Aspergillus nidulans* ammonia regulation protein (areA). For further examples, see Verdier, *Yeast* 6:271-297 (1990); MacKenzie *et al.*, *Journal of Gen. Microbiol.* 139:2295-2307 (1993), both of which are herein incorporated by reference in their entirety). A chaperone is a protein which assists another protein in folding properly (Hartl *et al.*, *TIBS* 19:20-25 (1994); Bergeron *et al.*, *TIBS* 19:124-128 (1994); Demolder *et al.*, *J. Biotechnology* 32:179-189 (1994); Craig, *Science* 260:1902-1903(1993); Gething and Sambrook, *Nature* 355:33-45 (1992); Puig and Gilbert, *J Biol. Chem.* 269:7764-7771 (1994); Wang and Tsou, *FASEB Journal* 7:1515-11157 (1993); Robinson *et al.*, *Bio/Technology* 1:381-384 (1994), all of which are herein incorporated by reference in their entirety). The nucleic acid sequence encoding a chaperone may be obtained from the genes encoding *Aspergillus oryzae* protein disulphide isomerase, *Saccharomyces cerevisiae* calnexin, *Saccharomyces cerevisiae* BiP/GRP78 and *Saccharomyces cerevisiae* Hsp70. For further examples, see Gething and Sambrook, *Nature* 355:33-45 (1992); Hartl *et al.*, *TIBS* 19:20-25 (1994). A processing protease is a protease that cleaves a propeptide to generate a mature biochemically active polypeptide (Enderlin and Ogrydziak, *Yeast* 10:67-79 (1994); Fuller *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:1434-1438 (1989); Julius *et al.*, *Cell* 37:1075-1089 (1984); Julius *et al.*, *Cell* 32:839-852 (1983), all of which are incorporated by reference in their entirety).

The nucleic acid sequence encoding a processing protease may be obtained from the genes encoding *Aspergillus niger* Kex2, *Saccharomyces cerevisiae* dipeptidylaminopeptidase, *Saccharomyces cerevisiae* Kex2 and *Yarrowia lipolytica* dibasic processing endoprotease (xpr6). Any factor that is functional in the fungal host cell of choice may be used in the present invention.

Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts and regeneration of the cell wall in a manner known per se. Suitable procedures for transformation of *Aspergillus* host cells are described in EP 238 023 and Yelton *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 81:1470-1474 (1984), both of which are herein incorporated by reference in their entirety. A suitable method of transforming *Fusarium* species is described by Malardier *et al.*, *Gene* 78:147-156 (1989), the entirety of which is herein incorporated by reference. Yeast may be transformed using the procedures described by Becker and Guarente, In: Abelson and Simon, (eds.), *Guide to Yeast Genetics and Molecular Biology, Methods Enzymol.* Volume 194, pp 182-187, Academic Press, Inc., New York; Ito *et al.*, *J. Bacteriology* 153:163 (1983); Hinnen *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 75:1920 (1978), all of which are herein incorporated by reference in their entirety.

The present invention also relates to methods of producing the protein or fragment thereof comprising culturing the recombinant fungal host cells under conditions conducive for expression of the protein or fragment thereof. The fungal cells of the present invention are cultivated in a nutrient medium suitable for production of the protein or fragment thereof using methods known in the art. For example, the cell may be cultivated by shake flask cultivation, small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under

conditions allowing the protein or fragment thereof to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art (*see, e.g., Bennett and LaSure (eds.), More Gene Manipulations in Fungi*, Academic Press, CA, (1991), the entirety of which is herein
5 incorporated by reference). Suitable media are available from commercial suppliers or may be prepared according to published compositions (*e.g., in catalogues of the American Type Culture Collection, Manassas, VA*). If the protein or fragment thereof is secreted into the nutrient medium, a protein or fragment thereof can be recovered directly from the medium. If the protein or fragment thereof is not secreted, it is recovered from cell lysates.

The expressed protein or fragment thereof may be detected using methods known in the art that are specific for the particular protein or fragment. These detection methods may include the use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example, if the protein or fragment thereof has enzymatic activity, an enzyme assay may be used. Alternatively, if polyclonal or monoclonal antibodies specific to the protein or fragment thereof are available, immunoassays may be employed using the antibodies to the protein or fragment thereof. The techniques of enzyme assay and immunoassay are well known to those skilled in the art.

The resulting protein or fragment thereof may be recovered by methods known in the arts. For example, the protein or fragment thereof may be recovered from the nutrient medium
20 by conventional procedures including, but not limited to, centrifugation, filtration, extraction, spray-drying, evaporation, or precipitation. The recovered protein or fragment thereof may then be further purified by a variety of chromatographic procedures, *e.g., ion exchange chromatography, gel filtration chromatography, affinity chromatography, or the like.*

(c) Mammalian Constructs and Transformed Mammalian Cells

The present invention also relates to methods for obtaining a recombinant mammalian host cell, comprising introducing into a mammalian host cell exogenous genetic material. The present invention also relates to a mammalian cell comprising a mammalian recombinant vector.

5 The present invention also relates to methods for obtaining a recombinant mammalian host cell, comprising introducing into a mammalian cell exogenous genetic material. In a preferred embodiment the exogenous genetic material includes a nucleic acid molecule of the present invention having a sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof or fragments of either or other nucleic acid molecule of the present invention.

10 Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC, Manassas, VA), such as HeLa cells, Chinese hamster ovary (CHO) cells, baby hamster kidney (BHK) cells and a number of other cell lines. Suitable promoters for mammalian cells are also known in the art and include viral promoters such as that from Simian Virus 40 (SV40) (Fiers *et al.*, *Nature* 273:113 (1978), the entirety of which is herein incorporated by reference), Rous sarcoma virus (RSV), adenovirus (ADV) and bovine papilloma virus (BPV). Mammalian cells may also require terminator sequences and poly-A addition sequences. Enhancer sequences which increase expression may also be included and sequences which promote amplification of
20 the gene may also be desirable (for example methotrexate resistance genes).

Vectors suitable for replication in mammalian cells may include viral replicons, or sequences which insure integration of the appropriate sequences encoding HCV epitopes into the host genome. For example, another vector used to express foreign DNA is vaccinia virus. In this

case, for example, a nucleic acid molecule encoding a protein or fragment thereof is inserted into the vaccinia genome. Techniques for the insertion of foreign DNA into the vaccinia virus genome are known in the art and may utilize, for example, homologous recombination. Such heterologous DNA is generally inserted into a gene which is non-essential to the virus, for example, the thymidine kinase gene (tk), which also provides a selectable marker. Plasmid vectors that greatly facilitate the construction of recombinant viruses have been described (see, for example, Mackett *et al*, *J Virol.* 49:857 (1984); Chakrabarti *et al.*, *Mol. Cell. Biol.* 5:3403 (1985); Moss, In: *Gene Transfer Vectors For Mammalian Cells* (Miller and Calos, eds., Cold Spring Harbor Laboratory, N.Y., p. 10, (1987); all of which are herein incorporated by reference in their entirety). Expression of the HCV polypeptide then occurs in cells or animals which are infected with the live recombinant vaccinia virus.

The sequence to be integrated into the mammalian sequence may be introduced into the primary host by any convenient means, which includes calcium precipitated DNA, spheroplast fusion, transformation, electroporation, biolistics, lipofection, microinjection, or other convenient means. Where an amplifiable gene is being employed, the amplifiable gene may serve as the selection marker for selecting hosts into which the amplifiable gene has been introduced. Alternatively, one may include with the amplifiable gene another marker, such as a drug resistance marker, e.g. neomycin resistance (G418 in mammalian cells), hygromycin in resistance etc., or an auxotrophy marker (HIS3, TRP1, LEU2, URA3, ADE2, LYS2, etc.) for use in yeast cells.

Depending upon the nature of the modification and associated targeting construct, various techniques may be employed for identifying targeted integration. Conveniently, the DNA may be digested with one or more restriction enzymes and the fragments probed with an appropriate

DNA fragment which will identify the properly sized restriction fragment associated with integration.

One may use different promoter sequences, enhancer sequences, or other sequence which will allow for enhanced levels of expression in the expression host. Thus, one may combine an enhancer from one source, a promoter region from another source, a 5'- noncoding region upstream from the initiation methionine from the same or different source as the other sequences and the like. One may provide for an intron in the non-coding region with appropriate splice sites or for an alternative 3'- untranslated sequence or polyadenylation site. Depending upon the particular purpose of the modification, any of these sequences may be introduced, as desired.

Where selection is intended, the sequence to be integrated will have with it a marker gene, which allows for selection. The marker gene may conveniently be downstream from the target gene and may include resistance to a cytotoxic agent, e.g. antibiotics, heavy metals, or the like, resistance or susceptibility to HAT, gancyclovir, etc., complementation to an auxotrophic host, particularly by using an auxotrophic yeast as the host for the subject manipulations, or the like. The marker gene may also be on a separate DNA molecule, particularly with primary mammalian cells. Alternatively, one may screen the various transformants, due to the high efficiency of recombination in yeast, by using hybridization analysis, PCR, sequencing, or the like.

For homologous recombination, constructs can be prepared where the amplifiable gene will be flanked, normally on both sides with DNA homologous with the DNA of the target region. Depending upon the nature of the integrating DNA and the purpose of the integration, the homologous DNA will generally be within 100kb, usually 50kb, preferably about 25kb, of the transcribed region of the target gene, more preferably within 2kb of the target gene. Where

modeling of the gene is intended, homology will usually be present proximal to the site of the mutation. The homologous DNA may include the 5'-upstream region outside of the transcriptional regulatory region or comprising any enhancer sequences, transcriptional initiation sequences, adjacent sequences, or the like. The homologous region may include a portion of the coding region, where the coding region may be comprised only of an open reading frame or combination of exons and introns. The homologous region may comprise all or a portion of an intron, where all or a portion of one or more exons may also be present. Alternatively, the homologous region may comprise the 3'-region, so as to comprise all or a portion of the transcriptional termination region, or the region 3' of this region. The homologous regions may extend over all or a portion of the target gene or be outside the target gene comprising all or a portion of the transcriptional regulatory regions and/or the structural gene.

The integrating constructs may be prepared in accordance with conventional ways, where sequences may be synthesized, isolated from natural sources, manipulated, cloned, ligated, subjected to in vitro mutagenesis, primer repair, or the like. At various stages, the joined sequences may be cloned and analyzed by restriction analysis, sequencing, or the like. Usually during the preparation of a construct where various fragments are joined, the fragments, intermediate constructs and constructs will be carried on a cloning vector comprising a replication system functional in a prokaryotic host, e.g., *E. coli* and a marker for selection, e.g., biocide resistance, complementation to an auxotrophic host, etc. Other functional sequences may also be present, such as polylinkers, for ease of introduction and excision of the construct or portions thereof, or the like. A large number of cloning vectors are available such as pBR322, the pUC series, etc. These constructs may then be used for integration into the primary mammalian host.

In the case of the primary mammalian host, a replicating vector may be used. Usually, such vector will have a viral replication system, such as SV40, bovine papilloma virus, adenovirus, or the like. The linear DNA sequence vector may also have a selectable marker for identifying transfected cells. Selectable markers include the neo gene, allowing for selection
5 with G418, the herpes tk gene for selection with HAT medium, the gpt gene with mycophenolic acid, complementation of an auxotrophic host, etc.

The vector may or may not be capable of stable maintenance in the host. Where the vector is capable of stable maintenance, the cells will be screened for homologous integration of the vector into the genome of the host, where various techniques for curing the cells may be employed. Where the vector is not capable of stable maintenance, for example, where a temperature sensitive replication system is employed, one may change the temperature from the permissive temperature to the non-permissive temperature, so that the cells may be cured of the vector. In this case, only those cells having integration of the construct comprising the amplifiable gene and, when present, the selectable marker, will be able to survive selection.

Where a selectable marker is present, one may select for the presence of the targeting construct by means of the selectable marker. Where the selectable marker is not present, one may select for the presence of the construct by the amplifiable gene. For the neo gene or the herpes tk gene, one could employ a medium for growth of the transformants of about 0.1-1 mg/ml of G418 or may use HAT medium, respectively. Where DHFR is the amplifiable gene,
20 the selective medium may include from about 0.01-0.5 μ M of methotrexate or be deficient in glycine-hypoxanthine-thymidine and have dialysed serum (GHT media).

The DNA can be introduced into the expression host by a variety of techniques that include calcium phosphate/DNA co-precipitates, microinjection of DNA into the nucleus,

electroporation, yeast protoplast fusion with intact cells, transfection, polycations, e.g., polybrene, polyornithine, etc., or the like. The DNA may be single or double stranded DNA, linear or circular. The various techniques for transforming mammalian cells are well known (see Keown *et al.*, *Methods Enzymol.* (1989); Keown *et al.*, *Methods Enzymol.* 185:527-537 (1990); Mansour *et al.*, *Nature* 336:348-352, (1988); all of which are herein incorporated by reference in their entirety).

(d) Insect Constructs and Transformed Insect Cells

The present invention also relates to an insect recombinant vectors comprising exogenous genetic material. The present invention also relates to an insect cell comprising an insect recombinant vector. The present invention also relates to methods for obtaining a recombinant insect host cell, comprising introducing into an insect cell exogenous genetic material. In a preferred embodiment the exogenous genetic material includes a nucleic acid molecule of the present invention having a sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof or fragments of either or other nucleic acid molecule of the present invention.

The insect recombinant vector may be any vector which can be conveniently subjected to recombinant DNA procedures and can bring about the expression of the nucleic acid sequence. The choice of a vector will typically depend on the compatibility of the vector with the insect host cell into which the vector is to be introduced. The vector may be a linear or a closed circular plasmid. The vector system may be a single vector or plasmid or two or more vectors or plasmids which together contain the total DNA to be introduced into the genome of the insect host. In addition, the insect vector may be an expression vector. Nucleic acid molecules can be suitably inserted into a replication vector for expression in the insect cell under a suitable

promoter for insect cells. Many vectors are available for this purpose and selection of the appropriate vector will depend mainly on the size of the nucleic acid molecule to be inserted into the vector and the particular host cell to be transformed with the vector. Each vector contains various components depending on its function (amplification of DNA or expression of DNA) and the particular host cell with which it is compatible. The vector components for insect cell transformation generally include, but are not limited to, one or more of the following: a signal sequence, origin of replication, one or more marker genes and an inducible promoter.

The insect vector may be an autonomously replicating vector, *i.e.*, a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, *e.g.*, a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one which, when introduced into the insect cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. For integration, the vector may rely on the nucleic acid sequence of the vector for stable integration of the vector into the genome by homologous or nonhomologous recombination. Alternatively, the vector may contain additional nucleic acid sequences for directing integration by homologous recombination into the genome of the insect host. The additional nucleic acid sequences enable the vector to be integrated into the host cell genome at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, there should be preferably two nucleic acid sequences which individually contain a sufficient number of nucleic acids, preferably 400bp to 1500bp, more preferably 800bp to 1000bp, which are highly homologous with the corresponding target sequence to enhance the probability of homologous recombination. These nucleic acid sequences may be any sequence that is homologous with a

target sequence in the genome of the insect host cell and, furthermore, may be non-encoding or encoding sequences.

5 Baculovirus expression vectors (BEVs) have become important tools for the expression of foreign genes, both for basic research and for the production of proteins with direct clinical applications in human and veterinary medicine (Doerfler, *Curr. Top. Microbiol. Immunol.* 131:51-68 (1968); Luckow and Summers, *Bio/Technology* 6:47-55 (1988a); Miller, *Annual Review of Microbiol.* 42:177-199 (1988); Summers, *Curr. Comm. Molecular Biology*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1988); all of which are herein incorporated by reference in their entirety). BEVs are recombinant insect viruses in which the coding sequence for a chosen foreign gene has been inserted behind a baculovirus promoter in place of the viral gene, e.g., polyhedrin (Smith and Summers, U.S. Pat. No., 4,745,051, the entirety of which is incorporated herein by reference).

10 The use of baculovirus vectors relies upon the host cells being derived from *Lepidopteran* insects such as *Spodoptera frugiperda* or *Trichoplusia ni*. The preferred *Spodoptera frugiperda* cell line is the cell line Sf9. The *Spodoptera frugiperda* Sf9 cell line was obtained from American Type Culture Collection (Manassas, VA.) and is assigned accession number ATCC CRL 1711 (Summers and Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Ag. Exper. Station Bulletin No. 1555 (1988), the entirety of which is herein incorporated by reference). Other insect cell systems, such as the silkworm *B. mori* may
20 also be used.

The proteins expressed by the BEVs are, therefore, synthesized, modified and transported in host cells derived from *Lepidopteran* insects. Most of the genes that have been inserted and produced in the baculovirus expression vector system have been derived from vertebrate species.

Other baculovirus genes in addition to the polyhedrin promoter may be employed to advantage in a baculovirus expression system. These include immediate-early (α), delayed-early (β), late (γ), or very late (δ), according to the phase of the viral infection during which they are expressed. The expression of these genes occurs sequentially, probably as the result of a "cascade" mechanism of transcriptional regulation. (Guarino and Summers, *J. Virol.* 57:563-571 (1986); Guarino and Summers, *J. Virol.* 61:2091-2099 (1987); Guarino and Summers, *Virol.* 162:444-451 (1988); all of which are herein incorporated by reference in their entirety).

Insect recombinant vectors are useful as intermediates for the infection or transformation of insect cell systems. For example, an insect recombinant vector containing a nucleic acid molecule encoding a baculovirus transcriptional promoter followed downstream by an insect signal DNA sequence is capable of directing the secretion of the desired biologically active protein from the insect cell. The vector may utilize a baculovirus transcriptional promoter region derived from any of the over 500 baculoviruses generally infecting insects, such as for example the Orders *Lepidoptera*, *Diptera*, *Orthoptera*, *Coleoptera* and *Hymenoptera*, including for example but not limited to the viral DNAs of *Autographa californica* MNPV, *Bombyx mori* NPV, *Trichoplusia ni* MNPV, *Rachiplusia ou* MNPV or *Galleria mellonella* MNPV, wherein said baculovirus transcriptional promoter is a baculovirus immediate-early gene IEl or IEN promoter; an immediate-early gene in combination with a baculovirus delayed-early gene promoter region selected from the group consisting of 39K and a *HindIII-k* fragment delayed-early gene; or a baculovirus late gene promoter. The immediate-early or delayed-early promoters can be enhanced with transcriptional enhancer elements. The insect signal DNA sequence may code for a signal peptide of a *Lepidopteran* adipokinetic hormone precursor or a signal peptide of the *Manduca sexta* adipokinetic hormone precursor (Summers, U.S. Patent No. 5,155,037; the

entirety of which is herein incorporated by reference). Other insect signal DNA sequences include a signal peptide of the *Orthoptera Schistocerca gregaria* locust adipokinetic hormone precursor and the *Drosophila melanogaster* cuticle genes CP1, CP2, CP3 or CP4 or for an insect signal peptide having substantially a similar chemical composition and function (Summers, U.S. Patent No. 5,155,037).

Insect cells are distinctly different from animal cells. Insects have a unique life cycle and have distinct cellular properties such as the lack of intracellular plasminogen activators in which are present in vertebrate cells. Another difference is the high expression levels of protein products ranging from 1 to greater than 500 mg/liter and the ease at which cDNA can be cloned into cells (Frasier, *In Vitro Cell. Dev. Biol.* 25:225 (1989); Summers and Smith, In: *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Ag. Exper. Station Bulletin No. 1555 (1988), both of which are incorporated by reference in their entirety).

Recombinant protein expression in insect cells is achieved by viral infection or stable transformation. For viral infection, the desired gene is cloned into baculovirus at the site of the wild-type polyhedron gene (Webb and Summers, *Technique* 2:173 (1990); Bishop and Posse, *Adv. Gene Technol.* 1:55 (1990); both of which are incorporated by reference in their entirety). The polyhedron gene is a component of a protein coat in occlusions which encapsulate virus particles. Deletion or insertion in the polyhedron gene results the failure to form occlusion bodies. Occlusion negative viruses are morphologically different from occlusion positive viruses and enable one skilled in the art to identify and purify recombinant viruses.

The vectors of present invention preferably contain one or more selectable markers which permit easy selection of transformed cells. A selectable marker is a gene the product of which provides, for example biocide or viral resistance, resistance to heavy metals, prototrophy to

auxotrophs and the like. Selection may be accomplished by co-transformation, *e.g.*, as described in WO 91/17243, a nucleic acid sequence of the present invention may be operably linked to a suitable promoter sequence. The promoter sequence is a nucleic acid sequence which is recognized by the insect host cell for expression of the nucleic acid sequence. The promoter
5 sequence contains transcription and translation control sequences which mediate the expression of the protein or fragment thereof. The promoter may be any nucleic acid sequence which shows transcriptional activity in the insect host cell of choice and may be obtained from genes encoding polypeptides either homologous or heterologous to the host cell.

For example, a nucleic acid molecule encoding a protein or fragment thereof may also be operably linked to a suitable leader sequence. A leader sequence is a nontranslated region of a mRNA which is important for translation by the fungal host. The leader sequence is operably linked to the 5' terminus of the nucleic acid sequence encoding the protein or fragment thereof. The leader sequence may be native to the nucleic acid sequence encoding the protein or fragment thereof or may be obtained from foreign sources. Any leader sequence which is functional in the insect host cell of choice may be used in the present invention.

A polyadenylation sequence may also be operably linked to the 3' terminus of the nucleic acid sequence of the present invention. The polyadenylation sequence is a sequence which when transcribed is recognized by the insect host to add polyadenosine residues to transcribed mRNA. The polyadenylation sequence may be native to the nucleic acid sequence encoding the protein or
20 fragment thereof or may be obtained from foreign sources. Any polyadenylation sequence which is functional in the fungal host of choice may be used in the present invention.

To avoid the necessity of disrupting the cell to obtain the protein or fragment thereof and to minimize the amount of possible degradation of the expressed polypeptide within the cell, it is

preferred that expression of the polypeptide gene gives rise to a product secreted outside the cell. To this end, the protein or fragment thereof of the present invention may be linked to a signal peptide linked to the amino terminus of the protein or fragment thereof. A signal peptide is an amino acid sequence which permits the secretion of the protein or fragment thereof from the insect host into the culture medium. The signal peptide may be native to the protein or fragment thereof of the invention or may be obtained from foreign sources. The 5' end of the coding sequence of the nucleic acid sequence of the present invention may inherently contain a signal peptide coding region naturally linked in translation reading frame with the segment of the coding region which encodes the secreted protein or fragment thereof.

At present, a mode of achieving secretion of a foreign gene product in insect cells is by way of the foreign gene's native signal peptide. Because the foreign genes are usually from non-insect organisms, their signal sequences may be poorly recognized by insect cells and hence, levels of expression may be suboptimal. However, the efficiency of expression of foreign gene products seems to depend primarily on the characteristics of the foreign protein. On average, nuclear localized or non-structural proteins are most highly expressed, secreted proteins are intermediate and integral membrane proteins are the least expressed. One factor generally affecting the efficiency of the production of foreign gene products in a heterologous host system is the presence of native signal sequences (also termed presequences, targeting signals, or leader sequences) associated with the foreign gene. The signal sequence is generally coded by a DNA sequence immediately following (5' to 3') the translation start site of the desired foreign gene.

The expression dependence on the type of signal sequence associated with a gene product can be represented by the following example: If a foreign gene is inserted at a site downstream from the translational start site of the baculovirus polyhedrin gene so as to produce a fusion

protein (containing the N-terminus of the polyhedrin structural gene), the fused gene is highly expressed. But less expression is achieved when a foreign gene is inserted in a baculovirus expression vector immediately following the transcriptional start site and totally replacing the polyhedrin structural gene.

5 Insertions into the region -50 to -1 significantly alter (reduce) steady state transcription which, in turn, reduces translation of the foreign gene product. Use of the pVL941 vector optimizes transcription of foreign genes to the level of the polyhedrin gene transcription. Even though the transcription of a foreign gene may be optimal, optimal translation may vary because of several factors involving processing: signal peptide recognition, mRNA and ribosome binding, glycosylation, disulfide bond formation, sugar processing, oligomerization, for example.

10 The properties of the insect signal peptide are expected to be more optimal for the efficiency of the translation process in insect cells than those from vertebrate proteins. This phenomenon can generally be explained by the fact that proteins secreted from cells are synthesized as precursor molecules containing hydrophobic N-terminal signal peptides. The signal peptides direct transport of the select protein to its target membrane and are then cleaved by a peptidase on the membrane, such as the endoplasmic reticulum, when the protein passes through it.

15 Another exemplary insect signal sequence is the sequence encoding for *Drosophila* cuticle proteins such as CP1, CP2, CP3 or CP4 (Summers, U.S. Patent No. 5,278,050; the
20 entirety of which is herein incorporated by reference). Most of a 9kb region of the *Drosophila* genome containing genes for the cuticle proteins has been sequenced. Four of the five cuticle genes contains a signal peptide coding sequence interrupted by a short intervening sequence (about 60 base pairs) at a conserved site. Conserved sequences occur in the 5' mRNA

untranslated region, in the adjacent 35 base pairs of upstream flanking sequence and at -200 base pairs from the mRNA start position in each of the cuticle genes.

Standard methods of insect cell culture, cotransfection and preparation of plasmids are set forth in Summers and Smith (Summers and Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agricultural Experiment Station Bulletin No. 1555, Texas A&M University (1987)). Procedures for the cultivation of viruses and cells are described in Volkman and Summers, *J. Virol* 19:820-832 (1975) and Volkman *et al.*, *J. Virol* 19:820-832 (1976); both of which are herein incorporated by reference in their entirety.

(e) Bacterial Constructs and Transformed Bacterial Cells

The present invention also relates to a bacterial recombinant vector comprising exogenous genetic material. The present invention also relates to a bacteria cell comprising a bacterial recombinant vector. The present invention also relates to methods for obtaining a recombinant bacteria host cell, comprising introducing into a bacterial host cell exogenous genetic material. . In a preferred embodiment the exogenous genetic material includes a nucleic acid molecule of the present invention having a sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 627 or complements thereof or fragments of either or other nucleic acid molecule of the present invention.

The bacterial recombinant vector may be any vector which can be conveniently subjected to recombinant DNA procedures. The choice of a vector will typically depend on the compatibility of the vector with the bacterial host cell into which the vector is to be introduced. The vector may be a linear or a closed circular plasmid. The vector system may be a single vector or plasmid or two or more vectors or plasmids which together contain the total DNA to be introduced into the genome of the bacterial host. In addition, the bacterial vector may be an

expression vector. Nucleic acid molecules encoding protein homologues or fragments thereof can, for example, be suitably inserted into a replicable vector for expression in the bacterium under the control of a suitable promoter for bacteria. Many vectors are available for this purpose and selection of the appropriate vector will depend mainly on the size of the nucleic acid to be
5 inserted into the vector and the particular host cell to be transformed with the vector. Each vector contains various components depending on its function (amplification of DNA or expression of DNA) and the particular host cell with which it is compatible. The vector components for bacterial transformation generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes and an inducible promoter.

In general, plasmid vectors containing replicon and control sequences that are derived from species compatible with the host cell are used in connection with bacterial hosts. The vector ordinarily carries a replication site, as well as marking sequences that are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is typically
10 transformed using pBR322, a plasmid derived from an *E. coli* species (see, e.g., Bolivar *et al.*, *Gene* 2:95 (1977); the entirety of which is herein incorporated by reference). pBR322 contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR322 plasmid, or other microbial plasmid or phage, also generally contains, or is modified to contain, promoters that can be used by the microbial organism for
15 expression of the selectable marker genes.
20

Nucleic acid molecules encoding protein or fragments thereof may be expressed not only directly, but also as a fusion with another polypeptide, preferably a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature polypeptide. In

general, the signal sequence may be a component of the vector, or it may be a part of the polypeptide DNA that is inserted into the vector. The heterologous signal sequence selected should be one that is recognized and processed (i.e., cleaved by a signal peptidase) by the host cell. For bacterial host cells that do not recognize and process the native polypeptide signal
5 sequence, the signal sequence is substituted by a bacterial signal sequence selected, for example, from the group consisting of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Generally, in cloning vectors this sequence is one that enables the vector to replicate independently of the host chromosomal DNA and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria.

Expression and cloning vectors also generally contain a selection gene, also termed a selectable marker. This gene encodes a protein necessary for the survival or growth of transformed host cells grown in a selective culture medium. Host cells not transformed with the vector containing the selection gene will not survive in the culture medium. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply
20 critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*. One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous protein homologue or fragment thereof produce a protein conferring drug resistance and thus survive the selection regimen.

The expression vector for producing a protein or fragment thereof can also contains an inducible promoter that is recognized by the host bacterial organism and is operably linked to the nucleic acid encoding, for example, the nucleic acid molecule encoding the protein homologue or fragment thereof of interest. Inducible promoters suitable for use with bacterial hosts include the β -lactamase and lactose promoter systems (Chang *et al.*, *Nature* 275:615 (1978); Goeddel *et al.*, *Nature* 281:544 (1979); both of which are herein incorporated by reference in their entirety), the arabinose promoter system (Guzman *et al.*, *J. Bacteriol.* 174:7716-7728 (1992); the entirety of which is herein incorporated by reference), alkaline phosphatase, a tryptophan (trp) promoter system (Goeddel, *Nucleic Acids Res.* 8:4057 (1980); EP 36,776; both of which are herein incorporated by reference in their entirety) and hybrid promoters such as the tac promoter (deBoer *et al.*, *Proc. Natl. Acad. Sci. (USA)* 80:21-25 (1983); the entirety of which is herein incorporated by reference). However, other known bacterial inducible promoters are suitable (Siebenlist *et al.*, *Cell* 20:269 (1980); the entirety of which is herein incorporated by reference).

Promoters for use in bacterial systems also generally contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the polypeptide of interest. The promoter can be removed from the bacterial source DNA by restriction enzyme digestion and inserted into the vector containing the desired DNA.

Construction of suitable vectors containing one or more of the above-listed components employs standard ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored and re-ligated in the form desired to generate the plasmids required. Examples of available bacterial expression vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as Bluescript™ (Stratagene, La Jolla, CA), in which, for example,

encoding an *A. nidulans* protein homologue or fragment thereof homologue, may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke and Schuster, *J. Biol. Chem.* 264:5503-5509 (1989), the entirety of which is herein incorporated by reference);
5 and the like. pGEX vectors (Promega, Madison Wisconsin U.S.A.) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems are designed to include heparin, thrombin or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

Suitable host bacteria for a bacterial vector include archaebacteria and eubacteria, especially eubacteria and most preferably *Enterobacteriaceae*. Examples of useful bacteria include *Escherichia*, *Enterobacter*, *Azotobacter*, *Erwinia*, *Bacillus*, *Pseudomonas*, *Klebsiella*, *Proteus*, *Salmonella*, *Serratia*, *Shigella*, *Rhizobia*, *Vitreoscilla* and *Paracoccus*. Suitable *E. coli* hosts include *E. coli* W3110 (American Type Culture Collection (ATCC) 27,325, Manassas, Virginia U.S.A.), *E. coli* 294 (ATCC 31,446), *E. coli* B and *E. coli* X1776 (ATCC 31,537).
15 These examples are illustrative rather than limiting. Mutant cells of any of the above-mentioned bacteria may also be employed. It is, of course, necessary to select the appropriate bacteria taking into consideration replicability of the replicon in the cells of a bacterium. For example, *E.*
20 *coli*, *Serratia*, or *Salmonella* species can be suitably used as the host when well known plasmids such as pBR322, pBR325, pACYC177, or pKN410 are used to supply the replicon. *E. coli* strain W3110 is a preferred host or parent host because it is a common host strain for recombinant

DNA product fermentations. Preferably, the host cell should secrete minimal amounts of proteolytic enzymes.

Host cells are transfected and preferably transformed with the above-described vectors and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

Numerous methods of transfection are known to the ordinarily skilled artisan, for example, calcium phosphate and electroporation. Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in section 1.82 of Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, New York: Cold Spring Harbor Laboratory Press, (1989), is generally used for bacterial cells that contain substantial cell-wall barriers. Another method for transformation employs polyethylene glycol/DMSO, as described in Chung and Miller (Chung and Miller, *Nucleic Acids Res.* 16:3580 (1988); the entirety of which is herein incorporated by reference). Yet another method is the use of the technique termed electroporation.

Bacterial cells used to produce the polypeptide of interest for purposes of this invention are cultured in suitable media in which the promoters for the nucleic acid encoding the heterologous polypeptide can be artificially induced as described generally, e.g., in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, New York: Cold Spring Harbor Laboratory Press, (1989). Examples of suitable media are given in U.S. Pat. Nos. 5,304,472 and 5,342,763; both of which are incorporated by reference in their entirety.

In addition to the above discussed procedures, practitioners are familiar with the standard resource materials which describe specific conditions and procedures for the construction,

manipulation and isolation of macromolecules (e.g., DNA molecules, plasmids, etc.), generation of recombinant organisms and the screening and isolating of clones, (see for example, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (1989); Mailga *et al.*, *Methods in Plant Molecular Biology*, Cold Spring Harbor Press (1995), the entirety of which is herein incorporated by reference; Birren *et al.*, *Genome Analysis: Analyzing DNA*, 1, Cold Spring Harbor, New York, the entirety of which is herein incorporated by reference).

(f) Computer Readable Media

The nucleotide sequence provided in SEQ ID NO: 1 through SEQ ID NO: 627 or fragment thereof, or complement thereof, or a nucleotide sequence at least 90% identical, preferably 95%, identical even more preferably 99% or 100% identical to the sequence provided in SEQ ID NO: 1 through SEQ ID NO: 627 or fragment thereof, or complement thereof, can be "provided" in a variety of mediums to facilitate use. Such a medium can also provide a subset thereof in a form that allows a skilled artisan to examine the sequences.

A preferred subset of nucleotide sequences are those nucleic acid sequences that encode a first nucleic acid molecule selected from the group consisting of a nucleic acid molecule that encodes a maize or soybean copalyl diphosphate synthase enzyme or complement thereof or fragment of either, a nucleic acid molecule that encodes a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize dehydroquinase synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a soybean dehydroquinase dehydratase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize putative

dehydroquinase dehydratase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean shikimate dehydrogenase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean shikimate kinase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize enolpyruvylshikimate-P-synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean chorismate synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean chorismate mutase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize tyrosine transaminase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean putative tyrosine transaminase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean transaminase A enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a soybean putative transaminase A enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean homogentisic acid dioxygenase enzyme or complement thereof or fragment of either; and a nucleic acid molecule that encodes a maize or soybean geranylgeranylpyrophosphate synthase enzyme or complement thereof or fragment of either.

A further preferred subset of nucleic acid sequences is where the subset of sequences which encode two proteins or fragments thereof, more preferably three proteins or fragments thereof, more preferably four proteins or fragments thereof, more preferably four proteins or fragments thereof, more preferably five proteins or fragments thereof, more preferably six

proteins or fragments thereof, more preferably seven proteins or fragments thereof, more preferably eight proteins or fragments thereof, more preferably nine proteins or fragments thereof, more preferably ten proteins or fragments thereof, more preferably eleven proteins or fragments thereof, more preferably twelve proteins or fragments thereof, more preferably thirteen proteins or fragments thereof, more preferably fourteen proteins or fragments thereof, more preferably fifteen proteins or fragments thereof, more preferably sixteen proteins or fragments thereof, and even more preferably seventeen proteins or fragments thereof. These nucleic acid sequences are selected from the group that encodes a maize or soybean copalyl diphosphate synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean deoxyarabiono-heptulosonate-P-synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean putative deoxyarabiono-heptulosonate-P-synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize dehydroquinase synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a soybean dehydroquinase dehydratase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize putative dehydroquinase dehydratase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean shikimate dehydrogenase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean shikimate kinase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize enolpyruvylshikimate-P-synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean chorismate synthase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean chorismate mutase enzyme or complement thereof or

fragment of either; a nucleic acid molecule that encodes a maize tyrosine transaminase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean putative tyrosine transaminase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean transaminase A enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a soybean putative transaminase A enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean 4-hydroxyphenylpyruvate dioxygenase enzyme or complement thereof or fragment of either; a nucleic acid molecule that encodes a maize or soybean homogentisic acid dioxygenase enzyme or complement thereof or fragment of either; and a nucleic acid molecule that encodes a maize or soybean geranylgeranylpyrophosphate synthase enzyme or complement thereof or fragment of either.

In one application of this embodiment, a nucleotide sequence of the present invention can be recorded on computer readable media. As used herein, "computer readable media" refers to any medium that can be read and accessed directly by a computer. Such media include, but are not limited to: magnetic storage media, such as floppy discs, hard disc, storage medium and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage media. A skilled artisan can readily appreciate how any of the presently known computer readable mediums can be used to create a manufacture comprising computer readable medium having recorded thereon a nucleotide sequence of the present invention.

As used herein, "recorded" refers to a process for storing information on computer readable medium. A skilled artisan can readily adopt any of the presently known methods for recording information on computer readable medium to generate media comprising the

nucleotide sequence information of the present invention. A variety of data storage structures are available to a skilled artisan for creating a computer readable medium having recorded thereon a nucleotide sequence of the present invention. The choice of the data storage structure will generally be based on the means chosen to access the stored information. In addition, a variety
5 of data processor programs and formats can be used to store the nucleotide sequence information of the present invention on computer readable medium. The sequence information can be represented in a word processing text file, formatted in commercially-available software such as WordPerfect and Microsoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like. A skilled artisan can readily adapt any number of data processor structuring formats (e.g. text file or database) in order to
10 obtain computer readable medium having recorded thereon the nucleotide sequence information of the present invention.

By providing one or more of nucleotide sequences of the present invention, a skilled artisan can routinely access the sequence information for a variety of purposes. Computer software is publicly available which allows a skilled artisan to access sequence information
15 provided in a computer readable medium. The examples which follow demonstrate how software which implements the BLAST (Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990), the entirety of which is herein incorporated by reference) and BLAZE (Brutlag *et al.*, *Comp. Chem.* 17:203-207 (1993), the entirety of which is herein incorporated by reference) search algorithms
20 on a Sybase system can be used to identify open reading frames (ORFs) within the genome that contain homology to ORFs or proteins from other organisms. Such ORFs are protein-encoding fragments within the sequences of the present invention and are useful in producing commercially important proteins such as enzymes used in amino acid biosynthesis, metabolism,

transcription, translation, RNA processing, nucleic acid and a protein degradation, protein modification and DNA replication, restriction, modification, recombination and repair.

The present invention further provides systems, particularly computer-based systems, which contain the sequence information described herein. Such systems are designed to identify commercially important fragments of the nucleic acid molecule of the present invention. As used herein, "a computer-based system" refers to the hardware means, software means and data storage means used to analyze the nucleotide sequence information of the present invention. The minimum hardware means of the computer-based systems of the present invention comprises a central processing unit (CPU), input means, output means and data storage means. A skilled artisan can readily appreciate that any one of the currently available computer-based system are suitable for use in the present invention.

As indicated above, the computer-based systems of the present invention comprise a data storage means having stored therein a nucleotide sequence of the present invention and the necessary hardware means and software means for supporting and implementing a search means. As used herein, "data storage means" refers to memory that can store nucleotide sequence information of the present invention, or a memory access means which can access manufactures having recorded thereon the nucleotide sequence information of the present invention. As used herein, "search means" refers to one or more programs which are implemented on the computer-based system to compare a target sequence or target structural motif with the sequence information stored within the data storage means. Search means are used to identify fragments or regions of the sequence of the present invention that match a particular target sequence or target motif. A variety of known algorithms are disclosed publicly and a variety of commercially available software for conducting search means are available can be used in the computer-based

systems of the present invention. Examples of such software include, but are not limited to, MacPattern (EMBL), BLASTIN and BLASTIX (NCBIA). One of the available algorithms or implementing software packages for conducting homology searches can be adapted for use in the present computer-based systems.

5 The most preferred sequence length of a target sequence is from about 10 to 100 amino acids or from about 30 to 300 nucleotide residues. However, it is well recognized that during searches for commercially important fragments of the nucleic acid molecules of the present invention, such as sequence fragments involved in gene expression and protein processing, may be of shorter length.

10 As used herein, "a target structural motif," or "target motif," refers to any rationally selected sequence or combination of sequences in which the sequences the sequence(s) are chosen based on a three-dimensional configuration which is formed upon the folding of the target motif. There are a variety of target motifs known in the art. Protein target motifs include, but are not limited to, enzymatic active sites and signal sequences. Nucleic acid target motifs include,
15 but are not limited to, promoter sequences, *cis* elements, hairpin structures and inducible expression elements (protein binding sequences).

20 Thus, the present invention further provides an input means for receiving a target sequence, a data storage means for storing the target sequences of the present invention sequence identified using a search means as described above and an output means for outputting the identified homologous sequences. A variety of structural formats for the input and output means can be used to input and output information in the computer-based systems of the present invention. A preferred format for an output means ranks fragments of the sequence of the present invention by varying degrees of homology to the target sequence or target motif. Such

presentation provides a skilled artisan with a ranking of sequences which contain various amounts of the target sequence or target motif and identifies the degree of homology contained in the identified fragment.

A variety of comparing means can be used to compare a target sequence or target motif with the data storage means to identify sequence fragments sequence of the present invention. For example, implementing software which implement the BLAST and BLAZE algorithms (Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990)) can be used to identify open frames within the nucleic acid molecules of the present invention. A skilled artisan can readily recognize that any one of the publicly available homology search programs can be used as the search means for the computer-based systems of the present invention.

Having now generally described the invention, the same will be more readily understood through reference to the following examples which are provided by way of illustration and are not intended to be limiting of the present invention, unless specified.

Example 1

The MONN01 cDNA library is a normalized library generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) total leaf tissue at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in

the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The older, more juvenile leaves, which are in a basal position, as well as the younger, more adult leaves, which are more apical are cut at the base of the leaves. The leaves are then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are crushed. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON001 cDNA library is generated from maize (B73, Illinois Foundation Seeds, Champaign, Illinois U.S.A.) immature tassels at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in a greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue from the maize plant is collected at the V6 stage. At that stage the tassel is an immature tassel of about 2-3 cm in length. The tassels are removed and frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON003 library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign, Illinois U.S.A.) roots at the V6 developmental stage. Seeds are planted at a depth of approximately 3 cm in coil into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth, the seedlings are transplanted into 10 inch pots containing the Metro 200 growing medium. Plants are watered daily before transplantation and approximately 3 times a week after transplantation. Peters 15-16-17 fertilizer is applied approximately three times per week after transplanting at a concentration of 150 ppm N. Two to three times during the life time of the plant from transplanting to flowering a total of approximately 900 mg Fe is added to each pot. Maize plants are grown in the green house in approximately 15hr day/9hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6 leaf development stage. The root system is cut from maize plant and washed with water to free it from the soil. The tissue is then immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON004 cDNA library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign, Illinois U.S.A.) total leaf tissue at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are

grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The older, more juvenile leaves, which are in a basal position, as well as the younger, more adult leaves, which are more apical are cut at the base of the leaves. The leaves are then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are crushed. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON005 cDNA library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign Illinois, U.S.A.) root tissue at the V6 development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The root system is cut from the mature maize plant and washed with water to free it from the soil. The tissue is immediately frozen in liquid nitrogen and the harvested tissue is then

stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON006 cDNA library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign Illinois, U.S.A.) total leaf tissue at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The older more juvenile leaves, which are in a basal position, as well as the younger more adult leaves, which are more apical are cut at the base of the leaves. The leaves are then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are crushed. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON007 cDNA library is generated from the primary root tissue of 5 day old maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) seedlings. Seeds are planted on a moist filter paper on a covered tray that is kept in the dark until germination (one day). After

germination, the trays, along with the moist paper, are moved to a greenhouse where the maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles for approximately 5 days. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. The primary root tissue is collected when the seedlings are 5 days old. At this stage, the primary root (radicle) is pushed through the coleorhiza which itself is pushed through the seed coat. The primary root, which is about 2-3 cm long, is cut and immediately frozen in liquid nitrogen and then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON008 cDNA library is generated from the primary shoot (coleoptile 2-3 cm) of maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) seedlings which are approximately 5 days old. Seeds are planted on a moist filter paper on a covered tray that is kept in the dark until germination (one day). Then the trays containing the seeds are moved to a greenhouse at 15hr daytime/9 hr nighttime cycles and grown until they are 5 days post germination. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Tissue is collected when the seedlings are 5 days old. At this stage, the primary shoot (coleoptile) is pushed through the seed coat and is about 2-3 cm long. The coleoptile is dissected away from the rest of the seedling, immediately frozen in liquid nitrogen and then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON009 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) leaves at the 8 leaf stage (V8 plant development stage). Seeds are

planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is 80°F and the nighttime temperature is 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 8-leaf development stage. The older more juvenile leaves, which are in a basal position, as well as the younger more adult leaves, which are more apical, are cut at the base of the leaves. The leaves are then pooled and then immediately transferred to liquid nitrogen containers in which the pooled leaves are crushed. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON010 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) root tissue at the V8 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house

in 15 hr day/9 hr night cycles. The daytime temperature is 80°F and the nighttime temperature is 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the V8 development stage. The root system is cut from this mature maize plant and washed with water to free it from the soil. The tissue is immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON011 cDNA library is generated from undeveloped maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) leaf at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The second youngest leaf which is at the base of the apical leaf of V6 stage maize plant is cut at the base and immediately transferred to liquid nitrogen containers in which the leaf is crushed. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON012 cDNA library is generated from 2 day post germination maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) seedlings. Seeds are planted on a moist filter paper on a covered tray that is kept in the dark until germination (one day). Then the trays containing the seeds are moved to the greenhouse and grown at 15hr daytime/9 hr nighttime cycles until 2 days post germination. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Tissue is collected when the seedlings are 2 days old. At the two day stage, the coleorhiza is pushed through the seed coat and the primary root (the radicle) is pierced the coleorhiza but is barely visible. Also, at this two day stage, the coleoptile is just emerging from the seed coat. The 2 days post germination seedlings are then immersed in liquid nitrogen and crushed. The harvested tissue is stored at -80°C until preparation of total RNA. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON013 cDNA library is generated from apical maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) meristem founder at the V4 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000

W sodium vapor lamps. Prior to tissue collection, the plant is at the 4 leaf stage. The lead at the apex of the V4 stage maize plant is referred to as the meristem founder. This apical meristem founder is cut, immediately frozen in liquid nitrogen and crushed. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the

5 cDNA library is constructed as described in Example 2.

The SATMON014 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) endosperm fourteen days after pollination. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium.

10 Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. After

15 the V10 stage, the maize plant ear shoots are ready for fertilization. At this stage, the ear shoots are enclosed in a paper bag before silk emergence to withhold the pollen. The ear shoots are pollinated and 14 days after pollination, the ears are pulled out and then the kernels are plucked out of the ears. Each kernel is then dissected into the embryo and the endosperm and the

20 aleurone layer is removed. After dissection, the endosperms are immediately frozen in liquid nitrogen and then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON016 library is a maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) sheath library collected at the V8 developmental stage. Seeds are planted in a depth of approximately 3 cm in solid into 2-3 inch pots containing Metro growing medium. After 2-3 weeks growth, they are transplanted into 10" pots containing the same. Plants are watered daily before transplantation and approximately the times a week after transplantation. Peters 15-16-17 fertilizer is applied approximately three times per week after transplanting, at a strength of 150 ppm N. Two to three times during the life time of the plant from transplanting to flowering, a total of approximately 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15hr day/9hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. When the maize plants are at the V8 stage the 5th and 6th leaves from the bottom exhibit fully developed leaf blades. At the base of these leaves, the ligule is differentiated and the leaf blade is joined to the sheath. The sheath is dissected away from the base of the leaf then the sheath is frozen in liquid nitrogen and crushed. The tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON017 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) embryo seventeen days after pollination. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth the seeds are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a

total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. After the V10 stage, the ear shoots of maize plant, which are ready for fertilization, are enclosed in a paper bag before silk emergence to withhold the pollen. The ear shoots are fertilized and 21 days after pollination, the ears are pulled out and the kernels are plucked out of the ears. Each kernel is then dissected into the embryo and the endosperm and the aleurone layer is removed. After dissection, the embryos are immediately frozen in liquid nitrogen and then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON019 (Lib3054) cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) culm (stem) at the V8 developmental stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. When the maize plant is at the V8 stage, the 5th and 6th leaves from the bottom have fully developed leaf blades. The region between the nodes of the 5th and the sixth leaves from

the bottom is the region of the stem that is collected. The leaves are pulled out and the sheath is also torn away from the stem. This stem tissue is completely free of any leaf and sheath tissue. The stem tissue is then frozen in liquid nitrogen and stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in

5 Example 2.

The SATMON020 cDNA library is from a maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) Hill Type II-Initiated Callus. Petri plates containing approximately 25 ml of Type II initiation media are prepared. This medium contains N6 salts and vitamins, 3% sucrose, 2.3 g/liter proline 0.1 g/liter enzymatic casein hydrolysate, 2mg/liter 2,4 – dichloro phenoxy-acetic acid (2,4, D), 15.3 mg/liter AgNO₃ and 0.8% bacto agar and is adjusted to pH 6.0 before autoclaving. At 9-11 days after pollination, an ear with immature embryos measuring approximately 1-2 mm in length is chosen. The husks and silks are removed and then the ear is broken into halves and placed in an autoclaved solution of Clorox/TWEEN 20 sterilizing solution. Then the ear is rinsed with deionized water. Then each embryo is extracted from the kernel. Intact embryos are placed in contact with the medium, scutellar side up). Multiple embryos are plated on each plate and the plates are incubated in the dark at 25°C. Type II calluses are friable, can be subcultured with a spatula, frequently regenerate via somatic embryogenesis and are relatively undifferentiated. As seen in the microscope, the Tape II calluses show color ranging from translucent to light yellow and heterogeneity on with respect to

20 embryoid structure as well as stage of embryoid development. Once Type II callus are formed, the calluses is transferred to type II callus maintenance medium without AgNO₃. Every 7-10 days, the callus is subcultured. About 4 weeks after embryo isolation the callus is removed from

the plates and then frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON021 cDNA library is generated from the immature maize (DK604, Dekalb Genetics, Dekalb Illinois, U.S.A.) tassel at the V8 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F . Supplemental lighting is provided by 1000 W sodium vapor lamps. As the maize plant enters the V8 stage, tassels which are 15-20 cm in length are collected and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON022 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) ear (growing silks) at the V8 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after

transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the plant is in the V8 stage. At this stage, some immature ear shoots are visible. The immature ear shoots (approximately 1 cm in length) are pulled out, frozen in liquid nitrogen and then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON23 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) ear (growing silk) at the V8 development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. When the tissue is harvested at the V8 stage, the length of the ear that is harvested is about 10-15 cm and the silks are just exposed (approximately 1 inch). The ear along with the silks is frozen in liquid nitrogen and then the tissue is stored at -80°C until

RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON024 cDNA library is generated from the immature maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) tassel at the V9 development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. As a maize plant enters the V9 stage, the tassel is rapidly developing and a 37 cm tassel along with the glume, anthers and pollen is collected and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON025 cDNA library is from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) Hill Type II-Regenerated Callus. Type II callus is grown in initiation media as described for SATMON020 and then the embryoids on the surface of the Type II callus are allowed to mature and germinate. The 1-2 gm fresh weight of the soft friable type callus containing numerous embryoids are transferred to 100 x 15 mm petri plates containing 25 ml of regeneration media. Regeneration media consists of Murashige and Skoog (MS) basal salts, modified White's vitamins (0.2 g/liter glycine and 0.5 g/liter myo-inositol) and 0.8% bacto agar

(6SMS0D)). The plates are then placed in the dark after covering with parafilm. After 1 week, the plates are moved to a lighted growth chamber with 16 hr light and 8 hr dark photoperiod. Three weeks after plating the Type II callus to 6SMS0D, the callus exhibit shoot formation. The callus and the shoots are transferred to fresh 6SMS0D plates for another 2 weeks. The callus and the shoots are then transferred to petri plates with reduced sucrose (3SMS0D). Upon distinct formation of a root and shoot, the newly developed green plants are then removed out with a spatula and frozen in liquid nitrogen containers. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON026 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) juvenile/adult shift leaves at the V8 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F . Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plants are at the 8-leaf development stage. Leaves are founded sequentially around the meristem over weeks of time and the older, more juvenile leaves arise earlier and in a more basal position than the younger, more adult leaves, which are in a

more apical position. In a V8 plant, some leaves which are in the middle portion of the plant exhibit characteristics of both juvenile as well as adult leaves. They exhibit a yellowing color but also exhibit, in part, a green color. These leaves are termed juvenile/adult shift leaves. The juvenile/adult shift leaves (the 4th, 5th leaves from the bottom) are cut at the base, pooled and transferred to liquid nitrogen in which they are then crushed. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON027 cDNA library is generated from 6 day maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) leaves. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the Metro 200 growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Prior to tissue collection, when the plant is at the 8-leaf stage, water is held back for six days. The older, more juvenile leaves, which are in a basal position, as well as the younger, more adult leaves, which are more apical, are all cut at the base of the leaves. All the leaves exhibit significant wilting. The leaves are then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are then crushed. The harvested tissue is then stored at -80°C until RNA

preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON028 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) roots at the V8 developmental stage that are subject to six days water stress. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the Metro 200 growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Prior to tissue collection, when the plant is at the 8-leaf stage, water is held back for six days. The root system is cut, shaken and washed to remove soil. Root tissue is then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are then crushed. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON029 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) seedlings at the etiolated stage. Seeds are planted on a moist filter paper on a covered tray that is kept in the dark for 4 days at approximately 70°F. Tissue is collected when the seedlings are 4 days old. By 4 days, the primary root has penetrated the coleorhiza and

is about 4-5 cm and the secondary lateral roots have also made their appearance. The coleoptile has also pushed through the seed coat and is about 4-5 cm long. The seedlings are frozen in liquid nitrogen and crushed. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

5 The SATMON030 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) root tissue at the V4 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth, they are transplanted into 10 inch pots containing the same. Plants are watered daily before transplantation and approximately 3 times a week after transplantation. Peters 15-16-17 fertilizer is applied approximately three times per week after transplanting, at a strength of 150 ppm N. Two to three times during the life time of the plant, from transplanting to flowering, a total of approximately 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15hr day/9hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 sodium vapor lamps. Tissue is collected when the maize plant is at the 4 leaf development stage. The root system is cut from the mature maize plant and washed with water to free it from the soil. The tissue is then immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

20 The SATMON031 cDNA library is generated from the maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) leaf tissue at the V4 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium.

After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is 80°F and the nighttime temperature is 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 4-leaf development stage. The third leaf from the bottom is cut at the base and immediately frozen in liquid nitrogen and crushed. The tissue is immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON033 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) embryo tissue 13 days after pollination. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. After the V10 stage, the ear shoots of the maize plant, which are ready for fertilization, are enclosed in

a paper bag before silk emergent to withhold the pollen. The ear shoots are pollinated and 13 days after pollination, the ears are pulled out and then the kernels are plucked out of the ears. Each kernel is then dissected into the embryo and the endosperm and the aleurone layer is removed. After dissection, the embryos are immediately frozen in liquid nitrogen and then
5 stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON034 cDNA library is generated from cold stressed maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) seedlings. Seeds are planted on a moist filter paper on a covered tray that is kept on at 10°C for 7 days. After 7 days, the temperature is shifted to 15°C
10 for one day until germination of the seed. Tissue is collected once the seedlings are 1 day old. At this point, the coleorhiza has just pushed out of the seed coat and the primary root is just making its appearance. The coleoptile has not yet pushed completely through the seed coat and is also just making its appearance. These 1 day old cold stressed seedlings are frozen in liquid nitrogen and crushed. The harvested tissue is then stored at -80°C until RNA preparation. The
15 RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMON~001 (Lib36, Lib83, Lib84) cDNA library is generated from maize leaves at the V8 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are
20 transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times

during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in a greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F.

Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue from the maize plant

5 is collected at the V8 stage. The older more juvenile leaves in a basal position as well as the younger more adult leaves which are more apical are all cut at the base, pooled and frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SATMONN01 cDNA library is generated from maize (B73, Illinois Foundation
10 Seeds, Champaign, Illinois U.S.A.) normalized immature tassels at the V6 plant development stage normalized tissue. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before
15 transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in a greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F.

Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue from the maize plant
20 is collected at the V6 stage. At that stage the tassel is an immature tassel of about 2-3 cm in length. The tassels are removed and frozen in liquid nitrogen. The harvested tissue is then

stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the normalized cDNA library is constructed as described in Example 2.

The SATMONN04 cDNA library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign, Illinois U.S.A.) normalized total leaf tissue at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The older, more juvenile leaves, which are in a basal position, as well as the younger, more adult leaves, which are more apical are cut at the base of the leaves. The leaves are then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are crushed. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the normalized cDNA library is constructed as described in Example 2.

The SATMONN05 cDNA library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign Illinois, U.S.A.) normalized root tissue at the V6 development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing

Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The root system is cut from the mature maize plant and washed with water to free it from the soil. The tissue is immediately frozen in liquid nitrogen and the harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the normalized cDNA library is constructed as described in Example 2.

The SATMONN06 cDNA library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign Illinois, U.S.A.) normalized total leaf tissue at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental

lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The older more juvenile leaves, which are in a basal position, as well as the younger more adult leaves, which are more apical are cut at the base of the leaves. The leaves are then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are crushed. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the normalized cDNA library is constructed as described in Example 2.

The CMZ029 (SATMON036) cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) endosperm 22 days after pollination. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. After the V10 stage, the ear shoots of the maize plant, which are ready for fertilization, are enclosed in a paper bag before silk emergent to withhold the pollen. The ear shoots are pollinated and 22 days after pollination, the ears are pulled out and then the kernels are plucked out of the ears. Each kernel is then dissected into the embryo and the endosperm and the alurone layer is removed. After dissection, the endosperms are immediately frozen in liquid nitrogen and

then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz030 (Lib143) cDNA library is generated from maize seedling tissue two days post germination. Seeds are planted on a moist filter paper on a covered try that is keep in the dark until germination. The trays are then moved to the bench top at 15 hr daytime/9 hr nighttime cycles for 2 days post-germination. The day time temperature is 80°F and the nighttime temperature is 70°F . Tissue is collected when the seedlings are 2 days old. At this stage, the colehrhiza has pushed through the seed coat and the primary root (the radicle) is just piercing the colehrhiza and is barely visible. The seedlings are placed at 42°C for 1 hour. Following the heat shock treatment, the seedlings are immersed in liquid nitrogen and crushed. The harvested tissue is stored at -80° until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz031 (Lib148) cDNA library is generated from maize pollen tissue at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F . Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from

V10+ stage plants. The ear shoots, which are ready for fertilization, are enclosed in a paper bag to withhold pollen. Twenty-one days after pollination, prior to removing the ears, the paper bag is shaken to collect the mature pollen. The mature pollen is immediately frozen in liquid nitrogen containers and the pollen is crushed. The harvested tissue is then stored at -80°C until
5 RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz033 (Lib189) cDNA library is generated from maize pooled leaf tissue. Samples are harvested from open pollinated plants. Tissue is collected from maize leaves at the anthesis stage. The leaves are collect from from 10-12 plants and frozen in liquid nitrogen. The
10 harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz034 (Lib3060) cDNA library is generated from maize mature tissue at 40 days post pollination plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are
15 transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime
20 temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from leaves located two leaves below the ear leaf. This sample represents those genes expressed

during onset and early stages of leaf senescence. The leaves are pooled and immediately transferred to liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

- 5 The CMz035 (Lib3061) cDNA library is generated from maize endosperm tissue at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F . Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from V10+ stage plants. The ear shoots, which are ready for fertilization, are enclosed in a paper bag prior to silk emergence to withhold pollen. Thirty-two days after pollination, the ears are pulled out and the kernels are removed from the cob. Each kernel is dissected into the embryo and the endosperm and the aleurone layer is removed. After dissection, the endosperms are immediately transferred to liquid nitrogen. The harvested tissue is then stored at 80°C until RNA preparation.
- 20 The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz036 (Lib3062) cDNA library is generated from maize husk tissue at the 8 week old plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before
5 transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from 8
10 week old plants. The husk is separated from the ear and immediately transferred to liquid nitrogen containers. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz037 (Lib3059) cDNA library is generated from maize pooled kernal at 12-15
15 days after pollination plant development stage. Sample were collected from field grown material. Whole kernels from hand pollinated (control pollination) are harvested as whole ears and immediately frozen on dry ice. Kernels from 10-12 ears were pooled and ground together in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is
20 purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz039 (Lib3066) cDNA library is generated from maize immature anther tissue at the 7 week old immature tassel stage. Seeds are planted at a depth of approximately 3 cm into 2-

3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 7 week old immature tassel stage. At this stage, prior to anthesis, the immature anthers are green and enclosed in the staminate spikelet. The developing anthers are dissected away from the 7 week old immature tassel and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz040 (Lib3067) cDNA library is generated from maize kernel tissue at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F.

Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from V10+ stage plants. The ear shoots, which are ready for fertilization, are enclosed in a paper bag before silk emergence to withhold pollen. Five to eight days after controlled pollination. The ears are pulled and the kernels removed. The kernels are immediately frozen in liquid nitrogen.

- 5 This sample represents genes expressed in early kernel development, during periods of cell division, amyloplast biogenesis and early carbon flow across the material to filial tissue. The harvested kernels tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz041 (Lib3068) cDNA library is generated from maize pollen germinating silk tissue at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F.

- Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from V10+ stage plants when the ear shoots are ready for fertilization at the silk emergence stage. The emerging silks are pollinated with an excess of pollen under controlled pollination conditions in the green house. Eighteen hours after pollination the silks are removed from the ears and immediately frozen in liquid nitrogen. This sample represents genes expressed in both pollen
- 20

and silk tissue early in pollination. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz042 (Lib3069) cDNA library is generated from maize ear tissue excessively
5 pollinated at the V10+ plant development stage. Seeds are planted at a depth of approximately 3
cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they
are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily
before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is
applied three times per week after transplanting at a strength of 150 ppm N. Two to three times
10 during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to
each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime
temperature is approximately 80°F and the nighttime temperature is approximately 70°F .
Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from
V10+ stage plants and the ear shoots which are ready for fertilization are at the silk emergence
15 stage. The immature ears are pollinated with an excess of pollen under controlled pollination
conditions. Eighteen hours post-pollination, the ears are removed and immediately transferred to
liquid nitrogen containers. The harvested tissue is then stored at -80°C until RNA preparation.
The RNA is purified from the stored tissue and the cDNA library is constructed as described in
Example 2.

20 The CMz044 (Lib3075) cDNA library is generated from maize microspore tissue at the
V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch
peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted

into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from immature anthers from 7 week old tassels. The immature anthers are first dissected from the 7 week old tassel with a scalpel on a glass slide covered with water. The microspores (immature pollen) are released into the water and are recovered by centrifugation. The microspore suspension is immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz045 (Lib3076) cDNA library is generated from maize immature ear megaspore tissue. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental

lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from immature ear (megaspore) obtained from 7 week old plants. The immature ears are harvested from the 7 week old plants and are approximately 2.5 to 3 cm in length. The kernels are removed from the cob immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz047 (Lib3078) cDNA library is generated from maize CO₂ treated high-exposure shoot tissue at the V10+ plant development stage. RX601 maize seeds are sterilized for 1 minute with a 10% clorox solution. The seeds are rolled in germination paper, and germinated in 0.5 mM calcium sulfate solution for two days at 30°C. The seedlings are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium at a rate of 2-3 seedlings per pot. Twenty pots are placed into a high CO₂ environment (approximately 1000 ppm CO₂). Twenty plants were grown under ambient greenhouse CO₂ (approximately 450 ppm CO₂). Plants are watered daily before transplantation and three times a week after transplantation. Peters 20-20-20 fertilizer is also lightly applied. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. At ten days post planting, the shoots from both atmosphere are frozen in liquid nitrogen and lightly ground. The roots are washed in deionized water to remove the support media and the tissue is immediately transferred to liquid nitrogen containers. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

10
20
30
40
50
60
70
80
90
100
110
120
130
140
150
160
170
180
190
200
210
220
230
240
250
260
270
280
290
300
310
320
330
340
350
360
370
380
390
400
410
420
430
440
450
460
470
480
490
500
510
520
530
540
550
560
570
580
590
600
610
620
630
640
650
660
670
680
690
700
710
720
730
740
750
760
770
780
790
800
810
820
830
840
850
860
870
880
890
900
910
920
930
940
950
960
970
980
990
1000
1010
1020
1030
1040
1050
1060
1070
1080
1090
1100
1110
1120
1130
1140
1150
1160
1170
1180
1190
1200
1210
1220
1230
1240
1250
1260
1270
1280
1290
1300
1310
1320
1330
1340
1350
1360
1370
1380
1390
1400
1410
1420
1430
1440
1450
1460
1470
1480
1490
1500
1510
1520
1530
1540
1550
1560
1570
1580
1590
1600
1610
1620
1630
1640
1650
1660
1670
1680
1690
1700
1710
1720
1730
1740
1750
1760
1770
1780
1790
1800
1810
1820
1830
1840
1850
1860
1870
1880
1890
1900
1910
1920
1930
1940
1950
1960
1970
1980
1990
2000
2010
2020
2030
2040
2050
2060
2070
2080
2090
2100
2110
2120
2130
2140
2150
2160
2170
2180
2190
2200
2210
2220
2230
2240
2250
2260
2270
2280
2290
2300
2310
2320
2330
2340
2350
2360
2370
2380
2390
2400
2410
2420
2430
2440
2450
2460
2470
2480
2490
2500
2510
2520
2530
2540
2550
2560
2570
2580
2590
2600
2610
2620
2630
2640
2650
2660
2670
2680
2690
2700
2710
2720
2730
2740
2750
2760
2770
2780
2790
2800
2810
2820
2830
2840
2850
2860
2870
2880
2890
2900
2910
2920
2930
2940
2950
2960
2970
2980
2990
3000
3010
3020
3030
3040
3050
3060
3070
3080
3090
3100
3110
3120
3130
3140
3150
3160
3170
3180
3190
3200
3210
3220
3230
3240
3250
3260
3270
3280
3290
3300
3310
3320
3330
3340
3350
3360
3370
3380
3390
3400
3410
3420
3430
3440
3450
3460
3470
3480
3490
3500
3510
3520
3530
3540
3550
3560
3570
3580
3590
3600
3610
3620
3630
3640
3650
3660
3670
3680
3690
3700
3710
3720
3730
3740
3750
3760
3770
3780
3790
3800
3810
3820
3830
3840
3850
3860
3870
3880
3890
3900
3910
3920
3930
3940
3950
3960
3970
3980
3990
4000
4010
4020
4030
4040
4050
4060
4070
4080
4090
4100
4110
4120
4130
4140
4150
4160
4170
4180
4190
4200
4210
4220
4230
4240
4250
4260
4270
4280
4290
4300
4310
4320
4330
4340
4350
4360
4370
4380
4390
4400
4410
4420
4430
4440
4450
4460
4470
4480
4490
4500
4510
4520
4530
4540
4550
4560
4570
4580
4590
4600
4610
4620
4630
4640
4650
4660
4670
4680
4690
4700
4710
4720
4730
4740
4750
4760
4770
4780
4790
4800
4810
4820
4830
4840
4850
4860
4870
4880
4890
4900
4910
4920
4930
4940
4950
4960
4970
4980
4990
5000
5010
5020
5030
5040
5050
5060
5070
5080
5090
5100
5110
5120
5130
5140
5150
5160
5170
5180
5190
5200
5210
5220
5230
5240
5250
5260
5270
5280
5290
5300
5310
5320
5330
5340
5350
5360
5370
5380
5390
5400
5410
5420
5430
5440
5450
5460
5470
5480
5490
5500
5510
5520
5530
5540
5550
5560
5570
5580
5590
5600
5610
5620
5630
5640
5650
5660
5670
5680
5690
5700
5710
5720
5730
5740
5750
5760
5770
5780
5790
5800
5810
5820
5830
5840
5850
5860
5870
5880
5890
5900
5910
5920
5930
5940
5950
5960
5970
5980
5990
6000
6010
6020
6030
6040
6050
6060
6070
6080
6090
6100
6110
6120
6130
6140
6150
6160
6170
6180
6190
6200
6210
6220
6230
6240
6250
6260
6270
6280
6290
6300
6310
6320
6330
6340
6350
6360
6370
6380
6390
6400
6410
6420
6430
6440
6450
6460
6470
6480
6490
6500
6510
6520
6530
6540
6550
6560
6570
6580
6590
6600
6610
6620
6630
6640
6650
6660
6670
6680
6690
6700
6710
6720
6730
6740
6750
6760
6770
6780
6790
6800
6810
6820
6830
6840
6850
6860
6870
6880
6890
6900
6910
6920
6930
6940
6950
6960
6970
6980
6990
7000
7010
7020
7030
7040
7050
7060
7070
7080
7090
7100
7110
7120
7130
7140
7150
7160
7170
7180
7190
7200
7210
7220
7230
7240
7250
7260
7270
7280
7290
7300
7310
7320
7330
7340
7350
7360
7370
7380
7390
7400
7410
7420
7430
7440
7450
7460
7470
7480
7490
7500
7510
7520
7530
7540
7550
7560
7570
7580
7590
7600
7610
7620
7630
7640
7650
7660
7670
7680
7690
7700
7710
7720
7730
7740
7750
7760
7770
7780
7790
7800
7810
7820
7830
7840
7850
7860
7870
7880
7890
7900
7910
7920
7930
7940
7950
7960
7970
7980
7990
8000
8010
8020
8030
8040
8050
8060
8070
8080
8090
8100
8110
8120
8130
8140
8150
8160
8170
8180
8190
8200
8210
8220
8230
8240
8250
8260
8270
8280
8290
8300
8310
8320
8330
8340
8350
8360
8370
8380
8390
8400
8410
8420
8430
8440
8450
8460
8470
8480
8490
8500
8510
8520
8530
8540
8550
8560
8570
8580
8590
8600
8610
8620
8630
8640
8650
8660
8670
8680
8690
8700
8710
8720
8730
8740
8750
8760
8770
8780
8790
8800
8810
8820
8830
8840
8850
8860
8870
8880
8890
8900
8910
8920
8930
8940
8950
8960
8970
8980
8990
9000
9010
9020
9030
9040
9050
9060
9070
9080
9090
9100
9110
9120
9130
9140
9150
9160
9170
9180
9190
9200
9210
9220
9230
9240
9250
9260
9270
9280
9290
9300
9310
9320
9330
9340
9350
9360
9370
9380
9390
9400
9410
9420
9430
9440
9450
9460
9470
9480
9490
9500
9510
9520
9530
9540
9550
9560
9570
9580
9590
9600
9610
9620
9630
9640
9650
9660
9670
9680
9690
9700
9710
9720
9730
9740
9750
9760
9770
9780
9790
9800
9810
9820
9830
9840
9850
9860
9870
9880
9890
9900
9910
9920
9930
9940
9950
9960
9970
9980
9990
10000

The CMz048 (Lib3079) cDNA library is generated from maize basal endosperm transfer layer tissue at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from V10+ maize plants. The ear shoots, which are ready for fertilization, are enclosed in a paper bag prior to silk emergence, to withhold the pollen. Kernels are harvested at 12 days post-pollination and placed on wet ice for dissection. The kernels are cross sectioned laterally, dissecting just above the pedicel region, including 1-2 mm of the lower endosperm and the basal endosperm transfer region. The pedicel and lower endosperm region containing the basal endosperm transfer layer is pooled and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz049(Lib3088) cDNA library is generated from maize immature anther tissue at the 7 week old immature tassel stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily

before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F.

Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 7 week old immature tassel stage. At this stage, prior to anthesis, the immature anthers are green and enclosed in the staminate spikelet. The developing anthers are dissected away from the 7 week old immature tassel and immediately transferred to liquid nitrogen container. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The CMz050 (Lib3114) cDNA library is generated from maize silk tissue at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the

maize plant is beyond the 10-leaf development stage and the ear shoots are approximately 15-20 cm in length. The ears are pulled and silks are separated from the ears and immediately transferred to liquid nitrogen containers. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON001 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) total leaf tissue at the V4 plant development stage. Leaf tissue from 38, field grown V4 stage plants is harvested from the 4th node. Leaf tissue is removed from the plants and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON002 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) root tissue at the V4 plant development stage. Root tissue from 76, field grown V4 stage plants is harvested. The root systems is cut from the soybean plant and washed with water to free it from the soil and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON003 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seedling hypocotyl axis tissue harvested 2 day post-imbibition. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium. Trays are placed in an environmental chamber and grown at 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the

nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Tissue is collected 2 days after the start of imbibition. The 2 days after imbibition samples are separated into 3 collections after removal of any adhering seed coat. At the 2 day stage, the hypocotyl axis is emerging from the soil. A few seedlings have cracked the soil surface and exhibited slight greening of the exposed cotyledons. The seedlings are washed in water to remove soil, hypocotyl axis harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON004 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seedling cotyledon tissue harvested 2 day post-imbibition. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium. Trays are placed in an environmental chamber and grown at 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Tissue is collected 2 days after the start of imbibition. The 2 days after imbibition samples are separated into 3 collections after removal of any adhering seed coat. At the 2 day stage, the hypocotyl axis is emerging from the soil. A few seedlings have cracked the soil surface and exhibited slight greening of the exposed cotyledons. The seedlings are washed in water to remove soil, hypocotyl axis harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON005 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seedling hypocotyl axis tissue harvested 6

hour post-imbibition. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium. Trays are placed in an environmental chamber and grown at 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Tissue is collected 6 hours after the start of imbibition. The 6 hours after imbibition samples are separated into 3 collections after removal of any adhering seed coat. The 6 hours after imbibition sample is collected over the course of approximately 2 hours starting at 6 hours post imbibition. At the 6 hours after imbibition stage, not all cotyledons have become fully hydrated and germination, or radicle protrusion, has not occurred. The seedlings are washed in water to remove soil, hypocotyl axis harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON006 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seedling cotyledons tissue harvest 6 hour post-imbibition. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium. Trays are placed in an environmental chamber and grown at 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Tissue is collected 6 hours after imbibition. The 6 hours after imbibition samples are separated into 3 collections after removal of any adhering seed coat. The 6 hours after imbibition sample is collected over the course of approximately 2 hours starting at 6 hours post-imbibition. At the 6 hours after imbibition, not all cotyledons have become fully hydrated and germination or radicle protrusion, have not occurred. The seedlings are washed in water to

remove soil, cotyledon harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON007 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seed tissue harvested 25 and 35 days post-flowering. Seed pods from field grown plants are harvested 25 and 35 days after flowering and the seeds extracted from the pods. Approximately 4.4g and 19.3g of seeds are harvested from the respective seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON008 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) leaf tissue harvested from 25 and 35 days post-flowering plants. Total leaf tissue is harvested from field grown plants. Approximately 19g and 29g of leaves are harvested from the fourth node of the plant 25 and 35 days post-flowering and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON009 cDNA library is generated from soybean cutlivar C1944 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) pod and seed tissue harvested 15 days post-flowering. Pods from field grown plants are harvested 15 days post-flowering. Approximately 3g of pod tissue is harvested and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON010 cDNA library is generated from soybean cultivar C1944 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) seed tissue harvested 40 days post-flowering. Pods from field grown plants are harvested 40 days post-flowering. Pods and seeds are separated, approximately 19g of seed tissue is harvested and immediately frozen in dry-ice.

5 The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON011 cDNA library is generated from soybean cultivars Cristalina (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) and FT108 (Monsoy, Brazil) (tropical germ plasma) leaf tissue. Leaves are harvested from plants grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C . Soil is checked and watered daily to maintain even moisture conditions. Approximately 30g of leaves are harvested from the 4th node of each of the Cristalina and FT108 cultivars and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON012 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) leaf tissue. Leaves from field grown plants are harvested from the fourth node 15 days post-flowering. Approximately 12g of leaves are harvested and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until

20 RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON013 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) root and nodule tissue. Approximately, 28g

of root tissue from field grown plants is harvested 15 days post-flowering. The root system is cut from the soybean plant, washed with water to free it from the soil and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

5 The SOYMON014 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seed tissue harvested 25 and 35 days after flowering. Seed pods from field grown plants are harvested 15 days after flowering and the seeds extracted from the pods. Approximately 5g of seeds are harvested from the respective seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

10 The SOYMON015 cDNA is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seed tissue harvested 45 and 55 days post-flowering. Seed pods from field grown plants are harvested 45 and 55 days after flowering and the seeds extracted from the pods. Approximately 19g and 31g of seeds are harvested from the respective seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

15 The SOYMON016 cDNA library is generated from soybean cultivar Asgrow 3244
20 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) root tissue. Approximately, 61g and 38g of root tissue from field grown plants is harvested 25 and 35 days post- flowering is harvested. The root system is cut from the soybean plant and washed with water to free it from the soil. The tissue is placed in 14ml polystyrene tubes and immediately frozen in dry-ice. The harvested

tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON017 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) root tissue. Approximately 28g of root
5 tissue from field grown plants is harvested 45 and 55 days post-flowering. The root system is cut from the soybean plant, washed with water to free it from the soil and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

10 The SOYMON018 cDNA is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) leaf tissue harvested 45 and 55 days post-flowering. Leaves from field grown plants are harvested 45 and 55 days after flowering from the fourth node. Approximately 27g and 33g of seeds are harvested from the respective seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA
15 preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

20 The SOYMON019 cDNA library is generated from soybean cultivars Cristalina (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) and FT108 (Monsoy, Brazil) (tropical germ plasma) root tissue. Roots are harvested from plants grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Approximately 50g and 56g of roots are harvested from each of the Cristalina and FT108 cultivars and immediately frozen in dry ice. The harvested tissue is then

stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON020 cDNA is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seed tissue harvested 65 and 75 days post-flowering.

5 Seed pods from field grown plants are harvested 45 and 55 days after flowering and the seeds extracted from the pods. Approximately 14g and 31g of seeds are harvested from the respective seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

10 The SOYMON021 cDNA library is generated from Soybean Cyst Nematode-resistant soybean cultivar Hartwig (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) root tissue. Plants are grown in tissue culture at room temperature. At approximately 6 weeks post-germination, the plants are exposed to sterilized Soybean Cyst Nematode eggs. Infection is then allowed to progress for 10 days. After the 10 day infection process, the tissue is harvested. Agar from the culture medium and nematodes are removed and the root tissue is immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON022 (Lib3030) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) partially opened flower tissue.

20 Partially to fully opened flower tissue is harvested from plants grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. A total of 3g of flower tissue is harvested and immediately

frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON023 cDNA library is generated from soybean genotype BW211S Null (Tohoku University, Morioka, Japan) seed tissue harvested 15 and 40 days post-flowering. Seed pods from field grown plants are harvested 15 and 40 days post-flowering and the seeds extracted from the pods. Approximately 0.7g and 14.2g of seeds are harvested from the respective seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON024 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) internode-2 tissue harvested 18 days post-imbibition. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium. The plants are grown in a greenhouse for 18 days after the start of imbibition at ambient temperature. Soil is checked and watered daily to maintain even moisture conditions. Stem tissue is harvested 18 days after the start of imbibition. The samples are divided into hypocotyl and internodes 1 through 5. The fifth internode contains some leaf bud material. Approximately 3 g of each sample is harvested and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON025 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) leaf tissue harvested 65 days post-flowering. Leaves are harvested from the fourth node of field grown plants 65 days post-flowering. Approximately 18.4g of leaf tissue is harvested and immediately frozen in dry ice. The

harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

SOYMON026 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) root tissue harvested 65 and 75 days post-flowering.

5 Approximately 27g and 40g of root tissue from field grown plants is harvested 65 and 75 days post-flowering. The root system is cut from the soybean plant, washed with water to free it from the soil and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

10 The SOYMON027 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seed tissue harvested 25 days post-flowering. Seed pods from field grown plants are harvested 25 days post-flowering and the seeds extracted from the pods. Approximately 17g of seeds are harvested from the seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA
15 preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON028 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) drought-stressed root tissue. The plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime
20 temperature is approximately 29°C and the nighttime temperature 24°C . Soil is checked and watered daily to maintain even moisture conditions. At the R3 stage of development, water is withheld from half of the plant collection (drought stressed population). After 3 days, half of the plants from the drought stressed condition and half of the plants from the control population are

harvested. After another 3 days (6 days post drought induction) the remaining plants are harvested. A total of 27g and 40g of root tissue is harvested and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

5 The SOYMON029 cDNA library is generated from Soybean Cyst Nematode-resistant soybean cultivar PI07354 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) root tissue. Late fall to early winter greenhouse grown plants are exposed to Soybean Cyst Nematode eggs. At 10 days post-infection, the plants are uprooted, rinsed briefly and the roots frozen in liquid nitrogen. Approximately 20 grams of root tissue is harvested from the infected plants. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

10 The SOYMON030 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) flower bud tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Flower buds are removed from the plant at the pedicel. A total of 100mg of flower buds are harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

15 The SOYMON031 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) carpel and stamen tissue. Seeds are planted

at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Flower buds are removed from the plant at the pedicel. Flowers are dissected to separate petals, sepals and reproductive structures (carpels and stamens). A total of 300mg of carpel and stamen tissue are harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON032 cDNA library is prepared from the Asgrow cultivar A4922 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) rehydrated dry soybean seed meristem tissue. Surface sterilized seeds are germinated in liquid media for 24 hours. The seed axis is then excised from the barely germinating seed, placed on tissue culture media and incubated overnight at 20°C in the dark. The supportive tissue is removed from the explant prior to harvest. Approximately 570mg of tissue is harvested and frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON033 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) heat-shocked seedling tissue without cotyledons. Seeds are imbibed and germinated in vermiculite for 2 days under constant illumination. After 48 hours, the seedlings are transferred to an incubator set at 40°C under constant illumination. After 30, 60 and 180 minutes seedlings are harvested and dissected. A portion of the seedling consisting of the root, hypocotyl and apical hook is frozen in liquid

nitrogen and stored at -80°C . The seedlings after 2 days of imbibition are beginning to emerge from the vermiculite surface. The apical hooks are dark green in appearance. Total RNA and poly A⁺ RNA is prepared from equal amounts of pooled tissue. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

5 The SOYMON034 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) cold-shocked seedling tissue without cotyledons. Seeds are imbibed and germinated in vermiculite for 2 days under constant illumination. After 48 hours, the seedlings are transferred to a cold room set at 5°C under constant illumination. After 30, 60 and 180 minutes seedlings are harvested and dissected. The seedlings after 2 days of imbibition are beginning to emerge from the vermiculite surface. The apical hooks are dark green in appearance. A portion of the seedling consisting of the root, hypocotyl and apical hook is frozen in liquid nitrogen and stored at -80°C . The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

10
15
20 The SOYMON035 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seed coat tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature 24°C . Soil is checked and watered daily to maintain even moisture conditions. Seeds are harvested from mid to nearly full maturation (seed coats are not yellowing). The entire embryo proper is removed from the seed coat sample and the seed coat tissue are harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON036 cDNA library is generated from soybean cultivars PI171451, PI227687 and PI229358 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) insect challenged leaves. Plants from each of the three cultivars are grown in screenhouse conditions. The screenhouse is divided in half and one half of the screenhouse is infested with soybean
5 looper and the other half infested with velvetbean caterpillar. A single leaf is taken from each of the representative plants at 3 different time points, 11 days after infestation, 2 weeks after infestation and 5 weeks after infestation and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. Total RNA and poly A+ RNA is isolated from pooled tissue consisting of equal quantities of all 18 samples (3 genotypes X 3 sample times X 2 insect genotypes). The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON037 cDNA library is generated from soybean cultivar A3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) etiolated axis and radical tissue. Seeds are planted in moist vermiculite, wrapped and kept at room temperature in complete darkness until harvest. Etiolated axis and hypocotyl tissue is harvested at 2, 3 and 4 days post-planting. A total of 1
15 gram of each tissue type is harvested at 2, 3 and 4 days after planting and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The SOYMON038 cDNA library is generated from soybean variety Asgrow A3237
20 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) rehydrated dry seeds. Explants are prepared for transformation after germination of surface-sterilized seeds on solid tissue media. After 6days, at 28°C and 18 hours of light per day, the germinated seeds are cold shocked at 4°C for 24 hours. Meristemic tissue and part of the hypocotyl is remove and cotyledon excised. The

prepared explant is then wounded for *Agrobacterium* infection. The 2 grams of harvested tissue is frozen in liquid nitrogen and stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

The Soy51 (LIB3027) normalized seed pool cDNA library is prepared from equal
5 amounts tissue harvested from SOYMON007, SOYMON015 and SOYMON020 prepared tissue. Single stranded and double stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is mixed with the
10 biotinylated RNA in a 1:10 molar ratio and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. The non-hybridized single stranded molecules remaining after hybrid capture are converted to double stranded form and represent the primary normalized library.

The Soy52 (LIB3028) cDNA library is generated from normalized flower DNA. Single
15 stranded DNA representing approximately 1×10^6 colony forming units of SOYMON022 harvested tissue is used as the starting material for normalization. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is
20 mixed with the biotinylated RNA in a 1:10 molar ratio and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet.

The non-hybridized single stranded molecules remaining after hybrid capture are converted to double stranded form and represent the primary normalized library.

The Soy53 (LIB3039) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seedling shoot apical meristem tissue.

5 Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature 24°C. Soil is checked and watered daily to maintain even moisture conditions. Apical tissue is harvested from seedling shoot meristem tissue, 7-8 days after the start of imbibition. The apex of each seedling is dissected to include the fifth node to the apical meristem. The fifth node corresponds to the third trifoliate leaf in the very early stages of development. Stipules completely envelop the leaf primordia at this time. A total of 200mg of apical tissue is harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

10
15
20 The Soy54 (LIB3040) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) heart to torpedo stage embryo tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature 24°C. Soil is checked and watered daily to maintain even moisture conditions. Seeds are collected and embryos removed from surrounding endosperm and maternal tissues. Embryos from globular to young torpedo stages (by corresponding analogy to *Arabidopsis*) are collected

with a bias towards the middle of this spectrum. Embryos which are beginning to show asymmetric development of cotyledons are considered the upper developmental boundary for the collection and are excluded. A total of 12 mg embryo tissue is frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

Soy55 (LIB3049) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) young seed tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature 24°C. Soil is checked and watered daily to maintain even moisture conditions. Seeds are collected from very young pods (5 to 15 days after flowering). A total of 100mg of seeds are harvested and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. The RNA is purified from the stored tissue and the cDNA library is constructed as described in Example 2.

Soy56 (LIB3029) non-normalized seed pool cDNA library is prepared from equal amounts tissue harvested from SOYMON007, SOYMON015 and SOYMON020 prepared tissue. Single stranded and double stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. The non-

hybridized single stranded molecules remaining after hybrid capture are not converted to double stranded form and represent a non-normalized seed pool for comparison to Soy51 cDNA libraries.

The Soy58 (LIB3050) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) drought stressed root tissue subtracted from control root tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature 24°C. Soil is checked and watered daily to maintain even moisture conditions. At the R3 stage of the plant drought is induced by withholding water. After 3 and 6 days root tissue from both drought stressed and control (watered regularly) plants are collected and frozen in dry-ice. The harvested tissue is stored at -80°C until RNA preparation. The RNA is prepared from the stored tissue and the subtracted cDNA library is constructed as described in Example 2.

The Soy59 (LIB3051) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) endosperm tissue. Seeds are germinated on paper towels under laboratory ambient light conditions. At 8, 10 and 14 hours after imbibition, the seed coats are harvested. The endosperm consists of a very thin layer of tissue affixed to the inside of the seed coat. The seed coat and endosperm are frozen immediately after harvest in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. The RNA is prepared from the stored tissue and the cDNA library is constructed as described in Example 2.

The Soy60 (LIB3072) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) drought stressed seed plus pod subtracted

from control seed plus pod tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 26°C and the nighttime temperature 21°C and 70% relative humidity. Soil is checked and
5 watered daily to maintain even moisture conditions. At the R3 stage of the plant drought is induced by withholding water. After 3 and 6 days seeds and pods from both drought stressed and control (watered regularly) plants are collected from the fifth and sixth node and frozen in dry-ice. The harvested tissue is stored at -80°C until RNA preparation. The RNA is prepared from the stored tissue and the subtracted cDNA library is constructed as described in Example 2.

6
10
15
20
The Soy61 (LIB3073) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) jasmonic acid treated seedling subtracted from control tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in a greenhouse. The daytime temperature is approximately 29.4°C and the nighttime temperature 20°C. Soil is checked and watered daily to maintain even moisture conditions. At 9 days post planting, the plantlets are sprayed with either control buffer of 0.1% Tween-20 or jasmonic acid (Sigma J-2500, Sigma, St. Louis, Missouri U.S.A.) at 1 mg/ml in 0.1% Tween-20. Plants are sprayed until runoff and the soil and the stem is soaked with the spraying solution. At 18 hours post application of jasmonic acid, the soybean plantlets appear growth retarded. After 18 hours, 24 hours and 48 hours post
20 treatment, the cotyledons are removed and the remaining leaf and stem tissue above the soil is harvested and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. To make RNA, the three sample timepoints were combined and ground. The RNA is prepared from the stored tissue and the subtracted cDNA library is constructed as described in

Example 2. For this library's construction, the eighth fraction of the cDNA size fractionation step was used for ligation.

The Soy62 (LIB3074) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) jasmonic acid treated seedling subtracted from control tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in a greenhouse. The daytime temperature is approximately 29.4°C and the nighttime temperature 20°C. Soil is checked and watered daily to maintain even moisture conditions. At 9 days post planting, the plantlets are sprayed with either control buffer of 0.1% Tween-20 or jasmonic acid (Sigma J-2500, Sigma, St. Louis, Missouri U.S.A.) at 1 mg/ml in 0.1% Tween-20. Plants are sprayed until runoff and the soil and the stem is soaked with the spraying solution. At 18 hours post application of jasmonic acid, the soybean plantlets appear growth retarded. After 18 hours, 24 hours and 48 hours post treatment, the cotyledons are removed and the remaining leaf and stem tissue above the soil is harvested and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. To make RNA, the three sample timepoints were combined and ground. The RNA is prepared from the stored tissue and the subtracted cDNA library is constructed as described in Example 2. For this library's construction, the ninth fraction of the cDNA size fractionation step was used for ligation.

The Soy65 (LIB3107) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) drought-stressed abscission zone tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature

24°C. Soil is checked and watered daily to maintain even moisture conditions. Plants are irrigated with 15-16-17 Peter's Mix. At the R3 stage of development, drought is imposed by withholding water. At 3, 4, 5 and 6 days, tissue is harvested and wilting is not obvious until the fourth day. Abscission layers from reproductive organs are harvested by cutting less than one millimeter proximal and distal to the layer and immediately frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. The RNA is prepared from the stored tissue and the cDNA library is constructed as described in Example 2.

The Soy66 (LIB3109) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) non-drought stressed abscission zone tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Plants are irrigated with 15-16-17 Peter's Mix. At 3, 4, 5 and 6 days, control abscission layer tissue is harvested. Abscission layers from reproductive organs are harvested by cutting less than one millimeter proximal and distal to the layer and immediately frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. The RNA is prepared from the stored tissue and the cDNA library is constructed as described in Example 2.

Soy67 (LIB3065) normalized seed pool cDNA library is prepared from equal amounts of tissue harvested from SOYMON007, SOYMON015 and SOYMON020 prepared tissue. Single stranded and double stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated

into the RNA during the synthesis reaction. The single stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio) and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. Captured
5 hybrids are eluted with water.

Soy68 (LIB3052) normalized seed pool cDNA library is prepared from equal amounts tissue harvested from SOYMON007, SOYMON015 and SOYMON020 prepared tissue. Single stranded and double stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio) and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. Captured
10 hybrids are eluted with water.
15

Soy69 (LIB3053) normalized cDNA library is generated from soybean cultivars Cristalina (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) and FT108 (Monsoy, Brazil) (tropical germ plasma) normalized leaf tissue. Leaves are harvested from plants grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The
20 daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Approximately 30g of leaves are harvested from the 4th node of each of the Cristalina and FT108 cultivars and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA

preparation. The RNA is prepared from the stored tissue and the normalized cDNA library is constructed as described in Example 2.

Soy70 (LIB3055) cDNA library is generated from soybean cultivars Cristalina (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) and FT108 (Monsoy, Brazil) (tropical germ plasma) leaf tissue. Leaves are harvested from plants grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Approximately 30g of leaves are harvested from the 4th node of each of the Cristalina and FT108 cultivars and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is prepared from the stored tissue and the cDNA library is constructed as described in Example 2.

Soy71 (LIB3056) cDNA library is generated from soybean cultivars Cristalina and FT108 (tropical germ plasma) root tissue. Roots are harvested from plants grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Approximately 50g and 56g of roots are harvested from each of the Cristalina and FT108 cultivars and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. The RNA is prepared from the stored tissue and the cDNA library is constructed as described in Example 2.

Soy73 (LIB3093) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) drought stressed leaf subtracted from control tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under

12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 26°C and the nighttime temperature 21°C and 70% relative humidity. Soil is checked and watered daily to maintain even moisture conditions. At the R3 stage of the plant drought is induced by withholding water. After 3 and 6 days seeds and pods from both drought stressed and control (watered regularly) plants are collected from the fifth and sixth node and frozen in dry-ice. The harvested tissue is stored at -80°C until RNA preparation. The RNA is prepared from the stored tissue and the subtraction cDNA library is constructed as described in Example 2.

The Soy76 (Lib3106) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) jasmonic acid and arachidonic treated seedling subtracted from control tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in a greenhouse. The daytime temperature is approximately 29.4°C and the nighttime temperature 20°C. Soil is checked and watered daily to maintain even moisture conditions. At 9 days post planting, the plantlets are sprayed with either control buffer of 0.1% Tween-20 or jasmonic acid (Sigma J-2500, Sigma, St. Louis, Missouri U.S.A.) at 1 mg/ml in 0.1% Tween-20. Plants are sprayed until runoff and the soil and the stem is soaked with the spraying solution. At 18 hours post application of jasmonic acid, the soybean plantlets appear growth retarded. Arachidonic treated seedlings are sprayed with 1m/ml arachidonic acid in 0.1% Tween-20. After 18hours, 24hours and 48 hours post treatment, the cotyledons are removed and the remaining leaf and stem tissue above the soil is harvested and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. To make RNA, the three sample timepoints were combined and ground. The RNA from the arachidonic treated seedlings is isolated separately. The RNA is prepared from the stored tissue and the subtraction cDNA library is constructed as described in Example 2.

For this subtraction library, fraction 10 of the size fractionated cDNA is ligated into the pSPORT vector (Invitrogen, Carlsbad California U.S.A.) in order to capture some of the smaller transcripts characteristic of antifungal proteins.

Soy77 (LIB3108) cDNA library is generated from soybean cultivar Asgrow 3244

5 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) jasmonic acid control tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in a greenhouse. The daytime temperature is approximately 29.4°C and the nighttime temperature 20°C. Soil is checked and watered daily to maintain even moisture conditions. At 9 days post planting, the plantlets are sprayed with either control buffer of 0.1% Tween-20 or jasmonic acid (Sigma J-2500, Sigma, St. Louis, Missouri U.S.A.) at 1 mg/ml in 0.1% Tween-20. Plants are sprayed until runoff and the soil and the stem is soaked with the spraying solution. At 18 hours post application of jasmonic acid, the soybean plantlets appear growth retarded. Arachidonic treated seedlings are sprayed with 1m/ml arachidonic acid in 0.1% Tween-20. After 18 hours, 24 hours and 48 hours post treatment, the cotyledons are removed and the remaining leaf and stem tissue above the soil is harvested and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. To make RNA, the three sample timepoints were combined and ground. The RNA from the arachidonic treated seedlings is isolated separately. The RNA is prepared from the stored tissue and the subtraction cDNA library is constructed as described in Example 2. For this subtraction cDNA library,
20 fraction 10 of the size fractionated cDNA is ligated into the pSPORT vector in order to capture some of the smaller transcripts characteristic of antifungal proteins.

Example 2

The stored RNA is purified using Trizol reagent from Life Technologies (Gibco BRL, Life Technologies, Gaithersburg, Maryland U.S.A.), essentially as recommended by the manufacturer. Poly A+ RNA (mRNA) is purified using magnetic oligo dT beads essentially as
5 recommended by the manufacturer (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.).

Construction of plant cDNA libraries is well-known in the art and a number of cloning strategies exist. A number of cDNA library construction kits are commercially available. The Superscript™ Plasmid System for cDNA synthesis and Plasmid Cloning (Gibco BRL, Life
10 Technologies, Gaithersburg, Maryland U.S.A.) is used, following the conditions suggested by the manufacturer.

Normalized libraries are made using essentially the Soares procedure (Soares *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 91:9228-9232 (1994), the entirety of which is herein incorporated by reference). This approach is designed to reduce the initial 10,000-fold variation in individual
15 cDNA frequencies to achieve abundances within one order of magnitude while maintaining the overall sequence complexity of the library. In the normalization process, the prevalence of high-abundance cDNA clones decreases dramatically, clones with mid-level abundance are relatively unaffected and clones for rare transcripts are effectively increased in abundance.

Normalized libraries are prepared from single-stranded and double-stranded DNA.

20 Single-stranded and double-stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single-stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated

into the RNA during the synthesis reaction. The single-stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio) and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. The non-
5 hybridized single-stranded molecules remaining after hybrid capture are converted to double stranded form and represent the primary normalized library.

For subtraction, target cDNA is made from the drought stressed tissue total RNA using the SMART cDNA synthesis system from Clontech (Clontech Laboratories, Palo Alto, California U.S.A.). Driver first strand cDNA is covalently linked to Dynabeads following a protocol similar to that described in the Dynal literature (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The target cDNA is then heat denatured and the second strand trapped using Dynabeads oligo-dT. The target second strand cDNA is then hybridized to the driver cDNA in 400µl 2X SSPE for two rounds of hybridization at 65°C and 20 hours. After each hybridization, the hybridization solution is removed from the system and the hybridized target cDNA removed from the driver by heat denaturation in water. After hybridization, the remaining cDNA is trapped with Dynabeads oligo-dT. The trapped cDNA is then amplified as in previous PCR based libraries and the resulting cDNA ligated into the pSPORT vector (Invitrogen, Carlsbad California U.S.A.).

Example 3

20 The cDNA libraries are plated on LB agar containing the appropriate antibiotics for selection and incubated at 37° for a sufficient time to allow the growth of individual colonies. Single colonies are individually placed in each well of a 96-well microtiter plates containing LB

liquid including the selective antibiotics. The plates are incubated overnight at approximately 37°C with gentle shaking to promote growth of the cultures. The plasmid DNA is isolated from each clone using Qiaprep plasmid isolation kits, using the conditions recommended by the manufacturer (Qiagen Inc., Santa Clara, California U.S.A.).

5 Template plasmid DNA clones are used for subsequent sequencing. For sequencing, the ABI PRISM dRhodamine Terminator Cycle Sequencing Ready Reaction Kit with AmpliTaq® DNA Polymerase, FS, is used (PE Applied Biosystems, Foster City, California U.S.A.).

Example 4

10 Nucleic acid sequences that encode for the following tocopherol synthesis pathway enzymes: deoxyarabiono-heptulosonate-P-synthase; putative deoxyarabiono-heptulosonate-P-synthase; dehydroquinase synthase; dehydroquinase dehydratase; putative dehydroquinase dehydratase; shikimate dehydrogenase; shikimate kinase; enolpyruvylshikimate-P-synthase; chorismate synthase; chorismate mutase; tyrosine transaminase; putative tyrosine transaminase; transaminase A; putative transaminase A; 4-hydroxyphenylpyruvate dioxygenase; homogentisic acid dioxygenase; and geranylgeranylpyrophosphate synthase are identified from the Monsanto EST PhytoSeq database using TBLASTN (default values)(TBLASTN compares a protein query against the six reading frames of a nucleic acid sequence). Matches found with BLAST P values equal or less than 0.001 (probability) or BLAST Score of equal or greater than 90 are classified as hits. If the program used to determine the hit is HMMSW then the score refers to HMMSW
20 score.

 In addition, the GenBank database is searched with BLASTN and BLASTX (default values) using ESTs as queries. EST that pass the hit probability threshold of $10e^{-8}$ for the

following enzymes are combined with the hits generated by using TBLASTN (described above) and classified by enzyme (see Table A below).

A cluster refers to a set of overlapping clones in the PhytoSeq database. Such an overlapping relationship among clones is designated as a "cluster" when BLAST scores from pairwise sequence comparisons of the member clones meets a predetermined minimum value or product score of 50 or more (Product Score = (BLAST SCORE x Percentage Identity)/(5 x minimum [length (Seq1), length (Seq2)])).

Since clusters are formed on the basis of single-linkage relationships, it is possible for two non-overlapping clones to be members of the same cluster if, for instance, they both overlap a third clone with at least the predetermined minimum BLAST score (stringency). A cluster ID is arbitrarily assigned to all of those clones which belong to the same cluster at a given stringency and a particular clone will belong to only one cluster at a given stringency. If a cluster contains only a single clone (a "singleton"), then the cluster ID number will be negative, with an absolute value equal to the clone ID number of its single member. Clones grouped in a cluster in most cases represent a contiguous sequence.

TABLE A*

deoxyarabiono-heptulosonate-P-synthase-maize								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
1	-700223776	700223776H1	SATMON011	g2398680	BLASTN	388	1e-51	77
2	-700260027	700260027H1	SATMON017	g169475	BLASTX	112	1e-10	75
3	-700356188	700356188H1	SATMON024	g2398679	BLASTX	93	1e-13	78
4	-700430072	700430072H1	SATMONN01	g2398679	BLASTX	180	1e-17	85
5	1228	700623827H1	SATMON034	g416252	BLASTN	1030	1e-105	87
6	1228	700452503H1	SATMON028	g416252	BLASTN	1141	1e-88	87
7	1228	700551557H1	SATMON022	g416252	BLASTN	547	1e-83	88
8	1228	700571345H1	SATMON030	g416252	BLASTN	712	1e-83	83
9	1228	700452527H1	SATMON028	g416252	BLASTN	908	1e-83	88
10	1228	700050505H1	SATMON003	g416252	BLASTN	1080	1e-83	90
11	1228	700551749H1	SATMON022	g416252	BLASTN	1029	1e-79	90
12	1228	700569172H1	SATMON030	g416252	BLASTN	993	1e-76	84
13	1228	700613721H1	SATMON033	g416252	BLASTN	491	1e-75	85
14	1228	700160395H1	SATMON012	g416252	BLASTN	969	1e-74	91
15	1228	701163236H1	SATMONN04	g416252	BLASTN	721	1e-73	84
16	1228	700267876H1	SATMON017	g416252	BLASTN	555	1e-65	86
17	1228	700096649H1	SATMON008	g169474	BLASTN	889	1e-65	79
18	1228	700346229H1	SATMON021	g166687	BLASTN	880	1e-64	78
19	1228	700259208H1	SATMON017	g169474	BLASTN	754	1e-63	76
20	1228	700454345H1	SATMON029	g416252	BLASTN	828	1e-62	86
21	1228	700151789H1	SATMON007	g416252	BLASTN	830	1e-62	87
22	1228	700803047H1	SATMON036	g169474	BLASTN	658	1e-61	77
23	1228	700049005H1	SATMON003	g416252	BLASTN	818	1e-61	89
24	1228	700617057H1	SATMON033	g170224	BLASTN	737	1e-60	79
25	1228	700204532H1	SATMON003	g2398680	BLASTN	486	1e-58	75
26	1228	700195344H1	SATMON014	g416252	BLASTN	774	1e-58	90
27	1228	700041845H1	SATMON004	g170224	BLASTN	806	1e-58	78
28	1228	700452030H1	SATMON028	g416252	BLASTN	574	1e-57	87
29	1228	700093451H1	SATMON008	g169474	BLASTN	511	1e-56	74
30	1228	700048096H1	SATMON003	g169474	BLASTN	774	1e-55	75
31	1228	700424054H1	SATMONN01	g170224	BLASTN	444	1e-54	81
32	1228	700448927H1	SATMON028	g416252	BLASTN	736	1e-54	83
33	1228	700579786H1	SATMON031	g169474	BLASTN	737	1e-52	74
34	1228	700168659H1	SATMON013	g169474	BLASTN	727	1e-51	79
35	1228	700150987H1	SATMON007	g166687	BLASTN	711	1e-50	81
36	1228	700022040H1	SATMON001	g294284	BLASTN	713	1e-50	80
37	1228	700449746H2	SATMON028	g416252	BLASTN	680	1e-49	90
38	1228	700166534H1	SATMON013	g169474	BLASTN	696	1e-49	76
39	1228	700257864H1	SATMON017	g169474	BLASTN	693	1e-48	78
40	1228	700452534H1	SATMON028	g416252	BLASTN	636	1e-45	85
41	1228	700042596H1	SATMON004	g416252	BLASTN	640	1e-45	84
42	1228	700151812H1	SATMON007	g416252	BLASTN	643	1e-45	89
43	1228	700421687H1	SATMONN01	g170224	BLASTN	646	1e-45	76
44	1228	700052344H1	SATMON003	g170224	BLASTN	657	1e-45	80
45	1228	701178585H1	SATMONN05	g166687	BLASTN	427	1e-44	80
46	1228	700239365H1	SATMON010	g170224	BLASTN	628	1e-43	80
47	1228	700153542H1	SATMON007	g169474	BLASTN	617	1e-42	76
48	1228	700380557H1	SATMON021	g416252	BLASTN	500	1e-41	90

49	1228	700570454H1	SATMON030	g170224	BLASTN	614	1e-41	79
50	1228	700264817H1	SATMON017	g169474	BLASTN	410	1e-39	75
51	1228	700153524H1	SATMON007	g169474	BLASTN	361	1e-37	77
52	1228	700618945H1	SATMON034	g416252	BLASTN	458	1e-36	79
53	1228	700193060H1	SATMON014	g1245452	BLASTN	547	1e-36	74
54	1228	700047557H1	SATMON003	g416252	BLASTN	511	1e-34	83
55	1228	700341009H1	SATMON020	g169474	BLASTN	324	1e-30	66
56	1228	700048589H1	SATMON003	g166689	BLASTN	409	1e-30	77
57	1228	700334956H1	SATMON019	g2398681	BLASTX	183	1e-26	85
58	1228	700027629H1	SATMON003	g169475	BLASTX	189	1e-19	87
59	29578	700219020H1	SATMON011	g2398678	BLASTN	694	1e-49	72
60	3007	700153267H1	SATMON007	g2398678	BLASTN	587	1e-40	71
61	3007	700352639H1	SATMON024	g2398678	BLASTN	506	1e-33	70
62	3007	700259481H1	SATMON017	g2398678	BLASTN	268	1e-13	66
63	31415	700219261H1	SATMON011	g166689	BLASTN	501	1e-32	77
64	32242	700090119H1	SATMON011	g2398680	BLASTN	803	1e-58	76
65	3227	700268010H1	SATMON017	g2398680	BLASTN	773	1e-55	73
66	3227	700450763H1	SATMON028	g166689	BLASTN	713	1e-50	73
67	3227	700241730H1	SATMON010	g2398680	BLASTN	686	1e-48	73
68	3227	700071633H1	SATMON007	g166689	BLASTN	628	1e-43	71
69	3227	700267002H1	SATMON017	g169474	BLASTN	446	1e-26	74
70	3227	700452543H1	SATMON028	g2546987	BLASTN	281	1e-25	75
71	5023	700378721H1	SATMON020	g2546988	BLASTX	183	1e-18	94
72	-L1487205	LIB148-064-Q1-E1-F6	LIB148	g2398679	BLASTX	296	1e-53	59
73	-L30622733	LIB3062-014-Q1-K1-D2	LIB3062	g2546987	BLASTN	484	1e-31	68
74	-L30622734	LIB3062-014-Q1-K1-D6	LIB3062	g416252	BLASTN	479	1e-29	68
75	-L30661773	LIB3066-011-Q1-K1-D6	LIB3066	g170224	BLASTN	747	1e-52	70
76	-L30664853	LIB3066-031-Q1-K1-F6	LIB3066	g169474	BLASTN	1139	1e-86	72
77	-L30685059	LIB3068-008-Q1-K1-A11	LIB3068	g166687	BLASTN	450	1e-26	64
78	-L30691358	LIB3069-002-Q1-K1-D6	LIB3069	g2398680	BLASTN	466	1e-28	60
79	1228	LIB3062-032-Q1-K1-F11	LIB3062	g416252	BLASTN	1508	1e-119	88
80	1228	LIB143-004-Q1-E1-H9	LIB143	g169474	BLASTN	1221	1e-92	78
81	1228	LIB3069-027-Q1-K1-E11	LIB3069	g169474	BLASTN	746	1e-91	77
82	1228	LIB143-045-Q1-E1-F10	LIB143	g170224	BLASTN	1056	1e-79	77
83	1228	LIB3069-001-Q1-K1-E2	LIB3069	g416252	BLASTN	598	1e-78	87
84	1228	LIB3068-006-Q1-K1-F10	LIB3068	g169474	BLASTN	1048	1e-78	74
85	1228	LIB3062-008-Q1-K1-A5	LIB3062	g169474	BLASTN	806	1e-75	79
86	1228	LIB148-045-Q1-E1-G7	LIB148	g2398678	BLASTN	735	1e-60	72
87	1228	LIB3068-034-Q1-K1-E6	LIB3068	g170224	BLASTN	819	1e-59	78
88	1228	LIB3061-040-Q1-K1-B8	LIB3061	g170224	BLASTN	771	1e-54	80
89	1228	LIB143-024-Q1-E1-B4	LIB143	g169474	BLASTN	736	1e-50	73
90	1228	LIB3068-032-Q1-K1-D10	LIB3068	g169474	BLASTN	598	1e-39	72
91	24030	LIB3066-047-Q1-K1-A9	LIB3066	g170225	BLASTX	121	1e-30	57
92	29578	LIB3066-011-Q1-K1-D4	LIB3066	g2546987	BLASTN	1212	1e-92	75
93	29578	LIB148-058-Q1-E1-F2	LIB148	g169474	BLASTN	680	1e-47	72
94	31415	LIB148-009-Q1-E1-E8	LIB148	g2398680	BLASTN	492	1e-29	74
95	32242	LIB148-018-Q1-E1-B2	LIB148	g2398680	BLASTN	1280	1e-97	76
96	32242	LIB3067-058-Q1-K1-C8	LIB3067	g2398680	BLASTN	1079	1e-81	76
97	5023	LIB3079-006-Q1-K1-D10	LIB3079	g2398681	BLASTX	143	1e-28	88

putative deoxyarabiono-heptulosonate-P-synthase-maize								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
98	-701178041	701178041H1	SATMONN05	g1742787	BLASTX	121	1e-9	49
99	13211	700267210H1	SATMON017	g1742787	BLASTX	72	1e-9	59

deoxyarabiono-heptulosonate-P-synthase-soybean								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
100	-700750583	700750583H1	SOYMON014	g169475	BLASTX	149	1e-13	78
101	-700756739	700756739H1	SOYMON014	g410315	BLASTX	170	1e-21	55
102	-700897290	700897290H1	SOYMON027	g1245452	BLASTN	1047	1e-78	93
103	-700953858	700953858H1	SOYMON022	g2398678	BLASTN	673	1e-47	74
104	-700958333	700958333H1	SOYMON022	g2398678	BLASTN	531	1e-35	77
105	-701212422	701212422H1	SOYMON035	g410487	BLASTN	527	1e-35	77
106	11948	701214211H1	SOYMON035	g2398678	BLASTN	860	1e-62	79
107	11948	700941217H1	SOYMON024	g2398678	BLASTN	843	1e-61	79
108	11948	700749762H1	SOYMON013	g2398678	BLASTN	659	1e-58	79
109	11948	701015341H1	SOYMON019	g169474	BLASTN	758	1e-54	78
110	11948	700787714H2	SOYMON011	g169474	BLASTN	613	1e-42	80
111	11948	700963862H1	SOYMON022	g2398678	BLASTN	581	1e-39	78
112	11948	701144405H1	SOYMON031	g2546987	BLASTN	266	1e-27	71
113	11948	700897376H1	SOYMON027	g2398679	BLASTX	136	1e-21	85
114	12144	700564714H1	SOYMON002	g170225	BLASTX	78	1e-14	54
115	12144	701036988H1	SOYMON029	g170225	BLASTX	64	1e-10	55
116	12144	701142430H1	SOYMON038	g170225	BLASTX	66	1e-10	52
117	18499	700746365H1	SOYMON013	g1245452	BLASTN	726	1e-62	85
118	18499	700565543H1	SOYMON002	g1245452	BLASTN	700	1e-49	88
119	19009	700953162H1	SOYMON022	g166689	BLASTN	679	1e-47	74
120	19009	700681944H1	SOYMON008	g2398680	BLASTN	613	1e-42	77
121	19576	701097076H1	SOYMON028	g1245452	BLASTN	1156	1e-87	92
122	19576	700669658H1	SOYMON006	g1245452	BLASTN	895	1e-65	90
123	19576	700656892H1	SOYMON004	g410487	BLASTN	410	1e-48	79
124	5102	700901033H1	SOYMON027	g2398678	BLASTN	894	1e-65	82
125	5102	700901290H1	SOYMON027	g2398678	BLASTN	837	1e-60	82
126	5102	701051386H1	SOYMON032	g170224	BLASTN	819	1e-59	81
127	5102	700755856H1	SOYMON014	g170224	BLASTN	782	1e-56	82
128	5102	701145291H1	SOYMON031	g170224	BLASTN	487	1e-48	80
129	5234	700565904H1	SOYMON002	g2398678	BLASTN	584	1e-77	84
130	5234	701138971H1	SOYMON038	g2398678	BLASTN	946	1e-70	84
131	5234	700725760H1	SOYMON009	g2398678	BLASTN	908	1e-66	82
132	5234	701097139H1	SOYMON028	g2398678	BLASTN	772	1e-55	79
133	5234	700952432H1	SOYMON022	g2398680	BLASTN	689	1e-48	81
134	5234	700996417H1	SOYMON018	g410487	BLASTN	468	1e-40	81
135	5699	701040351H1	SOYMON029	g166690	BLASTX	203	1e-21	69
136	5699	700847113H1	SOYMON021	g2398679	BLASTX	147	1e-13	72
137	5699	700967767H1	SOYMON033	g166690	BLASTX	149	1e-13	63
138	5699	700841638H1	SOYMON020	g166690	BLASTX	141	1e-12	62
139	5699	700891749H1	SOYMON024	g410486	BLASTX	127	1e-10	76
140	5699	700990984H1	SOYMON011	g2398679	BLASTX	127	1e-10	67
141	5699	700740310H1	SOYMON012	g410486	BLASTX	127	1e-10	76
142	5699	700834916H1	SOYMON019	g294285	BLASTX	117	1e-9	77
143	6819	700652910H1	SOYMON003	g1245452	BLASTN	1408	1e-109	89

144	6819	700761928H1	SOYMON015	g1245452	BLASTN	924	1e-68	89
145	6935	700987126H1	SOYMON009	g2398678	BLASTN	667	1e-46	72
146	6935	700734128H1	SOYMON010	g169474	BLASTN	533	1e-35	72

putative deoxyarabiono-heptulosonate-P-synthase -soybean

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
147	-700891658	700891658H1	SOYMON024	g1742787	BLASTX	119	1e-9	40
148	-701148391	701148391H1	SOYMON031	g1742787	BLASTX	109	1e-9	43
149	4075	700992239H1	SOYMON011	g1742787	BLASTX	66	1e-9	40
150	4075	700686128H1	SOYMON008	g1742787	BLASTX	62	1e-8	39
151	19576	LIB3029-012-Q1-B1-B5	LIB3029	g2546987	BLASTN	1280	1e-97	80
152	5699	LIB3052-011-Q1-N1-E8	LIB3052	g166690	BLASTX	187	1e-39	56

dehydroquinase synthase-maize

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
153	-700257536	700257536H1	SATMON017	g309862	BLASTX	102	1e-21	69
154	28069	700203301H1	SATMON003	g1789791	BLASTX	140	1e-16	50
155	7410	700222526H1	SATMON011	g1619336	BLASTX	149	1e-21	56
156	7410	700347409H1	SATMON023	g40968	BLASTX	83	1e-16	55
157	28069	LIB189-001-Q1-E1-D4	LIB189	g1789791	BLASTX	281	1e-48	57

putative dehydroquinase dehydratase-maize

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
158	-700237972	700237972H1	SATMON010	g535771	BLASTX	136	1e-18	69
159	11022	700155850H1	SATMON007	g535771	BLASTX	247	1e-27	65

dehydroquinase dehydratase-soybean

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
160	4639	700834936H1	SOYMON019	g535771	BLASTX	150	1e-20	55

Shikimate dehydrogenase-maize

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
158	-700237972	700237972H1	SATMON010	g535771	BLASTX	136	1e-18	69
159	11022	700155850H1	SATMON007	g535771	BLASTX	247	1e-27	65

Shikimate dehydrogenase-soybean

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
160	4639	700834936H1	SOYMON019	g535771	BLASTX	150	1e-20	55

Shikimate kinase-maize								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
161	-700050913	700050913H1	SATMON003	g19348	BLASTN	403	1e-28	69
162	-700104390	700104390H1	SATMON010	g19348	BLASTN	446	1e-26	64
163	-700452495	700452495H1	SATMON028	g19349	BLASTX	81	1e-13	57
164	-700619865	700619865H1	SATMON034	g19349	BLASTX	142	1e-12	65
165	15996	700030278H1	SATMON003	g19349	BLASTX	219	1e-33	66
166	15996	700257047H1	SATMON017	g19348	BLASTN	399	1e-32	66
167	15996	700237902H1	SATMON010	g19348	BLASTN	472	1e-28	64
168	15996	700155641H1	SATMON007	g19348	BLASTN	438	1e-27	66
169	15996	700224589H1	SATMON011	g19348	BLASTN	447	1e-27	64
170	18563	700205659H1	SATMON003	g19348	BLASTN	443	1e-39	67
171	18563	700243143H1	SATMON010	g19349	BLASTX	274	1e-31	64
172	18563	700264692H1	SATMON017	g19348	BLASTN	501	1e-31	63
173	18563	700106054H1	SATMON010	g19348	BLASTN	280	1e-27	63
174	18563	700026972H1	SATMON003	g19349	BLASTX	159	1e-26	66
175	18563	700160974H1	SATMON012	g19349	BLASTX	167	1e-17	57
176	6303	700088964H1	SATMON011	g19348	BLASTN	470	1e-28	62
177	6303	700572756H1	SATMON030	g19349	BLASTX	113	1e-18	60
178	15635	LIB36-001-Q1-E1-F1	LIB36	g19349	BLASTX	149	1e-28	31
179	18563	LIB3066-029-Q1-K1-G8	LIB3066	g19348	BLASTN	870	1e-63	67

Shikimate kinase-soybean								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
180	-700568344	700568344H1	SOYMON002	g19349	BLASTX	126	1e-15	42
181	-700792015	700792015H1	SOYMON011	g19348	BLASTN	652	1e-45	72
182	18190	700977239H1	SOYMON009	g19349	BLASTX	97	1e-10	41
183	18190	LIB3055-003-Q1-N1-D12	LIB3055	g19349	BLASTX	139	1e-28	36

Enolpyruvylshikimate-P-synthase-soybean								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
184	-700831419	700831419H1	SOYMON019	g169190	BLASTN	453	1e-50	80
185	-700845353	700845353H1	SOYMON021	g170373	BLASTN	629	1e-43	76
186	-700891187	700891187H1	SOYMON024	g170373	BLASTN	620	1e-42	74
187	-700976722	700976722H1	SOYMON009	g170373	BLASTN	774	1e-55	75
188	-700997285	700997285H1	SOYMON018	g170374	BLASTX	124	1e-11	86
189	-701048471	701048471H1	SOYMON032	g170228	BLASTN	886	1e-64	82
190	-701206839	701206839H1	SOYMON035	g17815	BLASTX	154	1e-14	88
191	17068	700942983H1	SOYMON024	g169190	BLASTN	571	1e-58	82
192	17068	701006194H1	SOYMON019	g169190	BLASTN	349	1e-53	79
193	18050	700906275H1	SOYMON022	g169190	BLASTN	868	1e-63	81
194	18050	701134508H1	SOYMON038	g169190	BLASTN	457	1e-60	80
195	3411	700556807H1	SOYMON001	g169190	BLASTN	568	1e-77	83
196	3411	700565035H1	SOYMON002	g169190	BLASTN	913	1e-67	79
197	3411	701008536H1	SOYMON019	g169190	BLASTN	622	1e-56	80
198	3411	701107917H1	SOYMON036	g170228	BLASTN	498	1e-32	84

Chorismate synthase-maize

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
199	-700104711	700104711H1	SATMON010	g976374	BLASTN	490	1e-30	70
200	10770	700092595H1	SATMON008	g410484	BLASTX	207	1e-21	70
201	10770	700088420H1	SATMON011	g410484	BLASTX	191	1e-19	68
202	10770	700333085H1	SATMON019	g410484	BLASTX	104	1e-13	67
203	2026	700282007H1	SATMON022	g410481	BLASTN	884	1e-64	77
204	2026	700077339H1	SATMON007	g410481	BLASTN	612	1e-59	75
205	2026	700571731H1	SATMON030	g410481	BLASTN	670	1e-54	77
206	2026	700348949H1	SATMON023	g410481	BLASTN	463	1e-53	77
207	2026	700090790H1	SATMON011	g18255	BLASTN	694	1e-49	71
208	2026	700236685H1	SATMON010	g18255	BLASTN	704	1e-49	77
209	2026	700166396H1	SATMON013	g410481	BLASTN	674	1e-47	76
210	2026	700466807H1	SATMON025	g18255	BLASTN	452	1e-43	71
211	2026	700335877H1	SATMON019	g410481	BLASTN	532	1e-35	74
212	4211	700259039H1	SATMON017	g18256	BLASTX	167	1e-18	60
213	4211	700457104H1	SATMON029	g410484	BLASTX	186	1e-18	75
214	4211	700153433H1	SATMON007	g410482	BLASTX	114	1e-14	68
215	4211	700073550H1	SATMON007	g18256	BLASTX	147	1e-13	61
216	4211	700224255H1	SATMON011	g18255	BLASTN	290	1e-13	67
217	4211	700440561H1	SATMON026	g410482	BLASTX	115	1e-8	66
218	9237	700105367H1	SATMON010	g410483	BLASTN	773	1e-55	73
219	9237	700337228H1	SATMON020	g18255	BLASTN	776	1e-55	73
220	9237	700159709H1	SATMON012	g18255	BLASTN	742	1e-53	76
221	9237	700242181H1	SATMON010	g410483	BLASTN	588	1e-40	71
222	9237	700168082H1	SATMON013	g18255	BLASTN	521	1e-34	73
223	9237	700169319H1	SATMON013	g18255	BLASTN	465	1e-29	78
224	9237	700172250H1	SATMON013	g18255	BLASTN	308	1e-15	71
225	9237	700584289H1	SATMON031	g410484	BLASTX	93	1e-8	69
226	-L1434254	LIB143-041-Q1-E1-F6	LIB143	g18255	BLASTN	541	1e-34	68
227	-L30781785	LIB3078-015-Q1-K1-E5	LIB3078	g410481	BLASTN	766	1e-55	67
228	2026	LIB3066-009-Q1-K1-C12	LIB3066	g410481	BLASTN	1124	1e-84	75
229	2026	LIB3078-012-Q1-K1-C8	LIB3078	g410481	BLASTN	992	1e-73	78
230	2026	LIB84-014-Q1-E1-D4	LIB84	g410481	BLASTN	517	1e-32	77
231	9237	LIB3067-014-Q1-K1-C5	LIB3067	g410484	BLASTX	219	1e-39	74

Chorismate synthase-soybean

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
232	-700829731	700829731H1	SOYMON019	g410482	BLASTX	163	1e-15	81
233	-700867002	700867002H1	SOYMON016	g18255	BLASTN	871	1e-63	82
234	-700941055	700941055H1	SOYMON024	g18257	BLASTN	621	1e-42	73
235	-700993596	700993596H1	SOYMON011	g18255	BLASTN	648	1e-56	80
236	-701107074	701107074H1	SOYMON036	g410482	BLASTX	119	1e-9	77
237	-701215158	701215158H1	SOYMON035	g18255	BLASTN	404	1e-33	82
238	11113	700792218H1	SOYMON011	g18257	BLASTN	549	1e-36	71
239	11113	701037327H1	SOYMON029	g18257	BLASTN	535	1e-35	70
240	20587	701042739H1	SOYMON029	g18257	BLASTN	537	1e-35	70
241	20587	700565645H1	SOYMON002	g18257	BLASTN	427	1e-31	73
242	24472	701053135H1	SOYMON032	g18255	BLASTN	927	1e-68	78
243	24472	700875233H1	SOYMON018	g410481	BLASTN	843	1e-61	81
244	6572	700652322H1	SOYMON003	g18255	BLASTN	722	1e-65	82

245	6572	701107063H1	SOYMON036	g18255	BLASTN	451	1e-58	81
246	6572	701139518H1	SOYMON038	g18255	BLASTN	682	1e-48	80
247	6572	700653111H1	SOYMON003	g410481	BLASTN	585	1e-39	81
248	6572	701008289H1	SOYMON019	g410481	BLASTN	545	1e-36	76
249	6572	701124777H1	SOYMON037	g18255	BLASTN	446	1e-27	80
250	6572	700556802H1	SOYMON001	g410482	BLASTX	161	1e-19	80
251	6572	700834126H1	SOYMON019	g410482	BLASTX	131	1e-16	77
252	6572	700645571H1	SOYMON009	g410482	BLASTX	172	1e-16	87
253	6572	700834378H1	SOYMON019	g18255	BLASTN	154	1e-15	77
254	6572	700990811H1	SOYMON011	g410484	BLASTX	107	1e-9	69
255	6572	LIB3030-002-Q1-B1-F12	LIB3030	g18255	BLASTN	854	1e-62	77

Chorismate mutase-maize

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
256	-700050713	700050713H1	SATMON003	g429153	BLASTX	159	1e-23	81
257	-700239884	700239884H1	SATMON010	g429153	BLASTX	146	1e-13	40
258	-700573382	700573382H1	SATMON030	g429152	BLASTN	349	1e-18	72
259	25556	700343477H1	SATMON021	g2352928	BLASTN	502	1e-35	70
260	25556	700194568H1	SATMON014	g429153	BLASTX	209	1e-21	65
261	25556	700196845H1	SATMON014	g429153	BLASTX	97	1e-10	46
262	32994	700089092H1	SATMON011	g429153	BLASTX	110	1e-15	48
263	32994	700203014H1	SATMON003	g429153	BLASTX	117	1e-9	43
264	3773	700048888H1	SATMON003	g429153	BLASTX	91	1e-24	72
265	3773	700090144H1	SATMON011	g429153	BLASTX	182	1e-18	55
266	3773	700221335H1	SATMON011	g429153	BLASTX	109	1e-10	49
267	8783	700574324H2	SATMON030	g429152	BLASTN	290	1e-13	73
268	8783	700164106H1	SATMON013	g429153	BLASTX	87	1e-9	58
269	25556	LIB3062-059-Q1-K1-H12	LIB3062	g2352928	BLASTN	502	1e-33	70
270	25556	LIB3062-023-Q1-K1-F12	LIB3062	g2352928	BLASTN	493	1e-32	70
271	25556	LIB3059-001-Q1-K2-E4	LIB3059	g2352928	BLASTN	443	1e-28	68
272	25556	LIB3069-042-Q1-K1-E10	LIB3069	g429152	BLASTN	260	1e-10	71
273	32994	LIB189-013-Q1-E1-G8	LIB189	g429153	BLASTX	150	1e-36	45
274	3773	LIB3062-011-Q1-K1-E11	LIB3062	g429152	BLASTN	621	1e-41	73
275	3773	LIB3061-006-Q1-K1-B5	LIB3061	g429152	BLASTN	408	1e-40	72
276	3773	LIB3061-035-Q1-K1-B12	LIB3061	g429152	BLASTN	319	1e-15	77
277	8783	LIB3059-017-Q1-K1-C2	LIB3059	g429152	BLASTN	357	1e-18	66

Chorismate mutase-soybean

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
278	-700649675	700649675H1	SOYMON003	g429153	BLASTX	207	1e-21	62
279	24797	701123012H1	SOYMON037	g3021541	BLASTN	525	1e-37	75
280	24797	701149634H1	SOYMON031	g3021541	BLASTN	520	1e-36	75
281	7212	700646325H1	SOYMON013	g429153	BLASTX	116	1e-8	65
282	-GM22414	LIB3030-009-Q1-B1-B5	LIB3030	g429153	BLASTX	134	1e-39	59
283	-GM29291	LIB3050-017-Q1-E1-E9	LIB3050	g2352930	BLASTN	473	1e-30	66
284	-GM30547	LIB3050-004-Q1-E1-G9	LIB3050	g429153	BLASTX	153	1e-29	64

tyrosine transaminase-maize								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
285	16305	700337451H1	SATMON020	g408894	BLASTX	134	1e-11	33
286	16305	700340103H1	SATMON020	g408894	BLASTX	93	1e-10	39

putative tyrosine transaminase-maize								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
287	14653	700220061H1	SATMON011	g2842484	BLASTX	349	1e-41	70
288	22902	700106817H1	SATMON010	g2842484	BLASTX	331	1e-38	58
289	22902	701181789H1	SATMONN06	g2842484	BLASTX	278	1e-31	62
290	6658	700442825H1	SATMON026	g2842484	BLASTX	209	1e-26	62
291	6658	700152030H1	SATMON007	g2842484	BLASTX	128	1e-18	53
292	6658	LIB3066-020-Q1-K1-F1	LIB3066	g2842484	BLASTX	348	1e-64	57

putative tyrosine transaminase-soybean								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
293	-700848909	700848909H1	SOYMON021	g2842484	BLASTX	281	1e-31	62
294	-700900410	700900410H1	SOYMON027	g2842484	BLASTX	119	1e-11	42
295	17700	700905146H1	SOYMON022	g2842484	BLASTX	315	1e-36	67
296	2201	700730931H1	SOYMON009	g2842484	BLASTX	174	1e-17	43
297	2201	700752627H1	SOYMON014	g2842484	BLASTX	102	1e-12	41
298	94	700658292H1	SOYMON004	g2842484	BLASTX	100	1e-18	53
299	6064	LIB3056-002-Q1-B1-A8	LIB3056	g2842484	BLASTX	124	1e-25	34
300	94	LIB3051-101-Q1-K1-H3	LIB3051	g2842484	BLASTX	205	1e-37	44

Transaminase A-maize								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
301	-700028003	700028003H1	SATMON003	g63066	BLASTX	125	1e-10	79
302	-700072842	700072842H1	SATMON007	g1001121	BLASTX	259	1e-28	50
303	-700194011	700194011H1	SATMON014	g435456	BLASTN	324	1e-18	73
304	-700196486	700196486H1	SATMON014	g20599	BLASTX	68	1e-10	74
305	-700331820	700331820H1	SATMON019	g20600	BLASTN	1192	1e-90	90
306	-700454550	700454550H1	SATMON029	g435458	BLASTN	198	1e-20	82
307	-700454567	700454567H1	SATMON029	g435458	BLASTN	333	1e-24	82
308	-700454642	700454642H1	SATMON029	g435458	BLASTN	269	1e-23	89
309	-700454849	700454849H1	SATMON029	g435458	BLASTN	318	1e-26	87
310	-700468560	700468560H1	SATMON025	g3328816	BLASTX	139	1e-19	58
311	-700476413	700476413H1	SATMON025	g2984217	BLASTX	156	1e-22	52
312	-700615109	700615109H1	SATMON033	g20598	BLASTN	256	1e-17	81
313	-701161385	701161385H1	SATMONN04	g435458	BLASTN	523	1e-45	80
314	10165	700341126H1	SATMON020	g20596	BLASTN	743	1e-71	91
315	10165	700160220H1	SATMON012	g20596	BLASTN	769	1e-55	92
316	10165	700158802H1	SATMON012	g20596	BLASTN	617	1e-42	94
317	10192	700204319H1	SATMON003	g2984217	BLASTX	148	1e-13	55
318	10329	700095671H1	SATMON008	g20600	BLASTN	816	1e-59	87
319	10329	700214146H1	SATMON016	g20596	BLASTN	610	1e-42	88

320	10329	700041823H1
321	10329	700094321H1
322	1148	700089060H1
323	1148	700044414H1
324	1148	700101429H1
325	1148	700221366H1
326	1148	700101604H1
327	1148	700041864H1
328	1148	700157048H1
329	1148	700581463H1
330	1148	700579938H1
331	1148	700432477H1
332	1148	700154706H1
333	1148	700043761H1
334	1148	700423679H1
335	1148	700424076H1
336	1148	701166426H1
337	16872	700211160H1
338	16872	700043705H1
339	16872	700208983H1
340	16872	700101375H1
341	17829	700194282H1
342	17829	700581970H1
343	18047	700206971H1
344	19241	700472363H1
345	19241	700472263H1
346	19241	700806145H1
347	319	700076939H1
348	319	700349974H1
349	319	700235923H1
350	319	700206180H1
351	319	700476547H1
352	319	700258893H1
353	319	700612236H1
354	319	700806537H1
355	319	700450338H1
356	319	700806243H1
357	319	700263732H1
358	319	700806094H1
359	319	700152610H1
360	319	700614581H1
361	319	700349161H1
362	319	700805964H1
363	319	700450544H1
364	319	700618252H1
365	319	700615189H1
366	319	700264196H1
367	4431	700211615H1
368	541	700073508H1
369	541	700098793H1
370	541	700101956H1
371	541	700100132H1
372	541	700799335H1
373	541	700446909H1

SATMON004	g20596	BLASTN	615	1e-42	78
SATMON008	g20596	BLASTN	559	1e-40	88
SATMON011	g633094	BLASTN	1397	1e-107	92
SATMON004	g633094	BLASTN	1272	1e-97	92
SATMON009	g633094	BLASTN	1221	1e-92	91
SATMON011	g633094	BLASTN	1205	1e-91	94
SATMON009	g633094	BLASTN	1167	1e-88	89
SATMON004	g633094	BLASTN	1159	1e-87	91
SATMON012	g633094	BLASTN	1121	1e-84	93
SATMON031	g633094	BLASTN	1124	1e-84	90
SATMON031	g633094	BLASTN	661	1e-83	91
SATMONN01	g633094	BLASTN	1050	1e-78	90
SATMON007	g633094	BLASTN	997	1e-74	90
SATMON004	g633094	BLASTN	905	1e-66	92
SATMONN01	g633094	BLASTN	555	1e-54	81
SATMONN01	g633094	BLASTN	228	1e-19	87
SATMONN04	g633094	BLASTN	221	1e-16	79
SATMON016	g633094	BLASTN	482	1e-56	88
SATMON004	g633094	BLASTN	293	1e-42	85
SATMON016	g633094	BLASTN	250	1e-15	84
SATMON009	g633094	BLASTN	154	1e-11	87
SATMON014	g1001309	BLASTX	107	1e-11	53
SATMON031	g1001309	BLASTX	107	1e-11	53
SATMON003	g1103380	BLASTX	107	1e-12	53
SATMON025	g20598	BLASTN	1010	1e-81	89
SATMON025	g20598	BLASTN	916	1e-78	89
SATMON036	g20598	BLASTN	947	1e-74	92
SATMON007	g20598	BLASTN	1102	1e-83	89
SATMON023	g20598	BLASTN	1018	1e-80	84
SATMON010	g20598	BLASTN	1017	1e-79	88
SATMON003	g20598	BLASTN	838	1e-78	86
SATMON025	g20598	BLASTN	794	1e-76	88
SATMON017	g20598	BLASTN	897	1e-73	89
SATMON022	g20598	BLASTN	820	1e-72	86
SATMON036	g20598	BLASTN	949	1e-70	87
SATMON028	g20598	BLASTN	912	1e-67	85
SATMON036	g20598	BLASTN	782	1e-66	87
SATMON017	g435456	BLASTN	662	1e-61	86
SATMON036	g20598	BLASTN	375	1e-59	91
SATMON007	g20598	BLASTN	806	1e-58	85
SATMON033	g20598	BLASTN	729	1e-51	89
SATMON023	g20598	BLASTN	270	1e-30	87
SATMON036	g20598	BLASTN	463	1e-29	79
SATMON028	g20598	BLASTN	280	1e-27	86
SATMON033	g20598	BLASTN	407	1e-26	86
SATMON033	g20598	BLASTN	309	1e-25	87
SATMON017	g20598	BLASTN	412	1e-25	84
SATMON016	g1001309	BLASTX	96	1e-9	32
SATMON007	g633094	BLASTN	1388	1e-106	91
SATMON009	g633094	BLASTN	1329	1e-101	90
SATMON009	g633094	BLASTN	1307	1e-100	89
SATMON009	g633094	BLASTN	1314	1e-100	93
SATMON036	g633094	BLASTN	1216	1e-92	95
SATMON027	g633094	BLASTN	1154	1e-87	91

374	541	700444305H1	SATMON027	g633094	BLASTN	988	1e-86	97
375	541	700222187H1	SATMON011	g633094	BLASTN	1116	1e-84	89
376	541	700093340H1	SATMON008	g633094	BLASTN	1121	1e-84	90
377	541	700576310H1	SATMON030	g633094	BLASTN	1107	1e-83	91
378	541	700443474H1	SATMON027	g633094	BLASTN	584	1e-82	93
379	541	700440955H1	SATMON026	g633094	BLASTN	803	1e-82	92
380	541	700446111H1	SATMON027	g633094	BLASTN	939	1e-81	87
381	541	700259835H1	SATMON017	g633094	BLASTN	1073	1e-80	87
382	541	700551206H1	SATMON022	g633094	BLASTN	968	1e-76	89
383	541	700445905H1	SATMON027	g633094	BLASTN	464	1e-75	89
384	541	700446192H1	SATMON027	g633094	BLASTN	774	1e-55	92
385	541	700614693H1	SATMON033	g633094	BLASTN	600	1e-54	80
386	7402	700439746H1	SATMON026	g20596	BLASTN	1353	1e-103	97
387	7402	700621225H1	SATMON034	g20596	BLASTN	709	1e-72	97
388	7402	700456918H1	SATMON029	g20596	BLASTN	968	1e-71	95
389	7402	700453876H1	SATMON029	g20600	BLASTN	761	1e-54	96
390	7402	700623616H1	SATMON034	g20596	BLASTN	432	1e-39	96
391	7402	700454592H1	SATMON029	g20600	BLASTN	380	1e-30	81
392	7402	700454593H1	SATMON029	g20600	BLASTN	310	1e-28	96
393	7482	700197666H1	SATMON014	g2621088	BLASTX	145	1e-24	55
394	7482	700615228H1	SATMON033	g3328816	BLASTX	201	1e-20	61
395	7482	700030129H1	SATMON003	g3328816	BLASTX	178	1e-17	56
396	7482	700579227H1	SATMON031	g2621088	BLASTX	132	1e-15	44
397	786	700476002H1	SATMON025	g20598	BLASTN	1119	1e-90	92
398	786	700461103H1	SATMON033	g20598	BLASTN	1196	1e-90	91
399	786	700240702H1	SATMON010	g20598	BLASTN	1174	1e-89	91
400	786	700470851H1	SATMON025	g20598	BLASTN	1138	1e-86	91
401	786	700262654H1	SATMON017	g20598	BLASTN	1138	1e-86	91
402	786	700452647H1	SATMON028	g20598	BLASTN	1115	1e-84	88
403	786	700194349H1	SATMON014	g20598	BLASTN	1115	1e-84	92
404	786	700472225H1	SATMON025	g20598	BLASTN	645	1e-82	86
405	786	700461203H1	SATMON033	g20598	BLASTN	1019	1e-82	90
406	786	700581588H1	SATMON031	g20598	BLASTN	561	1e-79	90
407	786	700194330H1	SATMON014	g20598	BLASTN	1043	1e-78	90
408	786	700194016H1	SATMON014	g20598	BLASTN	1044	1e-78	90
409	786	700157347H1	SATMON012	g20598	BLASTN	1049	1e-78	90
410	786	700195805H1	SATMON014	g20598	BLASTN	1049	1e-78	90
411	786	700160255H1	SATMON012	g20598	BLASTN	1040	1e-77	93
412	786	700582138H1	SATMON031	g20598	BLASTN	885	1e-75	88
413	786	700197148H1	SATMON014	g20598	BLASTN	1007	1e-75	90
414	786	700159366H1	SATMON012	g20598	BLASTN	1016	1e-75	91
415	786	701184326H1	SATMONN06	g20598	BLASTN	815	1e-72	89
416	786	700159491H1	SATMON012	g20598	BLASTN	979	1e-72	93
417	786	700104663H1	SATMON010	g20598	BLASTN	966	1e-71	86
418	786	700195003H1	SATMON014	g20598	BLASTN	779	1e-69	86
419	786	700218254H1	SATMON016	g20598	BLASTN	942	1e-69	89
420	786	700802451H1	SATMON036	g20598	BLASTN	581	1e-68	90
421	786	700157772H1	SATMON012	g20598	BLASTN	887	1e-65	90
422	786	700473425H1	SATMON025	g20598	BLASTN	466	1e-64	85
423	786	700800486H1	SATMON036	g20598	BLASTN	868	1e-63	91
424	786	700185039H1	SATMON014	g20598	BLASTN	859	1e-62	86
425	786	700800057H1	SATMON036	g20598	BLASTN	567	1e-59	85
426	786	700451832H1	SATMON028	g20598	BLASTN	501	1e-58	88
427	786	700799994H1	SATMON036	g20598	BLASTN	570	1e-55	91

428	786	700801486H1	SATMON036	g20598	BLASTN	750	1e-53	91
429	786	700802086H1	SATMON036	g20598	BLASTN	459	1e-51	89
430	786	700477105H1	SATMON025	g20598	BLASTN	708	1e-50	90
431	786	700260426H1	SATMON017	g20598	BLASTN	702	1e-49	84
432	786	700799811H1	SATMON036	g20598	BLASTN	409	1e-48	84
433	786	700427005H1	SATMONN01	g20598	BLASTN	691	1e-48	89
434	786	700803487H1	SATMON036	g20598	BLASTN	423	1e-46	83
435	786	700262695H1	SATMON017	g20598	BLASTN	367	1e-43	89
436	786	700471602H1	SATMON025	g20598	BLASTN	601	1e-41	90
437	786	701185813H2	SATMONN06	g20598	BLASTN	320	1e-39	83
438	786	700196744H1	SATMON014	g20598	BLASTN	490	1e-32	92
439	786	701184204H1	SATMONN06	g20598	BLASTN	247	1e-10	78
440	786	700622453H1	SATMON034	g20598	BLASTN	230	1e-8	79
441	786	700618768H1	SATMON034	g20598	BLASTN	230	1e-8	79
442	-L30591931	LIB3059-009-Q1-K1-C12	LIB3059	g20596	BLASTN	1989	1e-157	95
443	-L30593805	LIB3059-022-Q1-K1-H6	LIB3059	g20596	BLASTN	377	1e-56	79
444	-L30596704	LIB3059-055-Q1-K1-E5	LIB3059	g20596	BLASTN	733	1e-52	89
445	-L30624957	LIB3062-040-Q1-K1-H1	LIB3062	g633095	BLASTX	112	1e-27	56
446	-L30671766	LIB3067-014-Q1-K1-B8	LIB3067	g20596	BLASTN	1132	1e-122	86
447	-L30693715	LIB3069-012-Q1-K1-F3	LIB3069	g142538	BLASTX	98	1e-24	47
448	10329	LIB3079-007-Q1-K1-B3	LIB3079	g20596	BLASTN	1201	1e-97	87
449	10329	LIB143-052-Q1-E1-E4	LIB143	g20596	BLASTN	751	1e-53	86
450	1148	LIB3078-040-Q1-K1-H1	LIB3078	g633094	BLASTN	1675	1e-130	87
451	1148	LIB3062-040-Q1-K1-H3	LIB3062	g633094	BLASTN	1310	1e-100	88
452	1148	LIB143-054-Q1-E1-F1	LIB143	g633094	BLASTN	1234	1e-94	88
453	1148	LIB83-001-Q1-E1-A10	LIB83	g633094	BLASTN	1030	1e-77	81
454	16872	LIB36-018-Q1-E1-D12	LIB36	g633094	BLASTN	542	1e-69	85
455	25099	LIB3059-012-Q1-K1-G3	LIB3059	g1001309	BLASTX	130	1e-36	38
456	319	LIB143-022-Q1-E1-G3	LIB143	g20598	BLASTN	1698	1e-135	89
457	319	LIB143-048-Q1-E1-G12	LIB143	g20598	BLASTN	1562	1e-126	87
458	319	LIB143-001-Q1-E1-H6	LIB143	g20598	BLASTN	1462	1e-113	90
459	319	LIB143-002-Q1-E1-H2	LIB143	g20598	BLASTN	484	1e-66	88
460	32047	LIB148-034-Q1-E1-F3	LIB148	g435456	BLASTN	262	1e-12	68
461	32047	LIB148-032-Q1-E1-H8	LIB148	g435456	BLASTN	255	1e-11	71
462	541	LIB3062-033-Q1-K1-G2	LIB3062	g633094	BLASTN	1706	1e-133	90
463	541	LIB3062-033-Q1-K1-G3	LIB3062	g633094	BLASTN	1123	1e-94	84
464	541	LIB3060-005-Q1-K1-C1	LIB3060	g633094	BLASTN	1061	1e-90	84
465	7402	LIB3059-004-Q1-K1-F4	LIB3059	g20596	BLASTN	1461	1e-142	92
466	7482	LIB3059-049-Q1-K1-E5	LIB3059	g2621088	BLASTX	138	1e-48	51
467	786	LIB3061-042-Q1-K1-E8	LIB3061	g20598	BLASTN	1811	1e-142	88
468	786	LIB143-040-Q1-E1-D11	LIB143	g20598	BLASTN	1462	1e-113	92
469	786	LIB143-030-Q1-E1-D9	LIB143	g20598	BLASTN	1141	1e-101	90
470	786	LIB3068-035-Q1-K1-A4	LIB3068	g20598	BLASTN	533	1e-99	78
471	786	LIB143-017-Q1-E1-C8	LIB143	g20598	BLASTN	678	1e-92	82
472	786	LIB143-030-Q1-E1-D11	LIB143	g20598	BLASTN	1165	1e-88	86
473	786	LIB3061-048-Q1-K1-D7	LIB3061	g20598	BLASTN	299	1e-15	78
474	786	LIB3059-056-Q1-K1-B1	LIB3059	g20598	BLASTN	283	1e-12	74

Seq No.	Transaminase A-soybean							
	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
475	-700668054	700668054H1	SOYMON006	g3328816	BLASTX	172	1e-16	53
476	-700685655	700685655H1	SOYMON008	g387106	BLASTX	165	1e-15	62

477	-700729138	700729138H1
478	-700734818	700734818H1
479	-700787411	700787411H2
480	-700868646	700868646H1
481	-700874369	700874369H1
482	-700974412	700974412H1
483	-701009475	701009475H1
484	-701050301	701050301H1
485	-701061267	701061267H1
486	-701129551	701129551H1
487	13413	700904367H1
488	13413	700895714H1
489	13413	700727795H1
490	13503	700974712H1
491	13503	700895483H1
492	13503	700846207H1
493	14358	700909477H1
494	14358	700732673H1
495	14358	700890192H1
496	14358	700727008H1
497	15432	700567458H1
498	15529	701045375H1
499	15529	700567374H1
500	15529	701102885H1
501	15529	701213187H1
502	15529	701055675H1
503	15529	701052631H1
504	15529	701213639H1
505	1566	700651242H1
506	1566	700661083H1
507	1566	700668434H1
508	1566	700677640H1
509	1566	700655909H1
510	1566	700660728H1
511	1566	700807523H1
512	16634	700660070H1
513	16634	700746670H1
514	1703	700749933H1
515	1703	700793749H1
516	1703	701127031H1
517	1703	700997259H1
518	1703	700670783H1
519	25132	700678487H1
520	25132	701049020H1
521	25542	701151325H1
522	25542	700964436H1
523	26671	701106241H1
524	26671	701149504H1
525	27066	700605347H2
526	27066	701053078H1
527	6297	700971234H1
528	6297	701205146H1
529	6297	701137753H1
530	6297	700741154H1

SOYMON009	g2621088	BLASTX	136	1e-17	47
SOYMON010	g3201622	BLASTX	234	1e-25	54
SOYMON011	g20598	BLASTN	908	1e-66	90
SOYMON016	g435458	BLASTN	513	1e-33	75
SOYMON018	g2654093	BLASTN	808	1e-63	90
SOYMON005	g169914	BLASTN	249	1e-11	83
SOYMON019	g1001309	BLASTX	111	1e-15	49
SOYMON032	g169914	BLASTN	263	1e-11	75
SOYMON033	g169914	BLASTN	235	1e-35	88
SOYMON037	g169914	BLASTN	1232	1e-93	93
SOYMON022	g1001121	BLASTX	231	1e-24	52
SOYMON027	g2266762	BLASTX	175	1e-22	49
SOYMON009	g1001121	BLASTX	190	1e-19	48
SOYMON005	g169914	BLASTN	1358	1e-104	99
SOYMON027	g169914	BLASTN	1236	1e-94	97
SOYMON021	g169914	BLASTN	1136	1e-85	94
SOYMON022	g710595	BLASTN	1309	1e-100	98
SOYMON010	g710595	BLASTN	1296	1e-99	98
SOYMON024	g710595	BLASTN	913	1e-83	98
SOYMON009	g710595	BLASTN	553	1e-55	99
SOYMON002	g1001309	BLASTX	115	1e-8	31
SOYMON032	g3201622	BLASTX	189	1e-19	55
SOYMON002	g3201622	BLASTX	186	1e-18	55
SOYMON028	g3201622	BLASTX	172	1e-16	56
SOYMON035	g3201622	BLASTX	174	1e-16	55
SOYMON032	g3201622	BLASTX	166	1e-15	60
SOYMON032	g3201622	BLASTX	159	1e-14	53
SOYMON035	g3201622	BLASTX	110	1e-13	59
SOYMON003	g2654093	BLASTN	1433	1e-146	98
SOYMON005	g2654093	BLASTN	898	1e-102	95
SOYMON006	g2654093	BLASTN	1289	1e-98	99
SOYMON007	g2654093	BLASTN	758	1e-97	99
SOYMON004	g2654093	BLASTN	730	1e-95	100
SOYMON005	g2654093	BLASTN	634	1e-81	90
SOYMON016	g2654093	BLASTN	478	1e-31	87
SOYMON004	g2621088	BLASTX	111	1e-20	54
SOYMON013	g2621088	BLASTX	118	1e-18	53
SOYMON013	g2654093	BLASTN	1385	1e-106	100
SOYMON017	g2654093	BLASTN	1370	1e-105	100
SOYMON037	g2654093	BLASTN	716	1e-94	96
SOYMON018	g2654093	BLASTN	1089	1e-81	97
SOYMON006	g2654093	BLASTN	767	1e-79	93
SOYMON007	g2654093	BLASTN	1175	1e-104	98
SOYMON032	g2654093	BLASTN	1260	1e-96	100
SOYMON031	g1001309	BLASTX	96	1e-15	51
SOYMON022	g1001309	BLASTX	107	1e-13	51
SOYMON036	g1001309	BLASTX	121	1e-9	39
SOYMON031	g1001309	BLASTX	122	1e-9	36
SOYMON004	g169914	BLASTN	1147	1e-104	99
SOYMON032	g169914	BLASTN	833	1e-87	96
SOYMON005	g169914	BLASTN	1303	1e-99	99
SOYMON035	g169914	BLASTN	1269	1e-96	94
SOYMON038	g169914	BLASTN	335	1e-85	93
SOYMON012	g169914	BLASTN	1135	1e-85	100

531	6297	700954813H1	SOYMON022	g169914	BLASTN	1095	1e-84	100
532	6297	701000832H1	SOYMON018	g169914	BLASTN	410	1e-83	95
533	6297	701039262H1	SOYMON029	g169914	BLASTN	650	1e-82	97
534	6297	701108365H1	SOYMON036	g169914	BLASTN	1032	1e-80	97
535	6297	700953963H1	SOYMON022	g169914	BLASTN	1058	1e-79	92
536	6297	700971364H1	SOYMON005	g169914	BLASTN	865	1e-63	95
537	6297	701002832H1	SOYMON019	g169914	BLASTN	599	1e-62	90
538	6297	700650013H1	SOYMON003	g169914	BLASTN	686	1e-61	88
539	6297	701139166H1	SOYMON038	g169914	BLASTN	632	1e-43	83
540	6297	701055975H1	SOYMON032	g169914	BLASTN	611	1e-42	99
541	6297	701131513H1	SOYMON038	g169914	BLASTN	600	1e-41	96
542	6297	701065138H1	SOYMON034	g169914	BLASTN	432	1e-38	89
543	6297	701010254H2	SOYMON019	g169914	BLASTN	427	1e-36	88
544	7549	700666429H1	SOYMON005	g169914	BLASTN	1249	1e-95	96
545	7549	701001911H1	SOYMON018	g169914	BLASTN	819	1e-59	98
546	7585	701127651H1	SOYMON037	g2654093	BLASTN	1360	1e-104	100
547	7585	700668614H1	SOYMON006	g2654093	BLASTN	1341	1e-102	99
548	7585	701054030H1	SOYMON032	g2654093	BLASTN	1341	1e-102	99
549	7585	700890128H1	SOYMON024	g2654093	BLASTN	1285	1e-98	100
550	7585	701056607H1	SOYMON032	g2654093	BLASTN	1069	1e-96	96
551	7585	700973306H1	SOYMON005	g2654093	BLASTN	1250	1e-95	100
552	7585	700845404H1	SOYMON021	g2654093	BLASTN	890	1e-94	96
553	7585	700650253H1	SOYMON003	g2654093	BLASTN	1232	1e-93	98
554	7585	700672829H1	SOYMON006	g2654093	BLASTN	1188	1e-90	99
555	7585	700664509H1	SOYMON005	g2654093	BLASTN	1074	1e-87	97
556	7585	701056892H1	SOYMON032	g2654093	BLASTN	1158	1e-87	93
557	7585	700605686H2	SOYMON005	g2654093	BLASTN	1048	1e-86	97
558	7585	700894006H1	SOYMON024	g2654093	BLASTN	1052	1e-85	96
559	7585	700955412H1	SOYMON022	g2654093	BLASTN	625	1e-84	95
560	7585	700560909H1	SOYMON001	g2654093	BLASTN	1119	1e-84	93
561	7585	700895972H1	SOYMON027	g2654093	BLASTN	1105	1e-83	100
562	7585	700663309H1	SOYMON005	g2654093	BLASTN	888	1e-82	95
563	7585	700787774H2	SOYMON011	g2654093	BLASTN	943	1e-82	96
564	7585	701069589H1	SOYMON034	g2654093	BLASTN	539	1e-81	93
565	7585	700663096H1	SOYMON005	g2654093	BLASTN	498	1e-80	95
566	7585	700836390H1	SOYMON020	g2654093	BLASTN	898	1e-80	95
567	7585	700967858H1	SOYMON033	g2654093	BLASTN	978	1e-80	92
568	7585	701101575H1	SOYMON028	g2654093	BLASTN	1032	1e-80	97
569	7585	700750565H1	SOYMON014	g2654093	BLASTN	812	1e-79	95
570	7585	701064276H1	SOYMON034	g2654093	BLASTN	820	1e-75	90
571	7585	700995223H1	SOYMON011	g2654093	BLASTN	765	1e-68	89
572	7585	700756072H1	SOYMON014	g2654093	BLASTN	899	1e-66	93
573	7585	701147945H1	SOYMON031	g2654093	BLASTN	648	1e-64	95
574	7585	700888603H1	SOYMON					

putative Transaminase A-soybean								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
582	-700999272	700999272H1	SOYMON018	g1326254	BLASTX	153	1e-15	57
583	-GM17331	LIB3055-010-Q1-N1-G4	LIB3055	g169914	BLASTN	456	1e-27	85
584	-GM25144	LIB3040-027-Q1-E1-F2	LIB3040	g2654093	BLASTN	526	1e-65	85
585	-GM41298	LIB3051-109-Q1-K1-F6	LIB3051	g2654093	BLASTN	207	1e-29	83
586	14358	LIB3051-106-Q1-K1-G8	LIB3051	g710595	BLASTN	2246	1e-178	99
587	25132	LIB3051-063-Q1-K1-D12	LIB3051	g2654093	BLASTN	1347	1e-103	96
588	32509	LIB3056-012-Q1-N1-C3	LIB3056	g2648397	BLASTX	152	1e-29	43
589	6297	LIB3055-010-Q1-N1-G6	LIB3055	g169914	BLASTN	1721	1e-134	99
590	6297	LIB3055-010-Q1-N1-G7	LIB3055	g169914	BLASTN	1246	1e-123	97
591	6297	LIB3055-010-Q1-N1-G8	LIB3055	g169914	BLASTN	1120	1e-84	93
592	6297	LIB3049-021-Q1-E1-C8	LIB3049	g169914	BLASTN	864	1e-63	91
593	7585	LIB3051-105-Q1-K1-F8	LIB3051	g2654093	BLASTN	2108	1e-167	99
594	7585	LIB3028-010-Q1-B1-C7	LIB3028	g2654093	BLASTN	1973	1e-158	97
595	7585	LIB3030-001-Q1-B1-B7	LIB3030	g2654093	BLASTN	1117	1e-138	95
596	7585	LIB3051-040-Q1-K1-D4	LIB3051	g2654093	BLASTN	1166	1e-116	94
597	9138	LIB3065-001-Q1-N1-G1	LIB3065	g152149	BLASTX	168	1e-38	52

4-hydroxyphenylpyruvate dioxygenase-maize								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
598	-700428184	700428184H1	SATMONN01	g2695709	BLASTN	773	1e-55	83
599	-700578555	700578555H1	SATMON031	g2695710	BLASTX	144	1e-12	71
600	31568	LIB143-034-Q1-E1-C6	LIB143	g2695709	BLASTN	650	1e-47	74

4-hydroxyphenylpyruvate dioxygenase-soybean								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
601	-700655923	700655923H1	SOYMON004	g2145038	BLASTN	352	1e-45	77
602	11733	700833534H1	SOYMON019	g2145039	BLASTX	124	1e-17	60
603	13818	700961605H1	SOYMON022	g2145038	BLASTN	785	1e-56	82
604	13818	700906510H1	SOYMON022	g2145038	BLASTN	744	1e-53	82
605	-GM31671	LIB3051-002-Q1-E1-A1	LIB3051	g2145038	BLASTN	668	1e-44	74
606	-GM37087	LIB3051-068-Q1-K1-H8	LIB3051	g2695709	BLASTN	593	1e-50	76
607	11733	LIB3051-067-Q1-K1-E3	LIB3051	g2145038	BLASTN	726	1e-49	74

homogentisic acid dioxygenase-maize								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
608	-700215110	700215110H1	SATMON016	g2832726	BLASTX	157	1e-26	50
609	-701185447	701185447H1	SATMONN06	g2832726	BLASTX	216	1e-28	51
610	12601	700578778H1	SATMON031	g2832726	BLASTX	307	1e-35	67
611	1732	700469334H1	SATMON025	g2832726	BLASTX	146	1e-23	52
612	1732	700469267H1	SATMON025	g2832726	BLASTX	122	1e-19	53
613	8522	700466728H1	SATMON025	g2832726	BLASTX	189	1e-19	53
614	8522	700257246H1	SATMON017	g2832726	BLASTX	182	1e-18	53
615	-L30683918	LIB3068-049-Q1-K1-D6	LIB3068	g1561616	BLASTX	158	1e-43	69

homogentisic acid dioxygenase-soybean								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
616	-700854493	700854493H1	SOYMON023	g1561616	BLASTX	113	1e-14	63
617	24903	701206316H1	SOYMON035	g2832726	BLASTX	211	1e-22	54
618	24903	701204527H2	SOYMON035	g2832726	BLASTX	205	1e-21	54
619	24903	701106917H1	SOYMON036	g2832726	BLASTX	197	1e-20	54
620	24903	701204272H2	SOYMON035	g1561616	BLASTX	80	1e-10	67
621	26239	701208301H1	SOYMON035	g2832726	BLASTX	316	1e-36	69

geranylgeranylpyrophosphate synthase-maize								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
622	-700165387	700165387H1	SATMON013	g1419758	BLASTX	119	1e-17	67
623	-700622762	700622762H1	SATMON034	g1722699	BLASTX	115	1e-10	65
624	-L30782383	LIB3078-012-Q1-K1-D3	LIB3078	g1063276	BLASTX	149	1e-46	55

geranylgeranylpyrophosphate synthase-soybean								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
625	-700741352	700741352H1	SOYMON012	g1722699	BLASTX	154	1e-15	59
626	-701098728	701098728H2	SOYMON028	g643094	BLASTX	142	1e-26	69
627	-701210428	701210428H1	SOYMON035	g558924	BLASTN	639	1e-44	78

***Table Headings**

Cluster ID

A cluster ID is arbitrarily assigned to all of those clones which belong to the same cluster at a given stringency and a particular clone will belong to only one cluster at a given stringency.

- 5 If a cluster contains only a single clone (a "singleton"), then the cluster ID number will be negative, with an absolute value equal to the clone ID number of its single member. The cluster ID entries in the table refer to the cluster with which the particular clone in each row is associated.

Clone ID

10 The clone ID number refers to the particular clone in the PhytoSeq database. Each clone ID entry in the table refers to the clone whose sequence is used for (1) the sequence comparison whose scores are presented and/or (2) assignment to the particular cluster which is presented.

Note that a clone may be included in this table even if its sequence comparison scores fail to meet the minimum standards for similarity. In such a case, the clone is included due solely to its association with a particular cluster for which sequences of one or more other member clones possess the required level of similarity.

Library

15 The library ID refers to the particular cDNA library from which a given clone is obtained. Each cDNA library is associated with the particular tissue(s), line(s) and developmental stage(s) from which it is isolated.

NCBI gi

Each sequence in the GenBank public database is arbitrarily assigned a unique NCBI gi (National Center for Biotechnology Information GenBank Identifier) number. In this table, the

NCBI gi number which is associated (in the same row) with a given clone refers to the particular GenBank sequence which is used in the sequence comparison. This entry is omitted when a clone is included solely due to its association with a particular cluster.

Method

- 5 The entry in the "Method" column of the table refers to the type of BLAST search that is used for the sequence comparison. "CLUSTER" is entered when the sequence comparison scores for a given clone fail to meet the minimum values required for significant similarity. In such cases, the clone is listed in the table solely as a result of its association with a given cluster for which sequences of one or more other member clones possess the required level of similarity.

Score

Each entry in the "Score" column of the table refers to the BLAST score that is generated by sequence comparison of the designated clone with the designated GenBank sequence using the designated BLAST method. This entry is omitted when a clone is included solely due to its association with a particular cluster. If the program used to determine the hit is HMMSW then the score refers to HMMSW score.

P-Value

The entries in the P-Value column refer to the probability that such matches occur by chance.

%Ident

- 20 The entries in the "%Ident" column of the table refer to the percentage of identically matched nucleotides (or residues) that exist along the length of that portion of the sequences which is aligned by the BLAST comparison to generate the statistical scores presented. This entry is omitted when a clone is included solely due to its association with a particular cluster.